

Pacific Journal of Mathematics

CONDITIONS FOR ISOMORPHISM IN PARTIAL DIFFERENTIAL EQUATIONS

HAROLD H. JOHNSON

CONDITIONS FOR ISOMORPHISM IN PARTIAL DIFFERENTIAL EQUATIONS

H. H. JOHNSON

This paper studies systems of linear homogeneous p. d. e. in two independent variables with constant coefficients. For such systems powerful algebraic tools are available to obtain results which may indicate patterns for more general systems. Linear isomorphism is defined, and necessary and sufficient conditions for linear isomorphism between two systems are found. This result is obtained from the infinite prolongation of the systems, and two systems are isomorphic if and only if their infinite prolongations are isomorphic. One unexpected result is the important role played by lower-order coefficients which do not appear in such classical notions as ellipticity, hyperbolicity or characteristics. The classification problem for these p. d. e. is reduced to a problem in linear algebra involving a finite number of relations.

Usually notions such as jets, exterior forms, linear bundles, etc., in C^∞ or C^ω categories are used to express our definitions. Since we are concerned here with linear objects we shall use this opportunity to state the definitions in terms of linear algebra.

Let $z = (z^1, \dots, z^m)$ be a point in the real vector space V . We denote $\partial_j z = (\partial_j z^1, \dots, \partial_j z^m)$ where $j = 1$ or 2 . Let a and b be real m by m matrices. Consider systems Σ of the form $\partial_2 z = \partial_1 z a + z b$.

The total prolongation $P\Sigma$ of Σ is the system $\partial_2 z = z_1 a + b$, $z_1 = \partial_1 z$, $\partial_2 z_1 = \partial_1 z_1 a + z_1 b$, where $z_1 = (z_1^1, \dots, z_1^m)$. Let PV denote the real vector space of $2m$ -tuples (z, z_1) . Observe that the equation $\partial_2 z_1 = \partial_1 z_1 a + z_1 b$ is dictated by 'involutiveness' from $z_1 = \partial_1 z$, since $\partial_2 z_1 = \partial_{12} z = \partial_1(\partial_1 z a + z b)$. If $z = f(x^1, x^2)$ is any solution of Σ then $(z, z_1) = (f, \partial f / \partial x^1)$ is a solution of $P\Sigma$, and every solution of $P\Sigma$ arises in this way.

Similarly, $P^2\Sigma$ is the system on $P^2V = \{(z, z_1, z_2)\}$, $\partial_2 z = z_1 a + z b$, $\partial_2 z_1 = z_2 a + z_1 b$, $\partial_2 z_2 = \partial_1 z_2 a + z_2 b$, $z_1 = \partial_1 z$, $z_2 = \partial_1 z_1$. Then $P^n\Sigma$ is defined similarly on P^nV . Finally, $P^\infty\Sigma$ is the system on $z_\infty = (z, z_1, z_2, \dots)$

$$\partial_2 z_\infty = z_\infty \begin{pmatrix} b & 0 & 0 & 0 & 0 & \dots \\ a & b & 0 & 0 & 0 & \dots \\ 0 & a & b & 0 & 0 & \dots \\ 0 & 0 & a & b & 0 & \dots \\ 0 & 0 & 0 & a & b & \dots \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix} = z_\infty P,$$

$$\partial_1 z_\infty = z_\infty \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & \dots \\ I & 0 & 0 & 0 & 0 & \dots \\ 0 & I & 0 & 0 & 0 & \dots \\ 0 & 0 & I & 0 & 0 & \dots \\ 0 & 0 & 0 & I & 0 & \dots \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix} = z_\infty Q,$$

where $a, b, 0, I$ denote m by m real matrices, 0 the zero and I the identity matrices.

We shall denote by Σ' a system $\partial_2 y = \partial_1 y c + y d$ where y is a point in the real vector space U and dimension $(U) = m'$.

DEFINITION 1. Two systems Σ and Σ' are *isomorphic* if: (1) There is a linear transformation $\pi: P^r V \rightarrow U$,

$$\pi(z, z_1, \dots, z_r) = zk + z_1 k_1 + \dots + z_r k_r,$$

where the k_j are m by m' matrices so that the 'pull-back' of the equations $\partial_2 y = \partial_1 y c + y d$ in Σ' are in $P_r \Sigma$. That is,

$$\partial_2 zk + \dots + \partial_2 z_r k_r = (\partial_1 zk + \dots \partial_1 z_r k_r) c + (zk + \dots + z_r k_r) d$$

are equations in $P^r \Sigma$. Now if $z = f(x^1, x^2)$ is any solution of Σ then $y = fk + \partial_1 f k_1 + \dots + \partial_1^r f k_r$ is a solution of Σ' .

(2) Similarly, there is a linear transformation $\rho: P^s U \rightarrow V$ so that the pull-back of $\partial_2 z = \partial_1 z a + z b$ is in $P^s \Sigma'$.

(3) Finally, we require coherence in the following sense. Letting $p^s \pi: P^{r+s} V \rightarrow P^s U$ be defined by

$$p^s \pi(z, z_1, \dots, z_{r+s}) = (zk + \dots + z_r k_r, z_1 k + \dots + z_{r+1} k_r, \dots, z_s k + \dots + z_{r+s} k_r),$$

we require that $\rho \circ p^s \pi(z, z_1, \dots, z_{r+s}) = z$. This guarantees that if $z = f(x^1, x^2)$ is a solution of Σ corresponding to the solution $y = g(x^1, x^2)$ of Σ' under π , then g will correspond to f under ρ . We also require that $\pi \circ p^r \rho(y, \dots, y_{r+s}) = y$.

It is not difficult to generalize this to nonlinear systems using jet language. One may compare isomorphism with absolute equivalence. Using [2, Prop. 1] absolutely equivalent systems can be proved to be isomorphic.

DEFINITION 2. Two systems Σ and Σ' are *infinitely isomorphic* if there exists a matrix K with countably infinite columns and rows such that

- (1) each column has all but a finite number of elements zero;
- (2) K^{-1} exists satisfying (1);
- (3) the transformation $y_\infty = z_\infty K$ transforms $P^\infty \Sigma'$ to $P^\infty \Sigma$, that is, $P^\infty \Sigma$ is the system $\partial_2 z_\infty = z_\infty K P' K^{-1}$, $\partial_1 z_\infty = z_\infty K Q' K^{-1}$ when $P^\infty \Sigma'$ is $\partial_2 y_\infty = y_\infty P'$, $\partial_1 y_\infty = y_\infty Q'$.

PROPOSITION 1. *If Σ and Σ' are isomorphic they are infinitely isomorphic.*

Proof. From $\pi: P^r V \rightarrow U$ in Definition 1 we obtain

$$p^\infty \pi: P^\infty V \rightarrow P^\infty U$$

by

$$p^\infty \pi(z_\infty) = (zk + \dots + z_r k_r, z_1 k + \dots + z_{r+1} k_r, \dots, z_i k + \dots + z_{i+r} k_r, \dots)$$

which corresponds to a matrix K with finite columns. Similarly $p^\infty \rho$ corresponds to a matrix H , and $HK = KH = I$ follows from condition (3) in Definition 1.

We now obtain an algebraic condition equivalent to infinite isomorphism.

THEOREM 1. *Two systems Σ and Σ' are infinitely isomorphic if and only if*

- (1) $m = m'$,
- (2) *there exist m by m matrices k, k_1, \dots, k_r such that*

$$\begin{aligned} bk &= kd, \\ bk_1 + ak &= kc + k_1 d, \\ bk_2 + ak_1 &= k_1 c + k_2 d, \\ &\vdots \\ bk_r + ak_{r-1} &= k_{r-1} c + k_r d, \\ ak_r &= k_r c, \end{aligned}$$

and (3) the infinite matrix

$$K = \begin{pmatrix} k & 0 & 0 & 0 & \dots & \dots \\ k_1 & k & 0 & 0 & \dots & \dots \\ \cdot & k_1 & k & 0 & \dots & \dots \\ \cdot & \cdot & k_1 & k & \dots & \dots \\ \cdot & \cdot & \cdot & k_1 & \dots & \dots \\ k_r & \cdot & \cdot & \cdot & \dots & \dots \\ 0 & k_r & \cdot & \cdot & \dots & \dots \\ 0 & 0 & k_r & \cdot & \dots & \dots \\ \cdot & \cdot & \cdot & k_r & \dots & \dots \\ \cdot & \cdot & \cdot & \cdot & \dots & \dots \\ \cdot & \cdot & \cdot & \cdot & \dots & \dots \end{pmatrix}$$

has a column-finite inverse.

Proof. Assume (1), (2) and (3). Then K defines a linear transformation between $P^\infty V$ and $P^\infty U$, which gives an infinite isomorphism, since the equations in (2) are equivalent to $PK = KP'$, and (1) implies $Q = Q'$ while it is easy to check that $QK = KQ$.

Conversely, if K is a matrix defining an isomorphism between $P^\infty \Sigma$ and $P^\infty \Sigma'$, then $PK = KP'$ and $QK = KQ'$. Let W be the real vector space of countably infinite column vectors whose components are all zero except possibly a finite number. These matrices P, Q, K can be regarded as linear transformations on W to itself.

Then K defines a linear transformation, K_0 , on $W/Q'(W)$ to $W/Q(W)$ by $K_0(w + Q'(W)) = K(w) + Q(W)$, since if $w = Q'(v)$ then $K(w) = KQ'(v) = QK(v)$. This K_0 is onto because K is onto (it has an inverse), and K_0 is one-to-one because if

$$K(w) = Q(v) = QK(v_1) = KQ'(v_1),$$

then $w = Q'(v_1)$.

Since $W/Q(W)$ is isomorphic to $W/Q'(W)$, their dimensions are the same. But the dimension of $W/Q(W)$ is m , since and vector $(v_1, v_2, \dots)^t$, where the v_i are 1 by m , may be expressed as

$$(v_1, 0, 0, \dots)^t + Q(v_2, v_3, \dots)^t.$$

Similarly for Q' . Hence $m = m'$.

From $QK = Q'K = KQ'$, if one partitions K into m by m submatrices, it is immediate that K must have the form in (3). Then relations (2) follow at once from $PK = KP'$.

THEOREM 2. *If $P^\infty \Sigma$ and $P^\infty \Sigma'$ are infinitely isomorphic, then Σ and Σ' are isomorphic.*

Proof. Let k, k_1, \dots, k_r be as in Theorem 1. Then $\pi: P^r V \rightarrow U$ defined by $\pi(z, z_1, \dots, z_r) = zk + z_1k_1 + \dots + z_rk_r$ satisfies

$$\begin{aligned} & \partial_2(zk + z_1k_1 + \dots + z_rk_r) - \partial_1(zk + \dots + z_rk_r)c \\ & \quad - (zk + \dots + z_rk_r)d \\ & \equiv (z_1a + zb)k + (z_2a + z_1b)k_1 + \dots + (\partial_1z_1a + z_r b)k_r \\ & \quad - (z_1k + z_2k_1 + \dots + z_rk_{r-1} + \partial_1z_rk_r)c - (zk + z_1k_1 \\ & \quad + \dots + z_rk_r)d \quad (\text{modulo } P^r \Sigma) \\ & = z(bk - kd) + z_1(ak + bk_1 - kc - k_1d) \\ & \quad + \dots + z_r(ak_{r-1} + bk_r - k_{r-1}c - k_rd) + \partial_1z_r(ak_r - k_rc) = 0. \end{aligned}$$

Thus condition (1) of Definition 1 is satisfied.

If $H = K^{-1}$ the analogous relations for H define $\rho: P^S U \rightarrow V$ which satisfies condition (2) of Definition 1. Finally, condition (3) follows from $HK = KH = I$.

COROLLARY. *In order that Σ and Σ' be isomorphic it is necessary that $m = m'$ and b be similar to d .*

Proof. The matrix K^{-1} exists only if k^{-1} exists, so $k^{-1}bk = d$.

This last observation shows the importance of the O -order coefficients, which is rather unexpected in light of the classical notions such as characteristics which are invariants but in which b plays no part [3, 4, 5].

Observe that if Σ and Σ' are isomorphic using the relations in Theorem 2, then from $ak_r = k_r c$ one sees that for any eigenvector v of c , $ak_r v = k_r c v = \lambda k_r v$, so $k_r v$ is an eigenvector of a belonging to the same eigenvalue provided $k_r v \neq 0$. The author conjectures that in all cases the eigenvalues of Σ and Σ' coincide.

EXAMPLE 1. If Σ is $\partial_2 z^1 = z^1$, $\partial_2 z^2 = \partial_1 z^1$ and Σ' is $\partial_2 y^1 = y^1$, $\partial_2 y^2 = 0$, then taking $r = 0$ and

$$k = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

we find Σ and Σ' isomorphic. The Cauchy problem for Σ with initial data $z^1(x^1, 0) = \varphi(x^1)$ and $z^2(x^1, 0) = \psi(x^1)$ is well-posed, for its solution is $z^1 = \varphi(x^1) \exp(x^2)$, $z^2 = \varphi'(x^1) \exp(x^2) + \psi(x^1) - \varphi'(x^1)$.

EXAMPLE 2. Now let Σ be $\partial_2 z^1 = z^2$, $\partial_2 z^2 = \partial_1 z^1$ and Σ' be

$$\partial_2 y^1 = y^2, \partial_2 y^2 = 0.$$

These systems are not isomorphic, for Σ' has solutions

$$y^1 = x^2 \psi(x^1) + \varphi(x^1), y^2 = \psi(x^1),$$

while Σ is equivalent to $\partial_{22} z^1 = \partial_1 z^1$, whose solution for $z^1(x^1, 0) = 0$, $z^2(x^1, 0) = \psi(x^1)$ requires ψ that be infinitely differentiable [1, p.27].

REFERENCES

- 1 J. Hadamard, *Lectures on Cauchy's Problem in Linear Partial Differential Equations*, Dover, New York (1952).
2. H. H. Johnson, *Absolute equivalence of exterior differential systems*, Illinois J. Math., **10** (1966), 407-411.
3. ———, *The non-invariance of hyperbolicity for partial differential equations*, Pacific J. Math., **22** (1967) 419-430.

4. ———, *Determination of hyperbolicity by partial prolongations*, Pacific J. Math., **30** (1969) 670-695.
5. ———, *Partial prolongations and characteristics of differential equations*, Trans. Amer. Math. Soc., **146** (1969), 231-240.

Received September 21, 1970.

UNIVERSITY OF WASHINGTON

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Edward Arthur Bertram, <i>Permutations as products of conjugate infinite cycles</i>	275
David Blair, <i>Almost contact manifolds with Killing structure tensors</i>	285
Bruce Donald Calvert, <i>Nonlinear equations of evolution</i>	293
Bohumil Cenkľ and Giuliano Sorani, <i>Cohomology groups associated with the $\delta\bar{\delta}$ operator</i>	351
Martin Aaron Golubitsky and Bruce Lee Rothschild, <i>Primitive subalgebras of exceptional Lie algebras</i>	371
Thomas J. Jech, <i>Two remarks on elementary embeddings of the universe</i> ...	395
Harold H. Johnson, <i>Conditions for isomorphism in partial differential equations</i>	401
Solomon Leader, <i>Measures on semilattices</i>	407
Donald Steven Passman, <i>Group rings satisfying a polynomial identity. II</i>	425
Ralph Tyrrell Rockafellar, <i>Integrals which are convex functionals. II</i>	439
Stanisław Sławomir Świerczkowski, <i>Cohomology of group germs and Lie algebras</i>	471
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable. III</i>	483
Alan Curtiss Tucker, <i>Matrix characterizations of circular-arc graphs</i>	535