

Pacific Journal of Mathematics

**EVERY GENERALIZED PETERSEN GRAPH HAS A TAIT
COLORING**

FRANK CASTAGNA AND GEERT CALEB ERNST PRINS

EVERY GENERALIZED PETERSEN GRAPH HAS A TAIT COLORING

FRANK CASTAGNA AND GEERT PRINS

Watkins has defined a family of graphs which he calls generalized Petersen graphs. He conjectures that all but the original Petersen graph have a Tait coloring, and proves the conjecture for a large number of these graphs. In this paper it is shown that the conjecture is indeed true.

DEFINITIONS. Let n and k be positive integers, $k \leq n - 1$, $n \neq 2k$. The generalized Petersen graph $G(n, k)$ has $2n$ vertices, denoted by $\{0, 1, 2, \dots, n - 1; 0', 1', 2', \dots, \dots, (n - 1)'\}$ and all edges of the form $(i, i + 1)$, (i, i') , $(i', (i + k)')$ for $0 \leq i \leq n - 1$, where all numbers are read modulo n . $G(5, 2)$ is the Petersen graph. See Watkins [2].

The sets of edges $\{(i, i + 1)\}$ and $\{(i', (i + k)')\}$ are called the outer and inner rims respectively and the edges (i, i') are called the spokes.

A Tait coloring of a trivalent graph is an edge-coloring in three colors such that each color is incident to each vertex. A 2-factor of a graph is a bivalent spanning subgraph. A 2-factor consists of disjoint circuits. A Tait cycle of a trivalent graph is a 2-factor all of whose circuits have even length. A Tait cycle induces a Tait coloring and conversely.

The method that Watkins used in proving that many generalized Petersen graphs have a Tait coloring was to prove that certain color patterns on the spokes induce a Tait coloring. Our method for the remaining cases consists of the construction of 2-factors and of proof that these 2-factors are Tait cycles under appropriate conditions.

We restrict ourselves to the generalized Petersen graphs $G(n, k)$ with the properties:

$$n \text{ odd, } n \geq 7, (n, k) = 1, \text{ and } 2 < k < \frac{n - 1}{2}.$$

All other cases (and some special instances of the above) were dealt with by Watkins.

We construct three types of 2-factors. The first type is a Tait cycle when k is odd. The second type is a Tait cycle when k is even and $n \equiv 3 \pmod{4}$ and also when k is even and $n \equiv 1 \pmod{4}$ with k^{-1} even. (As $(n, k) = 1$, we define k^{-1} as the unique positive integer $< n$, for which $kk^{-1} \equiv 1 \pmod{n}$.) The third type takes care of the remaining graphs.

The principal tool in the proofs is the automorphism φ (henceforth fixed) of $G(n, k)$ defined by $\varphi(i) = n - i$; $\varphi(i') = (n - i)$. In each case φ induces an automorphism (also called φ) of the constructed 2-factor. To facilitate notation we write $n = 2m + 1$.

CONSTRUCTION 1. The subgraph H of $G(n, k)$ has the following edges:

(a) On the outer rim: $(m + k, m + k + 1), (m + k + 1, m + k + 2), \dots, (n - 1, 0), (0, 1), (1, 2), \dots, (m - k, m - k + 1), (m - k + 2, m - k + 3), (m - k + 4, m - k + 5), \dots, (m + k - 2, m + k - 1)$.

The last line may be written as $(m - k + 2j, m - k + 2j + 1)$, $1 \leq j \leq k - 1$.

(b) Spokes: $(m + k, (m + k)'), (m - k + 1, (m - k + 1)'), (m - k + 2, (m - k + 2)'), \dots, (m + k - 1, (m + k - 1)').$

(c) On the inner rim: $(i', (i + k)'), m + 1 \leq i \leq n - 1$
 $(i', (i - k)'), k \leq i \leq m.$

EXAMPLE. $G(11, 3)$




FIGURE 1

Clearly H is a 2-factor, and $\varphi(H) = H$. If C_0 is the circuit of H which contains 0, then $\varphi(C_0) = C_0$. If C_0 has odd length, then it must contain an odd number of edges of the form $(i, -i)$ and $(i', -i')$. The only candidates are:

- (A) $(m, m + 1)$
- (B) $\left(\left(n - \frac{k}{2} \right)', \left(\frac{k}{2} \right)' \right)$
- (C) $\left(\left(\frac{n - k}{2} \right)', \left(\frac{n + k}{2} \right)' \right).$

The edge (C) is not in H by our construction. Either the presence of (A) in H or the existence of edge (B) will imply that k is even. We conclude that if k is odd C_0 has even length.

Let $m - k + 2 \leq i \leq m + k - 1$. Then either $i', i, i + 1, (i + 1)'$ or $i', i, i - 1, (i - 1)'$ are 4 consecutive vertices on a circuit of H . We call such sets 4-sets. If every point of a circuit is on a 4-set, then the circuit has even length.

Now consider a vertex $i', m + k < i \leq n - 1$ or $0 \leq i < m - k + 1$, which is not on C_0 . The circuit of H which contains i' passes consecutively through the the vertices $i', (i + k)', (i + 2k)' \dots (i + rk)', (i + (r + 1)k)'$, where $i + rk < m - k + 1, i + (r + 1)k > m - k + 1, r \geq 0$. The vertex $(i + (r + 1)k)'$ is on a 4-set, and also $i + (r + 1)k \leq m$, hence the circuit continues through the vertices $i + (r + 1)k, i + (r + 1)k \pm 1, (i + (r + 1)k \pm 1)', (i + rk \pm 1)' \dots (i \pm 1)'$. The circuit continues to $(i \pm 1 - k)'$ and by an identical argument eventually returns and hits i' or $(i + 2)'$ or $(i - 2)'$. In the first case the circuit is complete and it is easily seen that it has even length. The other two cases lead to a contradiction; for assume (w.l.o.g) that the circuit is on $(i', (i + 1)', (i + 2)')$. Then by the above argument the circuit will eventually hit either $(i + 1)'$ again or else $(i + 3)'$. But the first case is impossible, because H is bivalent. Hence the circuit contains $(i + 3)'$ and further $(i + 4)' \dots (m - k + 1)'$, but this contradicts our assumption, as $(m - k + 1)'$ is on C_0 .

CONSTRUCTION 2. H has the following edges:

- (a) On the outer rim: $(n - 1, 0), (0, 1), (2, 3), \dots, (2j, 2j + 1) \dots (n - 3, n - 2)$.
- (b) Spokes: all, except $(0, 0')$.
- (c) On the inner rim: $(0', k'), (2k', 3k'), \dots (2jk', (2j + 1)k'), \dots, ((n - 1)k', 0')$.

(For the sake of clarity we have written ck' instead of the formally more correct $(ck)'$.)

EXAMPLE. $G(15, 4)$. See Figure 2.

Again, one checks easily that H is a 2-factor and that $\varphi(H) = H$. Looking at the edges $(A), (B)$, and (C) of Construction 1, we note that (C) is not an edge if k is even. If edge (A) occurs, then $m = (n - 1)/2$ is even and $n \equiv 1 \pmod{4}$. If edge (B) occurs, and we write $k/2 \equiv jk \pmod{n}, j < n$, then j is odd by our construction. But then $k \equiv 2jk \pmod{n} \Rightarrow (2j - 1) \equiv 0 \pmod{n} \Rightarrow n = 2j - 1 \Rightarrow n \equiv 1 \pmod{4}$.

Hence if $n \equiv 3 \pmod{4}$ and k is even none of the lines $(A), (B)$, and (C) occur, and we may conclude by the argument used in Construction 1 that the circuits through 0 and $0'$ have even length. All




FIGURE 2

the points of every other circuit belong to a 4-set, and hence also have even length. Therefore H is a Tait cycle if $n \equiv 3 \pmod{4}$ and k is even.

If $n \equiv 1 \pmod{4}$ and k and k^{-1} are both even, then the edge $((k+1)', 1') = (1', (k+1)') = (k^{-1}k', (k^{-1}+1)k')$ exists in H , and so does the edge $(-1', -(k+1)')$. We then obtain the circuit:

$$\begin{aligned} &0', k', k, k+1, (k+1)', 1', 1, 0, -1, -1', \\ &-(k+1)', -(k+1), -k, -k', 0' \end{aligned}$$

which has length 14 and contains both 0 and 0'.

We conclude that in this case H is again a Tait cycle.

CONSTRUCTION 3. For this construction we assume $n \equiv 1 \pmod{4}$, k even, k^{-1} odd and $> n/2$. This last assumption is no real restriction, because if k^{-1} is odd and $< n/2$, then Construction 1 gives a Tait cycle for $G(n, k^{-1})$ and Watkins has shown that $G(n, k)$ and $G(n, k^{-1})$ are isomorphic. Finally we need to assume $k > 2$; this restriction was not needed in Constructions 1 and 2.

H has the following edges:

On the outer rim: $(-1, 0), (0, 1), (2, 3), \dots, (k-4, k-3), (k-2, k-1), (k-1, k), (k+1, k+2), \dots, (n-k-2, n-k-1), (n-k, n-k+1), (n-k+1, n-k+2), (n-k+3, n-k+4), \dots, (n-3, n-2)$.

Spokes: all except $(0'0')$, $(k-1, (k-1)'), (n-k+1, (n-k+1)').$

On the inner rim: $(0', k'), (2k', 3k'), \dots, ((n-k^{-1})k', (n-k^{-1}+1)k'), ((n-k^{-1}+1)k', (n-k^{-1}+2)k'), ((n-k^{-1}+3)k', (n-k^{-1}+4)k'), \dots, ((k^{-1}-2)k', (k^{-1}-1)k'), ((k^{-1}-1)k', k^{-1}k'), ((k^{-1}+1)k', (k^{-1}+2)k'), \dots, ((n-1)k', 0')$.

EXAMPLE. $G(17, 4)$




FIGURE 3

H is a 2-factor, as long as $n - k^{-1} + 1 < k^{-1} - 1$, which assures that the constructed edges on the inner rim cover all vertices of the inner rim. But this condition holds whenever $k^{-1} > (n + 1/2)$ or alternatively when $k^{-1} > (n/2)$, and $k > 2$. It is clear that $\varphi(H) = H$.

Since $n \equiv 1 \pmod{4}$, m is even and $(m, m + 1)$ is not an edge of H . As $(n - k)/2$ is not an integer H does not have an edge $((n - k)/2)', (n + k)/2)'$. Finally, since $n - k^{-1} + 1 \leq (n - 1)/2 = m < m + 1 = (n + 1)/2 \leq k^{-1} - 1$, and m is even, H does not contain the edge $(mk', (m + 1)k') = (-k'/2, k'/2)$. As before we conclude that the circuits containing 0 and 0' have even length. The circuit containing 0 also contains $n - 1, (n - 1)', (k - 1)'$ and $1, 1', (n - k + 1)'$, while the circuit containing 0' also contains $k', k, k - 1, k - 2, (k - 2)'$ and $(n - k)', n - k, n - k + 1, n - k + 2, (n - k + 2)'$. Hence the other circuits only contain vertices of 4-sets and every circuit of H has even length.

We note that our constructions are not mutually exclusive. For example, Construction 1 also produces a Tait cycle, when k is even, and the largest positive integer q such that $qk < n$ is an odd number.

We conclude with a new conjecture. G.N. Robertson [1] has shown that $G(n, 2)$ is Hamiltonian unless $n \equiv 5 \pmod{6}$. As $G(n, 2) \cong G(n, (n + 1)/2) \cong G(n, (n - 1)/2) \cong G(n, n - 2)$ (see [2]), none of these graphs has a Hamiltonian if $n \equiv 5 \pmod{6}$. We conjecture that all other generalized Petersen graphs are Hamiltonian. In all examples that we have worked out $G(n, k)$ possesses a Hamiltonian H with $\varphi(H) = H$, but our three constructions are Hamiltonians only in a minority of cases.

REFERENCES

1. G. N. Robertson, *Graphs under Girth, Valency, and Connectivity Constraints* (Dissertation), University of Waterloo, Waterloo, Ontario, Canada, 1968.
2. Mark E. Watkins, *A theorem on Tait colorings with an application to the generalized Petersen graphs*, *J. Combinatorial Theory*, **6** (1969), 152-164.

Received June 16, 1970 and in revised form August 20, 1970.

WAYNE STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 40, No. 1

September, 1972

Alex Bacopoulos and Athanassios G. Kartsatos, <i>On polynomials approximating the solutions of nonlinear differential equations</i>	1
Monte Boisen and Max Dean Larsen, <i>Prüfer and valuation rings with zero divisors</i>	7
James J. Bowe, <i>Neat homomorphisms</i>	13
David W. Boyd and Hershy Kisilevsky, <i>The Diophantine equation $u(u + 1)(u + 2)(u + 3) = v(v + 1)(v + 2)$</i>	23
George Ulrich Brauer, <i>Summability and Fourier analysis</i>	33
Robin B. S. Brooks, <i>On removing coincidences of two maps when only one, rather than both, of them may be deformed by a homotopy</i>	45
Frank Castagna and Geert Caleb Ernst Prins, <i>Every generalized Petersen graph has a Tait coloring</i>	53
Micheal Neal Dyer, <i>Rational homology and Whitehead products</i>	59
John Fuelberth and Mark Lawrence Teply, <i>The singular submodule of a finitely generated module splits off</i>	73
Robert Gold, <i>Γ-extensions of imaginary quadratic fields</i>	83
Myron Goldberg and John W. Moon, <i>Cycles in k-strong tournaments</i>	89
Darald Joe Hartfiel and J. W. Spellmann, <i>Diagonal similarity of irreducible matrices to row stochastic matrices</i>	97
Wayland M. Hubbard, <i>Some results on blocks over local fields</i>	101
Alan Loeb Kostinsky, <i>Projective lattices and bounded homomorphisms</i>	111
Kenneth O. Leland, <i>Maximum modulus theorems for algebras of operator valued functions</i>	121
Jerome Irving Malitz and William Nelson Reinhardt, <i>Maximal models in the language with quantifier "there exist uncountably many"</i>	139
John Douglas Moore, <i>Isometric immersions of space forms in space forms</i>	157
Ronald C. Mullin and Ralph Gordon Stanton, <i>A map-theoretic approach to Davenport-Schinzel sequences</i>	167
Chull Park, <i>On Fredholm transformations in Yeh-Wiener space</i>	173
Stanley Poreda, <i>Complex Chebyshev alterations</i>	197
Ray C. Shiflett, <i>Extreme Markov operators and the orbits of Ryff</i>	201
Robert L. Snider, <i>Lattices of radicals</i>	207
Ralph Richard Summerhill, <i>Unknotting cones in the topological category</i>	221
Charles Irvin Vinsonhaler, <i>A note on two generalizations of QF - 3</i>	229
William Patterson Wardlaw, <i>Defining relations for certain integrally parameterized Chevalley groups</i>	235
William Jennings Wickless, <i>Abelian groups which admit only nilpotent multiplications</i>	251