

# Pacific Journal of Mathematics

**ON FREDHOLM TRANSFORMATIONS IN YEH-WIENER  
SPACE**

CHULL PARK

# ON FREDHOLM TRANSFORMATIONS IN YEH-WIENER SPACE

CHULL PARK

**Let  $C_Y$  denote the Yeh-Wiener space, i.e., the space of all real-valued continuous functions  $f(x, y)$  on  $I^2 \equiv [0, 1] \times [0, 1]$  such that  $f(0, y) = f(x, 0) \equiv 0$ . Yeh has defined a Gaussian probability measure on  $C_Y$  such that the mean of the process**

$$m(x, y) \equiv \int_{C_Y} f(x, y) d_Y f = 0$$

**and the covariance**

$$R(s, t, x, y) \equiv \int_{C_Y} f(s, t) f(x, y) d_Y f = (1/2) \min(s, x) \min(t, y).$$

**Consider now a linear transformation of  $C_Y$  onto  $C_Y$  of the form**

$$(1.1) \quad \begin{aligned} T: f(x, y) &\rightarrow g(x, y) \\ &= f(x, y) + \int_{I^2} K(x, y, s, t) f(s, t) ds dt, \end{aligned}$$

**which is often called a Fredholm transformation. The main purpose of this paper is to find the corresponding Radon-Nikodym derivative thus showing how the Yeh-Wiener integrals transform under the transformation.**

The transformations considered here contain the Volterra transformation

$$T_1[f(x, y)] = f(x, y) + \int_0^y \int_0^x K(x, y, s, t) f(s, t) ds dt$$

as a special case.

Such transformations in Wiener space have been studied a great deal by Cameron and Martin [1], Woodward [9], Segal [5], [6], and Shepp [7], and the results have proved very useful in the evaluation of various Wiener integrals.

The transformation theorems in this paper are based on stochastic integrals called the generalized Paley-Wiener-Zygmund (P.W.Z.) integrals given in [3] and [4]. For a function  $h(x, y) \in L^2(I^2)$  and  $f(x, y) \in C_Y$ , the generalized P.W.Z. integral is defined to be

$$(1.2) \quad \int_{I^2} h f d^* f \equiv \lim_{n \rightarrow \infty} \int_{I^2} (h f)_n df,$$

where  $(hf)_n$  is the  $n^{\text{th}}$  partial sum of the Fourier expansion of  $h(x, y)f(x, y)$  with respect to a C.O.N. set belonging to a class of C.O.N. systems  $\{\alpha_k(x, y)\}$  with each  $\alpha_k(x, y)$  of B.V. satisfying the condition

$$\lim_{n \rightarrow \infty} \sum_{j=1}^n \alpha_j(x, y) \int_0^y \int_0^x \alpha_j(s, t) ds dt = \frac{1}{4},$$

and the limit and on the right of (1.2) is an ordinary Riemann-Stieltjes (R-S) integral. It is known that the limit in (1.2) exists for almost all  $f$  in  $C_Y$  and it is essentially independent of the particular choice of the C.O.N. set in the class. (For details see [3] and [4].)

The Fredholm determinant  $D(K)$  of  $K(x, y, s, t) \in L^2(I^4)$  for  $\lambda = -1$  is defined by

$$D(K) \equiv 1$$

$$+ \sum_{k=1}^{\infty} \frac{1}{k!} \int_{I^{2k}} \begin{vmatrix} K(x_1, y_1, x_1, y_1) \cdots K(x_1, y_1, x_k, y_k) \\ \cdots \cdots \cdots \cdots \cdots \cdots \\ K(x_k, y_k, x_1, y_1) \cdots K(x_k, y_k, x_k, y_k) \end{vmatrix} dx_1 dy_1 \cdots dx_k dy_k.$$

## 2. Statement of main results.

**THEOREM I.** Suppose that each  $K_i(x, y, s, t)$ ,  $i = 1, 2, 3, 4$ , is continuous on  $I^4$  and absolutely continuous in  $x, y$  for each  $(s, t) \in I^2$  and  $K_1(0, y, s, t) = K_1(x, 0, s, t) = K_3(x, 0, s, t) = K_4(0, y, s, t) \equiv 0$ . Let  $K(x, y, s, t)$  be defined on  $I^4$  by

$$(2.1) \quad K(x, y, s, t) = \begin{cases} K_1(x, y, s, t) & \text{if } x < s, y < t \\ K_2(x, y, s, t) & \text{if } x > s, y > t \\ K_3(x, y, s, t) & \text{if } x > s, y < t \\ K_4(x, y, s, t) & \text{if } x < s, y > t \\ 2^{-1}(K_1 + K_3)(x, y, x, t) & \text{if } x = s, y < t \\ 2^{-1}(K_2 + K_4)(x, y, x, t) & \text{if } x = s, y > t \\ 2^{-1}(K_1 + K_4)(x, y, s, y) & \text{if } x < s, y = t \\ 2^{-1}(K_2 + K_3)(x, y, s, y) & \text{if } x > s, y = t \\ 4^{-1}(K_1 + K_2 + K_3 + K_4)(x, y, x, y) & \text{if } x = s, y = t, \end{cases}$$

where  $(K_1 + K_3)(x, y, x, t) = K_1(x, y, x, t) + K_3(x, y, x, t)$ , etc., and let

$$(2.2) \quad \begin{aligned} \phi(x, s, t) &\equiv K(x, t^+, s, t) - K(x, t^-, s, t) \\ \psi(y, s, t) &\equiv K(s^+, y, s, t) - K(s^-, y, s, t). \end{aligned}$$

To be definite at each jump discontinuity, let us agree that the partials take left-hand limit whenever it fails to exist at a point, i.e.,

$$(2.3) \quad \begin{aligned} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) &\equiv H(x, y, s, t) = \lim_{(u, v) \rightarrow (x^-, y^-)} \frac{\partial^2}{\partial v \partial u} K(u, v, s, t) \\ \frac{\partial}{\partial x} \phi(x, s, t) &\equiv A(x, s, t) = \lim_{u \rightarrow x^-} \frac{\partial}{\partial u} \phi(u, s, t), \\ \frac{\partial}{\partial y} \psi(y, s, t) &\equiv B(y, s, t) = \lim_{v \rightarrow y^-} \frac{\partial}{\partial v} \psi(v, s, t), \end{aligned}$$

and assume that there exists an integrable function  $M(s, t)$  such that for all  $(s, t) \in I^2$

$$(2.4) \quad \left. \begin{array}{l} \sup_{(x, y) \in I^2} |H(x, y, s, t)| \\ \text{var}_{(x, y) \in I^2} H(x, y, s, t) \\ \text{var}_{x \in I} H(x, 1, s, t) \\ \text{var}_{y \in I} H(1, y, s, t) \\ \text{var}_{x \in I} A(x, s, t) \\ \text{var}_{y \in I} B(y, s, t) \end{array} \right\} \leq M(s, t).$$

Also assume that

$$(2.5) \quad |A(x, s, t)|, |B(y, s, t)| \leq \beta, \text{ a constant, and}$$

$$(2.6) \quad D(K) \neq 0.$$

Then for any Yeh-Wiener measurable functional  $F(f)$ , we have under the Fredholm transformation (1.1)

$$(2.7) \quad \int_{C_Y} F(g) d_Y g \stackrel{*}{=} |D(K)| \int_{C_Y} F(Tf) \cdot \exp\{-\Phi(f)\} d_Y f,$$

where

$$(2.8) \quad \begin{aligned} \Phi(f) &= \int_{I^2} \left[ \frac{\partial^2}{\partial y \partial x} \int_{I^2} K(x, y, s, t) f(s, t) ds dt \right]^2 dx dy \\ &+ 2 \int_{I^2} \left[ \frac{\partial^2}{\partial y \partial x} \int_{I^2} K(x, y, s, t) f(s, t) ds dt \right] d^* f(x, y), \end{aligned}$$

and “ $\stackrel{*}{=}$ ” indicates the existence of one side implies that of the other and the equality.

**THEOREM II.** Let  $h(x, y) \in L^2$  on  $I^2$ ,  $K(x, y, s, t)$  and  $F(f)$  as in Theorem I. Then under the (nonlinear) transformation

$$(2.9) \quad \begin{aligned} L: f(x, y) \rightarrow g(x, y) &= f(x, y) + f_0(x, y) \\ &+ \int_{I^2} K(x, y, x, s, t) f(s, t) ds dt, \end{aligned}$$

where  $f_0(x, y) = \int_0^y \int_0^x h(s, t) ds dt$ , we have

$$\int_{C_Y} F(g) d_Y g \stackrel{*}{=} |D(K)| \int_{C_Y} F(Lf) \cdot \exp\{-\Psi(f)\} d_Y f ,$$

where

$$\begin{aligned} \Psi(f) &= \int_{I^2} \left[ h(x, y) + \frac{\partial^2}{\partial y \partial x} \int_{I^2} K(x, y, s, t) f(s, t) ds dt \right]^2 dx dy \\ &\quad + 2 \int_{I^2} \left[ h(x, y) + \frac{\partial^2}{\partial y \partial x} \int_{I^2} K(x, y, s, t) f(s, t) ds dt \right] d^* f(x, y) . \end{aligned}$$

**3. Definitions.** Corresponding to each continuous function  $f(x, y)$  on  $I^2$ , the ( $n^{\text{th}}$ ) quasi-polyhedric function  $f_{(n)}(x, y)$  of  $f(x, y)$  is defined to be

$$(3.1) \quad f_{(n)}(x, y) = a_{ij}xy + b_{ij}x + c_{ij}y + d_{ij}$$

on each square  $Q_{ij} \equiv [(i-1)/n, i/n] \times [(j-1)/n, j/n]$ ,  $i, j = 1, 2, \dots, n$ , where  $a_{ij}, b_{ij}, c_{ij}$ , and  $d_{ij}$  are so chosen that  $f_{(n)}(x, y)$  and  $f(x, y)$  agree at the vertices  $(i/n, j/n), (i/n, (j-1)/n), ((i-1)/n, j/n)$  and  $((i-1)/n, (j-1)/n)$ .

**REMARKS.** (i). Since the function  $f_{(n)}(x, y)$  is linear horizontally and vertically in each square  $Q_{ij}$ , we see that  $f_{(n)}(x, y)$  is continuous on  $I^2$ . Furthermore the sequence  $\{f_{(n)}(x, y)\}$  converges to  $f(x, y)$  uniformly on  $I^2$  as  $n \rightarrow \infty$ . Evaluating  $a_{ij}, b_{ij}, c_{ij}$ , and  $d_{ij}$  explicitly, and then combining terms, we have on each square  $Q_{ij}$ ,

$$\begin{aligned} (3.2) \quad f_{(n)}(x, y) &= f\left(\frac{i}{n}, \frac{j}{n}\right)[n^2xy - n(j-1)x - n(i-1)y + (i-1)(j-1)] \\ &\quad + f\left(\frac{i-1}{n}, \frac{j}{n}\right)[-n^2xy + n(j-1)x + niy - i(j-1)] \\ &\quad + f\left(\frac{i}{n}, \frac{j-1}{n}\right)[-n^2xy + njx + n(i-1)y - j(i-1)] \\ &\quad + f\left(\frac{i-1}{n}, \frac{j-1}{n}\right)[n^2xy - njx - niy + ij] . \end{aligned}$$

(ii) If  $K(x, y, s, t)$  is continuous on  $I^4$ , then for each  $(s, t) \in I^2$  we can think of  $K(x, y, s, t)$  as a function in  $x, y$  and so we have the ( $n^{\text{th}}$ ) quasi-polyhedric function  $K_{(n)}(x, y, s, t)$  in  $x, y$ , namely for  $(x, y) \in Q_{ij}; i, j = 1, 2, \dots, n$ , we have

$$\begin{aligned}
(3.3) \quad & K_{(n)}(x, y, s, t) \\
& = K\left(\frac{i}{n}, \frac{j}{n}, s, t\right)[n^2xy - n(j-1)x - n(i-1)y + (i-1)(j-1)] \\
& \quad + K\left(\frac{i-1}{n}, \frac{j}{n}, s, t\right)[-n^2xy + n(j-1)x + niy - i(j-1)] \\
& \quad + K\left(\frac{i}{n}, \frac{j-1}{n}, s, t\right)[-n^2xy + njx + n(i-1)y - j(i-1)] \\
& \quad + K\left(\frac{i-1}{n}, \frac{j-1}{n}, s, t\right)[n^2xy - njx - niy + ij] .
\end{aligned}$$

We also see that  $K_{(n)}(x, y, s, t) \rightharpoonup K(x, y, s, t)$  on  $I^4$  as  $n \rightarrow \infty$ , provided that  $K(x, y, s, t)$  is continuous on  $I^4$ . Here “ $\rightharpoonup$ ” indicates uniform convergence.

For each  $\varepsilon > 0$  let

$$\begin{aligned}
(3.4) \quad & \Omega_\varepsilon(s) = \operatorname{sgn} s - s/\varepsilon \quad \text{if } |s| \leq \varepsilon \\
& = 0 \quad \quad \quad \text{if } |s| > \varepsilon .
\end{aligned}$$

Let  $K(x, y, s, t)$  and  $\psi(y, s, t)$  be as in Theorem I. Then the function defined by

$$(3.5) \quad L_\varepsilon(x, y, s, t) \equiv K(x, y, s, t) + 2^{-1}[\Omega_\varepsilon(s-x) - \Omega_\varepsilon(s^+)]\psi(y, s, t)$$

is continuous in  $x, s$ . Now define

$$(3.6) \quad J(x, s, t) \equiv L_\varepsilon(x, t^+, s, t) - L_\varepsilon(x, t^-, s, t) ,$$

and

$$\begin{aligned}
(3.7) \quad & K_\varepsilon(x, y, s, t) \equiv L_\varepsilon(x, y, s, t) + 2^{-1}[\Omega_\varepsilon(t-y) - \Omega_\varepsilon(t^+)]J(x, s, t) \\
& = K(x, y, s, t) + 2^{-1}[\Omega_\varepsilon(s-x)\Omega_\varepsilon(s^+)]\psi(y, s, t) \\
& \quad + 2^{-1}[\Omega_\varepsilon(t-y) - \Omega_\varepsilon(t^+)]\phi(x, s, t) \\
& \quad + 4^{-1}[\Omega_\varepsilon(s-x) - \Omega_\varepsilon(s^+)][\Omega_\varepsilon(t-y) - \Omega_\varepsilon(t^+)](K_1 + K_2 \\
& \quad - K_3 - K_4)(s, t, s, t) .
\end{aligned}$$

Then  $K_\varepsilon(x, y, s, t)$  is continuous on  $I^4$ , and  $K_\varepsilon(0, y, s, t) = K_\varepsilon(x, 0, s, t) \equiv 0$ . Furthermore  $K_\varepsilon(x, y, s, t)$  is uniformly bounded in  $\varepsilon, x, y, s, t$ , and  $\lim_{\varepsilon \rightarrow 0^+} K_\varepsilon(x, y, s, t) = K(x, y, s, t)$ . Now, define

$$\begin{aligned}
(3.8) \quad & C_\varepsilon(x, s) = \varepsilon/2 \quad \text{if } -\varepsilon < s - x \leq \varepsilon \\
& = 0 \quad \text{otherwise} .
\end{aligned}$$

Then from (3.7), (3.4), (3.8), and (2.3) it follows

$$\begin{aligned}
(3.9) \quad H_\varepsilon(x, y, s, t) &\equiv \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) \\
&= H(x, y, s, t) + C_\varepsilon(x, s)B(y, s, t) + C_\varepsilon(y, t)A(x, s, t) \\
&\quad + C_\varepsilon(x, s) \cdot C_\varepsilon(y, t) \cdot (K_1 + K_2 - K_3 - K_4)(s, t, s, t),
\end{aligned}$$

with the understanding that whenever the partial derivatives are not defined at a point, then the value at the point to be the left-hand limit with respect to  $x$  and  $y$  (for the uniqueness sake) as in (2.3). Thus by using the mean-value property and the dominated convergence, we see that for almost all  $(x, y)$  in  $I^2$

$$\begin{aligned}
(3.10) \quad &\lim_{\varepsilon \rightarrow 0+} \int_{I^2} H_\varepsilon(x, y, s, t) f(s, t) ds dt \\
&= \int_{I^2} H(x, y, s, t) f(s, t) ds dt + \int_0^1 B(y, x, t) f(x, t) dt \\
&\quad + \int_0^1 A(x, s, y) f(s, y) ds + (K_1 + K_2 - K_3 - K_4)(x, y, x, y) \cdot f(x, y).
\end{aligned}$$

But the right hand side of (3.10) is exactly equal to

$$(\partial^2 / \partial y \partial x) \int_{I^2} K(x, y, s, t) f(s, t) ds dt.$$

Therefore for a.a.  $(x, y)$  in  $I^2$

$$\begin{aligned}
(3.11) \quad &\lim_{\varepsilon \rightarrow 0+} \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) f(s, t) ds dt \\
&= \frac{\partial^2}{\partial y \partial x} \int_{I^2} K(x, y, s, t) f(s, t) ds dt.
\end{aligned}$$

#### 4. Some Preliminary Lemmas.

**LEMMA 1.** Let  $K_{(n)}(x, y, s, t)$  be the  $n$ -th polyhedric function in  $(x, y)$  with the understanding that  $K_{(n)}(x, y, s, t) = 0$  for all  $(x, y, s, t) \notin I^4$ . For  $i, j, p, q = 1, 2, \dots, n$  let

$$(4.1) \quad A_{ijpq}^{(n)} \equiv \int_{(q-1)/n}^{(q+1)/n} \int_{(p-1)/n}^{(p+1)/n} K_{(n)}\left(\frac{i}{n}, \frac{j}{n}, s, t\right) (1 - |ns - p|)(1 - |nt - q|) ds dt.$$

Then for any  $f \in C_Y$  we obtain:

$$(i) \quad \int_{I^2} K_{(n)}\left(\frac{i}{n}, \frac{j}{n}, s, t\right) f_{(n)}(s, t) ds dt = \sum_{p,q=1}^n A_{ijpq}^{(n)} f\left(\frac{p}{n}, \frac{q}{n}\right),$$

$$(ii) \quad \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{(n)}(x, y, s, t) f_{(n)}(s, t) ds dt = n^2 \sum_{p,q=1}^n B_{ijpq}^{(n)} f\left(\frac{p}{n}, \frac{q}{n}\right)$$

for  $(x, y) \in \left(\frac{i-1}{n}, \frac{i}{n}\right) \times \left(\frac{j-1}{n}, \frac{j}{n}\right)$ , where

$$\begin{aligned} B_{ijpq}^{(n)} &\equiv A_{ijpq}^{(n)} - A_{i-1,jpq}^{(n)} - A_{i,j-1,pq}^{(n)} + A_{i-1,j-1,pq}^{(n)}, \\ \text{(iii)} \quad & \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{(n)}(x, y, s, t) f_{(n)}(s, t) ds dt \right]^2 dx dy \\ &= n^2 \sum_{i,j=1}^n \left[ \sum_{p,q=1}^n B_{ijpq}^{(n)} f\left(\frac{p}{n}, \frac{q}{n}\right) \right]^2, \\ \text{(iv)} \quad & \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{(n)}(x, y, s, t) f_{(n)}(s, t) ds dt \right] d f_{(n)}(x, y) \\ &= n^2 \sum_{i,j=1}^n \left[ \sum_{p,q=1}^n B_{ijpq}^{(n)} f\left(\frac{p}{n}, \frac{q}{n}\right) \right] \mathcal{A}_{ij} f_{(n)}, \end{aligned}$$

where

$$\begin{aligned} \mathcal{A}_{ij} f_{(n)} &\equiv f\left(\frac{i}{n}, \frac{j}{n}\right) - f\left(\frac{i-1}{n}, \frac{j}{n}\right) - f\left(\frac{i}{n}, \frac{j-1}{n}\right) \\ &\quad + f\left(\frac{i-1}{n}, \frac{j-1}{n}\right). \end{aligned}$$

The proof of this lemma is similar to that of corresponding results in [1]. Next, we consider a transformation of  $C_Y$  to  $C_Y$ :

$$T: g(x, y) = f_{(n)}(x, y) + \int_{I^2} K_{(n)}(x, y, s, t) f_{(n)}(s, t) ds dt.$$

Then by (i) of Lemma 1 and the fact that  $f_{(n)}(i/n, j/n) = f(i/n, j/n)$  at each  $i$  and  $j$ , we have

$$(4.2) \quad T: g\left(\frac{i}{n}, \frac{j}{n}\right) = f\left(\frac{i}{n}, \frac{j}{n}\right) + \sum_{p,q=1}^n A_{ijpq}^{(n)} f\left(\frac{p}{n}, \frac{q}{n}\right); \quad i, j = 1, 2, \dots, n.$$

The determinant  $\mathcal{A}(K_{(n)})$  of this transformation is given by

$$(4.3) \quad \mathcal{A}(K_{(n)}) \equiv \det(A_{IP}^* + \delta_{IP})_{I,P} = 1, 2, \dots, n^2$$

where  $A_{IP}^* = A_{ijpq}^{(n)}$  with  $I \equiv (i-1)n+j$ ,  $P \equiv (p-1)n+q$ ,  $1 \leq i, j, p, q \leq n$ .

**LEMMA 2.** *Let  $F(f)$  be a Yeh-Wiener measurable functional which depends only on the  $n^2$  values of  $f(x, y)$  at  $(x, y) = (i/n, j/n)$ ;  $i, j = 1, 2, \dots, n$ . Let the  $n^{\text{th}}$  quasi-polyhedric function in  $x, y, K_{(n)}(x, y, s, t)$ , satisfy that  $K_{(n)}(x, y, s, t) = 0$  if  $x = 0$  or  $y = 0$ , and that  $\mathcal{A}(K_{(n)}) \neq 0$ . Then*

$$\begin{aligned}
\int_{C_Y} F(g) d_Y g &\stackrel{*}{=} |\mathcal{A}(K_{(n)})| \int_{C_Y} F[f_{(n)} + \int_{I^2} K_{(n)}(\cdot, \cdot, s, t) f_{(n)}(s, t) ds dt] \\
&\quad \exp \left\{ - \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{(n)}(x, y, s, t) f_{(n)}(s, t) ds dt \right]^2 dx dy \right. \\
&\quad \left. - 2 \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{(n)}(x, y, s, t) f_{(n)}(s, t) ds dt \right] df_{(n)}(x, y) \right\} d_Y f .
\end{aligned}$$

In view of Lemma 1 and a Yeh-Wiener integral formula (see Theorem I, [11]) the proof is word for word identical to that of the corresponding lemma in [1].

**LEMMA 3.** If  $K(x, y, s, t)$  is continuous on  $I^4$ , and if the Fredholm determinant  $D(K) \neq 0$ , then

$$\lim_{n \rightarrow \infty} \mathcal{A}(K_{(n)}) = D(K) .$$

*Proof.* Using (4.3), we may expand  $\mathcal{A}(K_{(n)})$  as:

$$\begin{aligned}
(4.4) \quad \mathcal{A}(K_{(n)}) &= \det (A_{IP}^* + \delta_{IP})_{I, P=1, 2, \dots, n^2} \\
&= 1 + \sum_{P=1}^{n^2} A_{PP}^* + \frac{1}{2!} \sum_{P_1, P_2=1}^{n^2} \det (A_{P_i P_j}^*)_{i, j=1, 2} \\
&\quad + \dots + \frac{1}{(n^2)!} \sum_{P_1, \dots, P_{n^2}=1}^{n^2} \det (A_{P_i P_j}^*)_{i, j=1, \dots, n^2} \\
&= 1 + \sum_{p, q=1}^n A_{pqpq}^{(n)} + \frac{1}{2!} \sum_{p_1, q_1, p_2, q_2=1}^n \det (A_{p_i q_i p_j q_j}^{(n)})_{i, j=1, 2} \\
&\quad + \dots + \frac{1}{(n^2)!} \sum_{p_1, q_1, p_2, q_2, \dots, p_{n^2}, q_{n^2}=1}^n \det (A_{p_i q_i p_j q_j}^{(n)})_{i, j=1, 2, \dots, n^2} .
\end{aligned}$$

Let  $M$  be the bound for  $K$  on  $I^4$ . Then by (4.1) we have  $|n^2 A_{ijpq}^{(n)}| \leq M$  uniformly in  $n, i, j, p, q$ . Thus by Hadamard's inequality it follows:

$$(4.5) \quad |\det (A_{p_i q_i p_j q_j}^{(n)})_{i, j=1, \dots, N}| \leq (M/n^2)^N N^{N/2} .$$

Upon using (4.5) in (4.4), we conclude that  $|\mathcal{A}(K_{(n)})|$  is uniformly bounded by the convergent series  $1 + \sum_{N=1}^{\infty} M^N N^{N/2}/N!$ , and for each  $N$

$$\begin{aligned}
&\lim_{n \rightarrow \infty} \sum_{p_1, q_1, \dots, p_N, q_N=1}^n \det (A_{p_i q_i p_j q_j}^{(n)})_{i, j=1, \dots, N} \\
&= \int_{I^{2N}} \begin{vmatrix} K(s_1, t_1, s_1, t_1) \cdots K(s_1, t_1, s_N, t_N) \\ \cdots \cdots \cdots \cdots \cdots \cdots \\ K(s_N, t_N, s_1, t_1) \cdots K(s_N, t_N, s_N, t_N) \end{vmatrix} ds_1 dt_1 \cdots ds_N dt_N .
\end{aligned}$$

Hence the conclusion follows.

Similarly it follows:

LEMMA 4. Let  $K(x, y, s, t)$  be a bounded integrable function with  $D(K) \neq 0$  and let  $\{K_\lambda(x, y, s, t)\}$  be a set of Borel measurable functions which are uniformly bounded in  $\lambda, x, y, s, t$ , and let  $\lim_{\lambda \rightarrow 0^+} K_\lambda(x, y, s, t) = K(x, y, s, t)$ . Then  $\lim_{\lambda \rightarrow 0^+} D(K_\lambda) = D(K)$ .

LEMMA 5. For each  $(s, t) \in I^2$  let  $H(x, y, s, t)$  satisfy that

$$\sup_{(x,y) \in I^2} |H(x, y, s, t)|, \quad \text{var}_{(x,y) \in I^2} H(x, y, s, t), \quad \text{var}_{x \in I} H(x, 1, s, t), \quad \text{var}_{y \in I} H(1, y, s, t)$$

are all dominated by an integrable function  $M(s, t)$ . Then

$$(4.6) \quad \begin{aligned} & \int_{I^2} \left[ \int_{I^2} H(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\ &= \int_{I^2} f(s, t) \left[ \int_{I^2} H(x, y, s, t) df(x, y) ds dt \right]. \end{aligned}$$

*Proof.* Let “ $\|\cdot\|$ ” denote the supremum of the absolute value. Then from the fact that

$$(4.7) \quad \left. \begin{aligned} & \text{var}_{(x,y) \in I^2} \int_{I^2} H(x, y, s, t) f(s, t) ds dt \\ & \text{var}_{y \in I} \int_{I^2} H(x, 1, s, t) f(s, t) ds dt \\ & \text{var}_{y \in I} \int_{I^2} H(1, y, s, t) f(s, t) ds dt \end{aligned} \right\} \leq \|f\| \int_{I^2} M(s, t) ds dt < \infty,$$

the left member of (4.6) exists as an ordinary  $R-S$  integral. Hence for the net:  $x_i = i/n, y_j = j/n; i, j = 1, 2, \dots, n$ , and  $x_{i-1} \leqq x_i^* \leqq x_i$ ,  $y_{j-1} \leqq y_j^* \leqq y_j$ , we have with  $\Delta_{ij}f(x, y) = f(x_i, y_j) - f(x_{i-1}, y_j) - f(x_i, y_{j-1}) + f(x_{i-1}, y_{j-1})$ ,

$$(4.8) \quad \begin{aligned} & \int_{I^2} \left[ \int_{I^2} H(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\ &= \lim_{n \rightarrow \infty} \sum_{i,j=1}^n \left[ \int_I H(x_i^*, y_j^*, s, t) f(s, t) ds dt \right] \Delta_{ij}f(x, y) \\ &= \lim_{n \rightarrow \infty} \int_{I^2} f(s, t) \left[ \sum_{i,j=1}^n H(x_i^*, y_j^*, s, t) \Delta_{ij}f(x, y) \right] ds dt. \end{aligned}$$

But

$$(4.9) \quad \begin{aligned} & \left| \sum_{i,j=1}^n H(x_i^*, y_j^*, s, t) \Delta_{ij}f(x, y) \right| \\ & \leq \|f\| [\text{var}_{(x,y) \in I^2} H(x, y, s, t) + \text{var}_{x \in I} H(x, 1, s, t) + \text{var}_{y \in I} H(1, y, s, t) \\ & \quad + |H(1, 1, s, t)|] \\ & \leq 4 \|f\| M(s, t), \end{aligned}$$

and since  $M(s, t)$  is finite a.e. we see that for a.a.  $(s, t) \in I^2$

$$\lim_{n \rightarrow \infty} \sum_{i,j=1}^n H(x_i^*, y_j^*, s, t) \Delta_{ij} f(x, y) = \int_{I^2} H(x, y, s, t) df(x, y).$$

Thus (4.6) follows from this, (4.8) and (4.9) by dominated convergence.

**LEMMA 6.** *Let  $H(x, y, s, t)$  be as in Lemma 5, and let*

$$(4.10) \quad H^n(x, y, s, t) \equiv n^2 \int_{(q-1)/n}^{q/n} \int_{(p-1)/n}^{p/n} H(u, v, s, t) du dv$$

for  $(x, y) \in \left(\frac{p-1}{n}, \frac{p}{n}\right] \times \left(\frac{q-1}{n}, \frac{q}{n}\right]$ ,  $p, q = 1, 2, \dots, n$ .

Then for every  $f(x, y) \in C_Y$

$$(4.11) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \int_{I^2} \left[ \int_{I^2} H(x, y, s, t) f_{(n)}(s, t) ds dt \right]^2 dx dy \\ &= \int_{I^2} \left[ \int_{I^2} H(x, y, s, t) f(s, t) ds dt \right]^2 dx dy, \end{aligned}$$

$$(4.12) \quad \left| \int_{I^2} \left[ \int_{I^2} H^n(x, y, s, t) f_{(n)}(s, t) ds dt \right]^2 dx dy \right| \leq \left( \|f\| \int_{I^2} M(s, t) ds dt \right)^2,$$

$$(4.13) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \int_{I^2} \left[ \int_{I^2} H^n(x, y, s, t) f_{(n)}(s, t) ds dt \right] df_{(n)}(x, y) \\ &= \int_{I^2} \left[ \int_{I^2} H(x, y, s, t) f(s, t) ds dt \right] df(x, y). \end{aligned}$$

*Proof.* (4.11) and (4.12) are immediate. By (4.7) the function  $\int_{I^2} H(x, y, s, t) f(s, t) ds dt$  is of B.V. Now

$$\begin{aligned} & \int_{I^2} \left[ \int_{I^2} H^n(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\ &= \sum_{p,q=1}^n \int_{I^2} \left[ n^2 \int_{(p-1)/n}^{q/n} \int_{(q-1)/n}^{p/n} H(u, v, s, t) du dv \right] f(s, t) ds dt \cdot \Delta_{p,q} f, \end{aligned}$$

where  $\Delta_{p,q} f = f\left(\frac{p}{n}, \frac{q}{n}\right) - f\left(\frac{p-1}{n}, \frac{q}{n}\right) - f\left(\frac{p}{n}, \frac{q-1}{n}\right) + f\left(\frac{p-1}{n}, \frac{q-1}{n}\right)$ .

Since

$$\int_{I^2} \left[ n^2 \int_{(q-1)/n}^{q/n} \int_{(p-1)/n}^{p/n} H(u, v, s, t) du dv \right] f(s, t) ds dt$$

is the average value of  $\int_{I^2} H(x, y, s, t) f(s, t) ds dt$  in the square  $((q-1)/n, q/n] \times ((p-1)/n, p/n]$ , the existence of the R-S integral implies

$$(4.14) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \int_{I^2} \left[ \int_{I^2} H^n(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\ &= \int_{I^2} \left[ \int_{I^2} H(x, y, s, t) f(s, t) ds dt \right] df(x, y). \end{aligned}$$

Now,

$$(4.15) \quad \begin{aligned} \var_{(x,y) \in I^2} H^n(x, y, s, t) &\leq \var_{(x,y) \in I^2} H(x, y, s, t) \leq M(s, t) \\ \var_{x \in I} H^n(x, 1, s, t) &\leq \var_{x \in I} H(x, 1, s, t) \leq M(s, t) \\ \var_{y \in I} H^n(1, y, s, t) &\leq \var_{y \in I} H(1, y, s, t) \leq M(s, t) \\ |H^n(1, 1, s, t)| &\leq \sup_{(x,y) \in I^2} |H(x, y, s, t)| \leq M(s, t). \end{aligned}$$

Seeing the fact that  $H^n(x, y, s, t)$  is constant in  $x, y$  in each square  $\left(\frac{p-1}{n}, \frac{p}{n}\right] \left(\frac{q-1}{n}, \frac{q}{n}\right]$  and that  $f_{(n)}(x, y)$  and  $f(x, y)$  agree on the vertices of this square, we can write

$$\begin{aligned} & \int_{I^2} \left[ \int_{I^2} H^n(x, y, s, t) f_{(n)}(s, t) ds dt \right] df_{(n)}(x, y) \\ &= \sum_{p,q=1}^n \int_{I^2} H^n\left(\frac{p}{n}, \frac{q}{n}, s, t\right) f_{(n)}(s, t) ds dt \cdot \Delta_{pq} f \\ &= \int_{I^2} f_{(n)}(s, t) \left[ \sum_{p,q=1}^n H^n\left(\frac{p}{n}, \frac{q}{n}, s, t\right) \Delta_{pq} f \right] ds dt. \end{aligned}$$

Therefore

$$\begin{aligned} & \left| \int_{I^2} \left[ \int_{I^2} H^n(x, y, s, t) f(s, t) ds dt \right] df(x, y) \right. \\ & \quad \left. - \int_{I^2} \left[ \int_{I^2} H^n(x, y, s, t) f_{(n)}(s, t) ds dt \right] df_{(n)}(x, y) \right| \\ &= \left| \int_{I^2} [f(s, t) - f_{(n)}(s, t)] \left[ \sum_{p,q=1}^n H^n\left(\frac{p}{n}, \frac{q}{n}, s, t\right) \cdot \Delta_{pq} f \right] ds dt \right| \\ &\leq 4 \|f - f_n\| \|f\| \int_{I^2} M(s, t) ds dt, \end{aligned}$$

thus obtaining

$$(4.16) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \int_{I^2} \left[ \int_{I^2} H^n(x, y, s, t) f_{(n)}(s, t) ds dt \right] df_{(n)}(x, y) \\ &= \lim_{n \rightarrow \infty} \int_{I^2} \left[ \int_{I^2} H^n(x, y, s, t) f(s, t) ds dt \right] df(x, y). \end{aligned}$$

Hence (4.13) follows from (4.14) and (4.16).

**COROLLARY.** Let  $K(x, y, s, t)$  be as in Theorem I,  $K_\epsilon(x, y, s, t)$  the

corresponding function defined by (3.7), and  $K_{\varepsilon,(n)}(x, y, s, t)$  the  $n^{\text{th}}$  quasi-polyhedric function of  $K_\varepsilon$  in  $x, y$ . Then

$$(4.17) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{\varepsilon,(n)}(x, y, s, t) f_{(n)}(s, t) ds dt \right]^2 dx dy \\ &= \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) f(s, t) ds dt \right]^2 dx dy , \end{aligned}$$

$$(4.18) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{\varepsilon,(n)}(x, y, s, t) f_{(n)}(s, t) ds dt \right] df_{(n)}(x, y) \\ &= \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) f(s, t) ds dt \right] df(x, y) . \end{aligned}$$

*Proof.* In view of (3.9), (3.8), and (2.4) we obtain

$$(4.19) \quad \begin{aligned} \sup_{(x,y) \in I^2} |H_\varepsilon(x, y, s, t)| &\leq M(s, t) + \frac{\beta}{\varepsilon} + \frac{1}{4\varepsilon^2} |(K_1 + K_2 - K_3 \\ &\quad - K_4)(s, t, s, t)| , \\ \text{var}_{(x,y) \in I^2} H_\varepsilon(x, y, s, t) &\leq M(s, t) + \frac{2}{\varepsilon} M(s, t) + \frac{1}{\varepsilon^2} |(K_1 + K_2 - K_3 \\ &\quad - K_4)(s, t, s, t)| , \\ \text{var}_{x \in I} H_\varepsilon(x, 1, s, t) &\leq M(s, t) + \frac{1}{2\varepsilon} [2\beta + M(s, t)] + \frac{1}{2\varepsilon^2} |(K_1 + K_2 \\ &\quad - K_3 - K_4)(s, t, s, t)| , \\ \text{var}_{y \in I} H_\varepsilon(1, y, s, t) &\leq M(s, t) + \frac{1}{2\varepsilon} [2\beta + M(s, t)] + \frac{1}{2\varepsilon^2} |(K_1 + K_2 \\ &\quad - K_3 - K_4)(s, t, s, t)| . \end{aligned}$$

Let  $M_\varepsilon(s, t)$  be the sum of the four right members of (4.19). From (3.3) it is obvious that

$$(4.20) \quad \frac{\partial^2}{\partial x \partial y} K_{\varepsilon,(n)}(x, y, s, t) = n^2 \int_{(q-1)/n}^{q/n} \int_{(p-1)/n}^{p/n} \frac{\partial^2}{\partial v \partial u} K_\varepsilon(u, v, s, t) du dv$$

for  $(p-1)/n < x \leq p/n, (q-1)/n < y \leq q/n; p, q = 1, \dots, n$ . Hence from (4.10), (4.20), and the fact that  $(\partial^2/\partial v \partial u) K_\varepsilon(u, v, s, t) \equiv H_\varepsilon(u, v, s, t)$ , we see that

$$(4.21) \quad \frac{\partial^2}{\partial y \partial x} H_{\varepsilon,(n)}(x, y, s, t) \equiv H^n_\varepsilon(x, y, s, t) .$$

Thus the conclusions follow directly from Lemma 6.

5. Some discussions on resolvent kernels. Let us denote the resolvent kernel of  $K(x, y, s, t)$  in the Fredholm transformation (1.1) by  $K^*(x, y, s, t)$ , i.e., if

$$(5.1) \quad g(x, y) = f(x, y) + \int_{I^2} K(x, y, s, t) f(s, t) ds dt ,$$

then

$$(5.2) \quad f(x, y) = g(x, y) + \int_{I^2} K^*(x, y, s, t) g(s, t) ds dt .$$

Let  $K(x, y, s, t)$  be a bounded  $L^2$ -kernel on  $I^4$  with  $D(K) \neq 0$ , and set

$$(5.3) \quad \mathcal{D}(x, y, s, t) \equiv K^*(x, y, s, t) \cdot D(K) .$$

Then, using the familiar results for resolvent kernels for Fredholm integral equations with  $\lambda = -1$  (see [8], pp. 66–75), we can establish

$$(5.4) \quad K(x, y, s, t) + K^*(x, y, s, t) = - \int_{I^2} K(x, y, u, v) \cdot K^*(u, v, s, t) du dv ,$$

$$(5.5) \quad \mathcal{D}(x, y, s, t) = \sum_{n=0}^{\infty} C_n(x, y, s, t) / n!$$

where  $C_0(x, y, s, t) = -K(x, y, s, t)$ , and other  $C_n$ 's are found successively by

$$(5.6) \quad \begin{aligned} C_n(x, y, s, t) &= J_n(K) \cdot K(x, y, s, t) \\ &\quad - n \int_{I^2} K(x, y, u, v) C_{n-1}(u, v, s, t) du dv , \\ J_n(K) &= - \int_{I^2} \cdots \int_{I^2} \left| \begin{array}{c} K(x_1, y_1, x_1, y_1) \cdots K(x_1, y_1, x_n, y_n) \\ \cdots \cdots \cdots \cdots \cdots \cdots \\ K(x_n, y_n, x_1, y_1) \cdots K(x_n, y_n, x_n, y_n) \end{array} \right| \\ &\quad dx_1 dy_1 \cdots dx_n dy_n . \end{aligned}$$

By Hadamard's inequality it follows from (5.6) that

$$|C_n(x, y, s, t)| \leq (n+1)^{(n+1)/2} \|K\|^{n+1} .$$

Therefore the series in (5.5) is absolutely and uniformly convergent. If  $K(x, y, s, t)$  satisfies the assumptions in Theorem I, then

$$\int_{I^2} K(x, y, u, v) C_n(u, v, s, t) du dv$$

is continuous on  $I^4$  for  $n = 0, 1, 2, \dots$ , and hence from (5.6) we see that the jumps for  $C_n(u, v, s, t)$  coincides with those for  $K(x, y, s, t)$ , and thus it takes average value at each jump, and so does  $\sum_{n=0}^{\infty} C_n(x, y, s, t) / n!$  by uniform convergence. By absolute convergence

we may rearrange the terms in the series (5.5), and then use the uniform convergence to obtain

$$\begin{aligned}
 (5.7) \quad \mathcal{D}(x, y, s, t) &= \sum_{n=0}^{\infty} \frac{1}{n!} \left[ J_n \cdot K(x, y, s, t) \right. \\
 &\quad \left. - n \int_{I^2} K(x, y, u, v) \cdot C_{n-1}(u, v, s, t) du dv \right] \\
 &= C \cdot K(x, y, s, t) - \int_{I^2} K(x, y, u, v) \cdot \mathcal{D}(u, v, st) du dv ,
 \end{aligned}$$

where  $C = \sum_{n=0}^{\infty} J_n(K)/n!$ . Corresponding to each  $K \in (x, y, s, t)$ , define  $K_{(\varepsilon)}^*(x, y, s, t)$  by

$$\begin{aligned}
 (5.8) \quad K_{(\varepsilon)}^*(x, y, s, t) &= D(K) \mathcal{D}_{(\varepsilon)}(x, y, s, t) \\
 &= D(K) \left[ C \cdot K_{\varepsilon}(x, y, s, t) \right. \\
 &\quad \left. - \int_{I^2} K_{\varepsilon}(x, y, u, v) \cdot \mathcal{D}(u, v, s, t) du dv \right] .
 \end{aligned}$$

Then  $K_{(\varepsilon)}^*(x, y, s, t)$  is continuous on  $I^4$ , and since  $\mathcal{D}(u, v, s, t)$  is bounded, we have that  $K_{(\varepsilon)}^*(x, y, s, t)$  is uniformly bounded in  $\varepsilon, x, y, s, t$ , and from (3.9), (4.19), and (5.8) it follows that

$$\begin{aligned}
 (5.9) \quad H_{(\varepsilon)}^*(x, y, s, t) &\equiv \frac{\partial^2}{\partial y \partial x} K_{(\varepsilon)}^*(x, y, s, t) \\
 &= D(K) \left[ C \cdot H_{\varepsilon}(x, y, s, t) \right. \\
 &\quad \left. - \int_{I^2} H_{\varepsilon}(x, y, u, v) \cdot \mathcal{D}(u, v, s, t) du dv \right] ,
 \end{aligned}$$

and  $\sup_{(x, y) \in I^2} |H_{(\varepsilon)}^*(x, y, s, t)|$ ,  $\text{var}_{(x, y) \in I^2} H_{(\varepsilon)}^*(x, y, s, t)$ ,  $\text{var}_{x \in I} H_{(\varepsilon)}^*(x, 1, s, t)$ , and  $\text{var}_{y \in I} H_{(\varepsilon)}^*(1, y, s, t)$  are all dominated by  $|D(K)|(|C| + \|\mathcal{D}\|)$ .  $M_{\varepsilon}(s, t)$ . Furthermore, by Lemma 4

$$(5.10) \quad \lim_{\varepsilon \rightarrow 0+} D(K_{(\varepsilon)}^*) = D(K^*) = D^{-1}(K) .$$

**6. Additional lemmas.** Utilizing (3.7), (3.9), (3.11), (5.8), and (5.9), we have the following

**LEMMA 7.** *If  $K(x, y, s, t)$  satisfies the hypotheses of Theorem I, then*

$$\begin{aligned}
 (6.1) \quad &\left| \int_{I^2} [K_{\varepsilon}(x, y, s, t) - K(x, y, s, t)] f(s, t) ds dt \right| \\
 &\leq 4\varepsilon(1 + \varepsilon) \|f\| \cdot \|K\| , \\
 &\int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{\varepsilon}(x, y, s, t) f(s, t) ds dt \right]^2 dx dy
 \end{aligned}$$

$$\begin{aligned}
&\leq \|f\|^2 \left[ \int_{I^2} M(s, t) ds dt + 2\beta + 4\|K\| \right]^2, \\
(6.2) \quad &\lim_{\varepsilon \rightarrow 0^+} \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) f(s, t) ds dt \right]^2 dx dy \\
&= \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) f(s, t) ds dt \right]^2 dx dy,
\end{aligned}$$

$$\begin{aligned}
(6.3) \quad &\left| \int_{I^2} [K_{(\varepsilon)}^*(x, y, s, t) - K^*(x, y, s, t)] f(s, t) ds dt \right| \\
&\leq 4\varepsilon(1 + \varepsilon) \|f\| \|K\| |D(K)| \cdot (|C| + \|\mathcal{D}\|),
\end{aligned}$$

$$\begin{aligned}
(6.4) \quad &\int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{(\varepsilon)}^*(x, y, s, t) f(s, t) ds dt \right]^2 dx dy \\
&\leq (\|f\| |D(K)|)^2 (|C| + \|\mathcal{D}\|) \left[ \int_{I^2} M(s, t) ds dt + 2\beta + 4\|K\| \right]^2,
\end{aligned}$$

$$\begin{aligned}
(6.5) \quad &\lim_{\varepsilon \rightarrow 0^+} \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{(\varepsilon)}^*(x, y, s, t) f(s, t) ds dt \right]^2 dx dy \\
&= \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) f(s, t) ds dt \right]^2 dx dy.
\end{aligned}$$

The following two lemmas are the key results:

LEMMA 8. Let  $K(x, y, s, t)$  be as in Theorem I. Then

$$\int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) f(s, t) ds dt \right] d^* f(x, y)$$

defined as in (1.2) converges for a.a.  $f$  in  $C_Y$ , and any two C.O.N. sets in the class lead to the same value for a.a.  $f$  in  $C_Y$ . Furthermore, we have

$$\begin{aligned}
(6.6) \quad &\text{l.i.m.}_{\varepsilon \rightarrow 0^+} \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\
&= \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) f(s, t) ds dt \right] d^* f(x, y) \text{ on } C_Y,
\end{aligned}$$

where the mean convergence is in  $L^2$ -sense.

*Proof.* First we observe that

$$\int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) f(s, t) ds dt \right] d^* f(x, y)$$

$$\begin{aligned}
&= \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) f(s, t) ds dt \right] d^* f(x, y) \\
(6.7) \quad &+ \int_{I^2} \left[ \int_0^1 B(y, x, t) f(x, t) dt \right] d^* f(x, y) \\
&+ \int_{I^2} \left[ \int_0^1 A(x, s, y) f(s, y) ds \right] d^* f(x, y) \\
&+ \int_{I^2} (K_1 + K_2 - K_3 - K_4)(x, y, x, y) \cdot f(x, y) d^* f(x, y) ,
\end{aligned}$$

and the existence and the consistency of each of last three expressions follow in the similar manner as in the generalized P.W.Z. integral  $\int_{I^2} h f d^* f$ . On account of (3.9) and Lemma 5 we may write

$$\begin{aligned}
&\int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\
&= \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\
(6.8) \quad &+ \int_{I^2} f(s, t) \left[ \int_{I^2} C_\varepsilon(x, s) B(y, s, t) df(x, y) \right] ds dt \\
&+ \int_{I^2} f(s, t) \left[ \int_{I^2} C_\varepsilon(y, t) A(x, s, t) df(x, y) \right] ds dt \\
&+ \int_{I^2} (K_1 + K_2 - K_3 - K_4)(s, t, s, t) \left[ \int_{I^2} C_\varepsilon(x, s) C_\varepsilon(y, t) df(x, y) \right] ds dt .
\end{aligned}$$

Obviously,

$$\begin{aligned}
&\int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\
&= \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K(x, y, s, t) f(s, t) ds dt \right] d^* f(x, y) \text{ for a.a. } f \in C_Y .
\end{aligned}$$

Thus to establish (6.6) we need only show that

$$\begin{aligned}
(6.9) \quad &\text{l.i.m.}_{\varepsilon \rightarrow 0^+} \int_{I^2} f(s, t) \left[ \int_{I^2} C_\varepsilon(x, s) B(y, s, t) df(x, y) \right] ds dt \\
&= \int_{I^2} \left[ \int_0^1 B(y, x, t) f(x, t) dt \right] d^* f(x, y) \text{ on } C_Y ,
\end{aligned}$$

and so on. To see this we observe that the Yeh-Wiener integral

$$\begin{aligned}
&\int_{C_Y} \left\{ \int_{I^2} f(s, t) \left[ \int_{I^2} C_\varepsilon(x, s) B(y, s, t) df(x, y) \right] ds dt \right\}^2 d_Y f \\
&= \frac{1}{4} \left\{ \int_{I^4} \left[ \int_s^1 \int_t^1 C_\varepsilon(x, s') B(y, s', t') ds' dt' \right]^2 ds dt dx dy
\end{aligned}$$

$$\begin{aligned}
& + \left[ \int_{I^2} \left( \int_0^t \int_0^s C_\varepsilon(x, s) B(y, s, t) dx dy \right) ds dt \right]^2 \\
& + \int_{I^4} \left( \int_0^t \int_0^s C_\varepsilon(x, s') B(y, s', t') dx dy \right) \\
& \quad \times \left( \int_0^{t'} \int_0^{s'} C_\varepsilon(x, s) B(y, s, t) dx dy \right) ds dt ds' dt' \Big\} , \\
& \int_{C_Y} \left\{ \int_{I_1} \left[ \int_0^1 B(y, x, t) f(x, t) dt \right] d^* f(x, y) \right\}^2 d_Y f \\
& = \frac{1}{4} \left\{ \int_{I^3} \left[ \int_x^1 \left( \int_t^1 B(y, x', t') dt' \right)^2 dx' \right] dx dt dy \right. \\
& \quad \left. + \frac{1}{4} \left[ \int_{I^2} \left( \int_y^1 B(y, x, t) dt \right) dx dy \right]^2 \right\} ,
\end{aligned}$$

and

$$\begin{aligned}
& \int_{C_Y} \left\{ \int_{I^2} f(s, t) \left[ \int_{I^2} C_\varepsilon(x, s) B(y, s, t) df(x, y) \right] ds dt \right\} \\
& \quad \cdot \left\{ \int_{I^2} \left[ \int_0^1 B(y, x, t) f(x, t) dt \right] d^* f(x, y) \right\} d_Y f \\
& = \frac{1}{4} \left\{ \int_{I^3} \left[ \int_{I^2} B(y, x, t') \min(s, x) C_\varepsilon(x, s) B(y, s, t) dx dy \right] \min(t, t') ds dt dt' \right. \\
& \quad + \int_{I^3} \left( \int_0^t \int_0^s B(y', x', t) \left[ \int_0^{t'} \int_0^{s'} C_\varepsilon(x, s) B(y, s, t) dx dy \right] dx' dy' \right) ds dt dt' \\
& \quad \left. + \frac{1}{2} \int_{I^4} \left( \int_0^{t'} B(y', x', t') \left[ \int_0^t \int_0^s C_\varepsilon(s, x) B(y, s, t) dx dy \right] dy' \right) dx' ds dt dt' \right\} .
\end{aligned}$$

The techniques leading to the evaluation of above integrals can be found in [3] and [4]. Using these and then taking the limit as  $\varepsilon \rightarrow 0^+$ , we get

$$\lim_{\varepsilon \rightarrow 0^+} \int_{C_Y} \left\{ \int_{I^2} f(s, t) \left[ \int_{I^2} C_\varepsilon(x, s) B(y, s, t) df(x, y) \right] ds dt \right\} \\
- \left\{ \int_{I^2} \left[ \int_0^1 B(y, x, t) f(x, t) dt \right] d^* f(x, y) \right\}^2 d_Y f = 0 ,$$

which is exactly the same as (6.9).

**LEMMA 8'.** *Let  $K(x, y, s, t)$  satisfy the hypotheses of Theorem I. Then for its resolvent kernel  $K^*$  we have*

$$(6.10) \quad \int_{I^2} \left[ \frac{\partial^2}{\partial y \partial x} \int_{I^2} K^*(x, y, s, t) f(x, t) ds dt \right] d^* f(x, y)$$

*converges for almost all  $f$  in  $C_Y$  and is essentially independent of the particular choice of the C.O.N. set, and*

$$(6.11) \quad \begin{aligned} & \text{l.i.m.}_{\varepsilon \rightarrow 0^+} \left[ \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{(\varepsilon)}^*(x, y, s, t) f(s, t) ds dt \right] df(x, y) \right. \\ & \quad \left. = \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K^*(x, y, s, t) f(s, t) ds dt \right] d^*f(x, y) \quad \text{on } C_Y. \right. \end{aligned}$$

*Proof.* The claim on (6.10) can be treated as that in (6.7). As for (6.11) we may use (5.9) with  $H_\varepsilon(x, y, s, t) \equiv \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t)$  to see that

$$\begin{aligned} & \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{(\varepsilon)}^*(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\ & = D(K) \left\{ C \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) f(s, t) ds dt \right] df(x, y) \right. \\ & \quad \left. - \int_{I^2} \left[ \int_{I^2} \left( \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, u, v) \mathcal{D}(u, v, s, t) du dv \right) f(s, t) ds dt \right] df(x, y) \right\}, \end{aligned}$$

and the first term in the braces converges in the mean to

$$C \int_{I^2} \left[ (\partial^2 / \partial y \partial x) \int_{I^2} K(x, y, s, t) f(s, t) ds dt \right] d^*f(x, y)$$

on  $C_Y$  by the preceding lemma. Therefore, in view of (5.3) and (5.7), it remains to show that

$$\begin{aligned} & \text{l.i.m.}_{\varepsilon \rightarrow 0^+} \left[ \int_{I^2} \left( \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, u, v) \mathcal{D}(u, v, s, t) du dv \right) f(s, t) ds dt \right] df(x, y) \\ & = \int_{I^2} \left[ \frac{\partial^2}{\partial y \partial x} \int_{I^2} \left( \int_{I^2} K(x, y, u, v) \mathcal{D}(u, v, s, t) du dv \right) f(s, t) ds dt \right] d^*f(x, y) \end{aligned}$$

on  $C_Y$  whose proof is essentially on the same lines as that of Lemma 8.

**LEMMA 9.** *Let  $F(f)$  be a nonnegative Yeh-Wiener measurable functional on  $C_Y$ , and let  $K(x, y, s, t)$  satisfy the hypotheses of Theorem I. Then under the Fredholm transformation  $T$  in (1.1), we have*

$$(6.12) \quad \int_{C_Y} F(g) d_Y g \geq |D(K)| \int_{C_Y} F(Tf) \cdot \exp \{-\Phi(f)\} d_Y f,$$

where  $\Phi(f)$  is given by (2.8).

*Proof.* First we assume that  $F(f)$  is a bounded nonnegative functional continuous on  $C_Y$  with respect to the uniform topology, and is dependent only on the  $n^2$  values of  $f$  at  $(x, y) = (i/n, j/n)$ ;  $i, j = 1, 2, \dots, n$ . Since  $\lim_{\varepsilon \rightarrow 0^+} D(K_\varepsilon) = D(K) \neq 0$  and  $\lim_{n \rightarrow \infty} D(K_{\varepsilon(n)}) = D(K_\varepsilon)$

by Lemma 4, there exist  $\varepsilon'$  and  $N(\varepsilon)$  such that  $0 < \varepsilon < \varepsilon'$  and  $n \geq N(\varepsilon)$  imply  $D(K_\varepsilon) \neq 0$  and  $D(K_{\varepsilon, (n)}) \neq 0$ . For such  $\varepsilon$  and  $n$  we may apply Lemma 2 to obtain

$$(6.13) \quad \begin{aligned} \int_{C_Y} F(g_n) d_Y g &= |D(K_{\varepsilon, (n)})| \int_{C_Y} F \left[ f_{(n)} + \int_{I^2} K_{\varepsilon, (n)}(\cdot, \cdot, s, t) f_{(n)}(s, t) ds dt \right] \\ &\quad \cdot \exp \left\{ - \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{\varepsilon, (n)}(x, y, s, t) f_{(n)}(s, t) ds dt \right]^2 dx dy \right. \\ &\quad \left. - 2 \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{\varepsilon, (n)}(x, y, s, t) f_{(n)}(s, t) ds dt \right] df_{(n)}(x, y) \right\} d_Y f. \end{aligned}$$

As  $n \rightarrow \infty$ ,  $f_{(n)}(x, y) \rightharpoonup f(x, y)$  and  $K_{\varepsilon, (n)}(x, y, s, t) \rightharpoonup K_\varepsilon(x, y, s, t)$ . Hence  $g_n(x, y) \equiv f_{(n)}(x, y) + \int_{I^2} K_{\varepsilon, (n)}(x, y, s, t) f_{(n)}(s, t) ds dt \rightharpoonup g(x, y) = f(x, y) + \int_{I^2} K_\varepsilon(x, y, s, t) f(s, t) ds dt$ . Therefore  $F(g_n) \rightarrow F(g)$  boundedly. Thus  $\lim_n \int_{C_Y} F(g_n) d_Y g = \int_{C_Y} F(g) d_Y g$ . Now, by Fatou's lemma it follows from (6.13) with the help of Lemma 3, (4.11), and (4.13) that

$$(6.14) \quad \begin{aligned} \int_{C_Y} F(g) d_Y g &\geq |D(K_\varepsilon)| \int_{C_Y} F \left[ f + \int_{I^2} K_\varepsilon(\cdot, \cdot, s, t) f(s, t) ds dt \right] \\ &\quad \cdot \exp \left\{ - \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) f(s, t) ds dt \right]^2 dx dy \right. \\ &\quad \left. - 2 \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_\varepsilon(x, y, s, t) f(s, t) ds dt \right] df(x, y) \right\} d_Y f. \end{aligned}$$

By (6.1)  $\int_{I^2} K_\varepsilon(x, y, s, t) f(s, t) ds dt \rightharpoonup \int_{I^2} K(x, y, s, t) f(s, t) ds dt$  as  $\varepsilon \rightarrow 0^+$ ,

and hence

$$F \left[ f + \int_{I^2} K_\varepsilon(\cdot, \cdot, s, t) f(s, t) ds dt \right] \rightarrow F \left[ f + \int_{I^2} K(\cdot, \cdot, s, t) f(s, t) ds dt \right]$$

boundedly. Since "mean convergence" implies the existence of an almost everywhere convergent subsequence, it follows from (6.6) that there exists a monotone sequence  $\varepsilon_n \downarrow 0$  such that

$$(6.15) \quad \begin{aligned} \lim_{n \rightarrow \infty} \int_{I^2} \left[ \int_{I^2} \frac{\partial^2}{\partial y \partial x} K_{\varepsilon_n}(x, y, s, t) f(s, t) ds dt \right] df(x, y) \\ = \int_{I^2} \left[ \frac{\partial^2}{\partial y \partial x} \int_{I^2} K(x, y, s, t) f(s, t) ds dt \right] d^* f(x, y) \text{ for a.a. } f \in C_Y. \end{aligned}$$

Now we replace  $\varepsilon_n$  for  $\varepsilon$  in (6.14), and then use Fatou's Lemma together with (6.2) and (6.15) to arrive at (6.12). This completes the proof for the case when  $F(f)$  is nonnegative, bounded, and continuous in uniform topology. To obtain the result for arbitrary nonnegative measurable

functionals, we go through, as usual, the following steps: after proving it for the preceding case, we go to characteristic functionals of intervals, then 0-sets, then  $0_\delta$ -sets, and then null-sets. (In the last case we get equality rather than inequality, both sides being zero.) Then to characteristic functionals of measurable sets, nonnegative simple functionals, nonnegative bounded Yeh-Wiener measurable functionals, and then finally nonnegative Yeh-Wiener measurable functionals. For the details of these steps see [2: pp. 391-392].

On account of (5.9), (5.10), (6.3), (6.5), and (6.11) we can establish the following on the same lines as above:

**LEMMA 9'.** *Let  $F(g)$  be a nonnegative Yeh-Wiener measurable functional on  $C_Y$ , and let  $K(x, y, s, t)$  be as in Theorem I and  $K^*(x, y, s, t)$  its resolvent kernel. Then under the transformation*

$$T^{-1}: g(x, y) \rightarrow f(x, y) = g(x, y) + \int_{I^2} K^*(x, y, s, t)g(s, t)dsdt ,$$

we have

$$\int_{C_Y} F(f)d_Y f \geq |D^{-1}(K)| \int_{C_Y} F(T^{-1}g) \cdot \exp\{-\Phi^*(g)\}d_Y g ,$$

where

$$(6.16) \quad \begin{aligned} \Phi^*(g) &= \int_{I^2} \left[ \frac{\partial^2}{\partial x \partial y} \int_{I^2} K^*(x, y, s, t)g(s, t)dsdt \right]^2 dx dy \\ &\quad + 2 \int_{I^2} \left[ \frac{\partial^2}{\partial y \partial x} \int_{I^2} K^*(x, y, s, t)g(s, t)dsdt \right] d^*g(x, y) . \end{aligned}$$

**7. Proof of theorems.** We prove Theorem I for nonnegative Yeh-Wiener measurable functionals first. In this case by Lemma 9 we have under the transformation  $T$  in (1.1),

$$(7.1) \quad \int_{C_Y} F(g)d_Y g \geq |D(K)| \int_{C_Y} F[Tf] \exp\{-\Phi(f)\}d_Y f ,$$

and upon applying Lemma 9' on the right-hand side of (7.1), we obtain under the transformation

$$(7.2) \quad \begin{aligned} T^{-1}: g(x, y) \rightarrow f(x, y) &= g(x, y) + \int_{I^2} K^*(x, y, s, t)g(s, t)dsdt \\ &\quad \int_{C_Y} F[Tf] \exp\{-\Phi(f)\}d_Y f \\ &\geq |D^{-1}(K)| \int_{C_Y} F(T \circ T^{-1}g) \exp\{-\Phi(T^{-1}g)\} \cdot \exp\{-\Phi^*(g)\}d_Y g . \end{aligned}$$

Hence if we show that

$$(7.3) \quad \Phi(T^{-1}g) + \Phi^*(g) = 0 \quad \text{for a.a. } g \text{ in } C_Y ,$$

then it will follow from (7.1) and (7.2) that

$$\begin{aligned} \int_{C_Y} F(g) d_Y g &\geq |D(K)| \int_{C_Y} F[Tf] \exp\{-\Phi(f)\} d_Y f \\ &\geq \int_{C_Y} F(g) d_Y g , \end{aligned}$$

and hence the theorem will follow for the case. Observe now that if  $T(f) = g$  and so  $T^{-1}(g) = f$ , then

$$(7.4) \quad \begin{aligned} \int_{r^2} K(x, y, s, t) f(s, t) ds dt &= g(x, y) - f(x, y) , \\ \int_{r^2} K^*(x, y, s, t) g(s, t) ds dt &= f(x, y) - g(x, y) , \end{aligned}$$

and from (3.10) and (3.11), we see that the left-hand side of the first equation in (7.4) is absolutely continuous, and

$$\frac{\partial^2}{\partial y \partial x} \int_{r^2} K(x, y, s, t) f(s, t) ds dt$$

is continuous. Therefore  $(\partial^2/\partial y \partial x)[g(x, y) - f(x, y)]$  is also continuous. Therefore by the corollary to Theorem 4 in [3], we see that

$$(7.5) \quad \begin{aligned} \int_{r^2} \frac{\partial^2}{\partial y \partial x} [g(x, y) - f(x, y)] d^*[g(x, y) - f(x, y)] \\ \equiv \int_{r^2} \left\{ \frac{\partial^2}{\partial y \partial x} [g(x, y) - f(x, y)] \right\}^2 dx dy . \end{aligned}$$

Hence from (2.8), (7.4), and (7.5) it follows that

$$(7.6) \quad \begin{aligned} \Phi(T^{-1}g) &= \Phi(f) \\ &= \int_{r^2} \left\{ \frac{\partial^2}{\partial y \partial x} [g(x, y) - f(x, y)] \right\}^2 dx dy \\ &\quad + 2 \int_{r^2} \frac{\partial^2}{\partial y \partial x} [g(x, y) - f(x, y)] d^* f(x, y) \\ &= \int_{r^2} \left\{ \frac{\partial^2}{\partial y \partial x} [g(x, y) - f(x, y)] \right\}^2 dx dy \\ &\quad + 2 \int_{r^2} \frac{\partial^2}{\partial y \partial x} [g(x, y) - f(x, y)] d^* f(x, y) \\ &\quad - 2 \int_{r^2} \left\{ \frac{\partial^2}{\partial y \partial x} [g(x, y) - f(x, y)] \right\}^2 dx dy , \end{aligned}$$

and from (6.16) and (7.4)

$$(7.7) \quad \begin{aligned} \Phi^*(g) &= \int_{\mathbb{R}^2} \left\{ \frac{\partial^2}{\partial y \partial x} [f(x, y) - g(x, y)] \right\}^2 dx dy \\ &\quad + 2 \int_{\mathbb{R}^2} \frac{\partial^2}{\partial y \partial x} [f(x, y) - g(x, y)] d^* g(x, y). \end{aligned}$$

Therefore (7.3) follows from (7.6) and (7.7). Hence the theorem holds for any nonnegative Yeh-Wiener measurable functionals. For arbitrary real Yeh-Wiener measurable functionals the theorem will also hold by considering the positive part and the negative part separately. For complex functionals we get the same result once we consider real part and imaginary part separately. This completes the proof of Theorem I.

To prove Theorem II we consider the following two transformations

$$\begin{aligned} I_1: f(x, y) &\rightarrow g(x, y) = f(x, y) + f_0(x, y), \\ I_2: f(x, y) &\rightarrow h(x, y) = f(x, y) + \int_{\mathbb{R}^2} K(x, y, s, t) f(s, t) ds dt. \end{aligned}$$

The theorem now follows by the use of Cameron-Martin translation theorem (see [11] or more precisely Theorem 1.4 in [4] on  $L_1$  and then our Theorem I for  $L_2$ .

**REMARK.** As mentioned in the introduction our theorems in one-dimensional version have slightly different forms from the ones given in [1], the difference being in the expressions of  $\Phi(f)$  and  $\Psi(f)$ . The  $\Phi(f)$  in [1] is given by

$$(7.8) \quad \begin{aligned} \Phi(f) &= \int_0^1 \left[ \frac{d}{dx} \int_0^1 K(x, s) f(s) ds \right]^2 dx \\ &\quad + 2 \int_0^1 \left[ \int_0^1 \frac{\partial}{\partial x} K(x, s) f(s) ds \right] df(x) + \int_0^1 J(x) d[f^2(x)], \end{aligned}$$

and ours is in the form

$$(7.9) \quad \Phi(f) = \int_0^1 \left[ \frac{d}{dx} \int_0^1 K(x, s) f(s) ds \right]^2 dx + 2 \int_0^1 \left[ \frac{d}{dx} \int_0^1 K(x, s) f(s) ds \right] d^* f(x).$$

However by the assumptions given on  $K(x, s)$  in [1], we have that

$$\begin{aligned} \frac{d}{dx} \int_0^1 K(x, s) f(s) ds &= \frac{d}{dx} \int_0^x K_2(x, s) f(s) ds + \int_x^1 K_1(x, s) f(s) ds \\ &= [K_2(x, x) - K_1(x, x)] f(x) + \int_0^1 \frac{\partial}{\partial x} K(x, s) f(s) ds \end{aligned}$$

$$= J(x)f(x) + \int_0^1 \frac{\partial}{\partial x} K(x, s)f(s)ds .$$

But assuming absolute continuity of  $J(x)$  with  $J'(x) \in L^2$ , it follows from Theorem 5 in [3] that

$$\int_0^1 J(x)f(x)d^*f(x) = \frac{1}{2} \int_0^1 J(x)d[f^2(x)] \quad \text{for a.a. } f \in C_w .$$

Also as mentioned earlier

$$\int_0^1 \left[ \int_0^1 \frac{\partial}{\partial x} K(x, s)f(s)ds \right] d^*f(x) = \int_0^1 \left[ \int_0^1 \frac{\partial}{\partial x} K(x, s)f(s)ds \right] df(x)$$

for almost all  $f$  in  $C_w$ . Thus (7.8) and (7.9) represent essentially the same thing.

#### REFERENCES

1. R. H. Cameron and W. T. Martin, *Transformations of Wiener integrals under a general class of linear transformations*, Trans. Amer. Math. Soc., **58** (1945), 184-219.
2. ———, *On transformations of Wiener integrals under transformations*, Ann. of Math., **45** (1944), 386-396.
3. C. Park, *A generalized Paley-Wiener-Zygmund integral and its applications*, Proc. Amer. Math. Soc., **23** (1969), 388-400.
4. ———, *Generalized Riemann-Stieltjes integral over the K.Y.W. space of functions of two variables*, Thesis, Univ. of Minnesota (1968).
5. I. E. Segal, *Distributions in Hilbert space and canonical systems of operators*, Trans. Amer. Math. Soc., **88** (1958), 12-41.
6. ———, *Algebraic integration theory*, Bull. Amer. Math. Soc., **71** (1965), 419-489.
7. L. A. Shepp, *Radon-Nikodym derivatives of Gaussian measures*, Ann. Math. Statist., **37** (1966), 321-354.
8. F. G. Tricomi, *Integral equations*, Interscience Publishers (1957).
9. D. A. Woodward, *A general class of linear transformations of Wiener integrals*, Trans. Amer. Math. Soc., **100** (1961), 459-480.
10. J. Yeh, *Wiener measure in a space of functions of two variables*, Trans. Amer. Math. Soc., **95** (1960), 433-450.
11. ———, *Cameron-Martin translation theorems in the Wiener space of functions of two variables*, Trans. Amer. Math. Soc., **107** (1963), 1427-1436.
12. ———, *Orthogonal development of functionals and related theorems in the Wiener space of functions of two variables*, Pacific J. Math., **13** (1963), 1427-1436.

Received October 23, 1970. This research was performed at Miami University and the U.S.A.F. Aerospace Research Laboratories while in the capacity of an Ohio State University Research Foundation visiting Research Associate under Contract F33615 67 C 1758.

MIAMI UNIVERSITY



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

C. R. HOBBY  
University of Washington  
Seattle, Washington 98105

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

# Pacific Journal of Mathematics

Vol. 40, No. 1

September, 1972

Alex Bacopoulos and Athanassios G. Kartsatos, <i>On polynomials approximating the solutions of nonlinear differential equations</i> .....	1
Monte Boisen and Max Dean Larsen, <i>Priifer and valuation rings with zero divisors</i> .....	7
James J. Bowe, <i>Neat homomorphisms</i> .....	13
David W. Boyd and Hershy Kisilevsky, <i>The Diophantine equation <math>u(u+1)(u+2)(u+3) = v(v+1)(v+2)</math></i> .....	23
George Ulrich Brauer, <i>Summability and Fourier analysis</i> .....	33
Robin B. S. Brooks, <i>On removing coincidences of two maps when only one, rather than both, of them may be deformed by a homotopy</i> .....	45
Frank Castagna and Geert Caleb Ernst Prins, <i>Every generalized Petersen graph has a Tait coloring</i> .....	53
Micheal Neal Dyer, <i>Rational homology and Whitehead products</i> .....	59
John Fuelberth and Mark Lawrence Teply, <i>The singular submodule of a finitely generated module splits off</i> .....	73
Robert Gold, <i><math>\Gamma</math>-extensions of imaginary quadratic fields</i> .....	83
Myron Goldberg and John W. Moon, <i>Cycles in <math>k</math>-strong tournaments</i> .....	89
Darald Joe Hartfiel and J. W. Spellmann, <i>Diagonal similarity of irreducible matrices to row stochastic matrices</i> .....	97
Wayland M. Hubbart, <i>Some results on blocks over local fields</i> .....	101
Alan Loeb Kostinsky, <i>Projective lattices and bounded homomorphisms</i> .....	111
Kenneth O. Leland, <i>Maximum modulus theorems for algebras of operator valued functions</i> .....	121
Jerome Irving Malitz and William Nelson Reinhardt, <i>Maximal models in the language with quantifier "there exist uncountably many"</i> .....	139
John Douglas Moore, <i>Isometric immersions of space forms in space forms</i> .....	157
Ronald C. Mullin and Ralph Gordon Stanton, <i>A map-theoretic approach to Davenport-Schinzel sequences</i> .....	167
Chull Park, <i>On Fredholm transformations in Yeh-Wiener space</i> .....	173
Stanley Poreda, <i>Complex Chebyshev alterations</i> .....	197
Ray C. Shiflett, <i>Extreme Markov operators and the orbits of Ryff</i> .....	201
Robert L. Snider, <i>Lattices of radicals</i> .....	207
Ralph Richard Summerhill, <i>Unknotting cones in the topological category</i> .....	221
Charles Irvin Vinsonhaler, <i>A note on two generalizations of QF-3</i> .....	229
William Patterson Wardlaw, <i>Defining relations for certain integrally parameterized Chevalley groups</i> .....	235
William Jennings Wickless, <i>Abelian groups which admit only nilpotent multiplications</i> .....	251