

Pacific Journal of Mathematics

COMPLEX CHEBYSHEV ALTERATIONS

STANLEY POREDA

COMPLEX CHEBYSHEV ALTERATIONS

S. J. POREDA

P. Chebyshev's famous Alternation Theorem for best uniform approximation to continuous real valued functions on an interval is generalized to include best approximation to a class of continuous complex valued functions on an ellipse.

1. Preliminary remarks and definitions. For a continuous complex valued function f defined on a compact set E in the plane and, for $n \in \mathbb{Z}^+$, let $p_n(f, E)$ denote the polynomial of degree n , of best uniform approximation to f on E and let;

$$\rho_n(f, E) = \max_{z \in E} |f(z) - p_n(f, E)(z)|.$$

Chebyshev's Alternation Theorem [1, p. 29] states that if f is a continuous real valued function on an interval $[a, b]$, and p_n is a polynomial of degree n , $n \in \mathbb{Z}^+$, then $p_n = p_n(f, [a, b])$ if and only if, there exists $n + 2$ points,

$\{x_i\}_{i=1}^{n+2}$, $a \leq x_1 < x_2 < \dots < x_{n+2} \leq b$, with the property that $|f(x) - p_n(x)|$ attains its maximum on $[a, b]$ at these points and $f(x_i) - p_n(x_i) = -[f(x_{i+1}) - p_n(x_{i+1})]$ for $i = 1, 2, \dots, n + 1$.

The sets we consider here are ellipses which are of course a generalization of intervals. So, for $a \geq 0$, let $E_a = \{z + a/z : |z| = 1\}$. Now let $\mathcal{F}_n(E_a)$ denote those complex valued functions f , not themselves polynomials of degree n , continuous on E_a , having the property that there exists $n + 2$ points $\{\xi_k\}_{k=1}^{n+2}$ in E_a , such that $p_n(f, E_a) = p_n(f, \{\xi_k\}_{k=1}^{n+2})$. It is known [1, p. 22] that there always exists a set $D \subset E_a$, consisting of $n + k$ points, $2 \leq k \leq n + 3$, such that $p_n(f, E_a) = p_n(f, D)$. Furthermore, to this author's knowledge, every example of best uniform approximation to rational functions on infinite sets in the plane (e.g., [3], [4] and [5]) is one in which such a set consisting of $n + 2$ points exists or, can be shown equivalent to such an example.

2. Main theorem. Given $n + 2$ points $\{\xi_k\}_{k=1}^{n+2}$ in E_a let z_k be such that $\xi_k = z_k + a/z_k$, $|z_k| = 1$ and if $a = 1$, $0 \leq \text{Arg } z_k \leq \pi$ for $k = 1, 2, \dots, n + 2$. The z'_k s are uniquely determined. Now let

$$\Phi_k = z_k^{-n/2} \prod_{\substack{j=1 \\ j \neq k}}^{n+2} [(z_k z_j - a) / |z_k z_j - a|] \text{ for}$$

$k = 1, 2, \dots, n + 2$ where $0 \leq \arg z^{1/2} < \pi$.

THEOREM 1. *If f is continuous on E_a and p_n is a polynomial of degree n , $n \in Z^+$, then $f \in \mathcal{F}_n(E_a)$ and $p_n = p_n(f, E_a)$ if and only if there exists $n + 2$ points $\{\xi_k\}_{k=1}^{n+2}$ in E_a , with $0 \leq \text{Arg } \xi_1 < \text{Arg } \xi_2 < \dots < \text{Arg } \xi_{n+2} < 2\pi$ if $a \neq 1$ or $-2 \leq \xi_1 < \xi_2 < \dots < \xi_{n+2} \leq 2$ if $a = 1$, where $|f(\xi) - p_n(\xi)|$ attains its maximum on E_a and, $[f(\xi_i) - p_n(\xi_i)]/\Phi_i = -[f(\xi_{i+1}) - p_n(\xi_{i+1})]/\Phi_{i+1}$ for $i = 1, 2, \dots, n + 1$ where the Φ_i 's are defined in terms of the ξ_i 's as above.*

Proof. In order to prove our theorem we make use of a lemma which is a reformulation of a result [2] due to T. S. Motzkin and J. L. Walsh.

LEMMA. *A necessary and sufficient condition that the given numbers $\{\sigma_k\}_{k=1}^{n+2}$ be the deviations of some function f defined on the $n + 2$ points $\{\xi_k\}_{k=1}^{n+2}$ and its polynomial of degree n of best uniform approximation to f on these points is that for some $\rho \geq 0$;*

- (1) $|\sigma_k| = \rho$ for $k = 1, 2, \dots, n + 2$ and,
- (2) $\arg \sigma_k = \arg \omega'(\xi_k) + \theta_0$ for $k = 1, 2, \dots, n + 2$ if $\rho > 0$ where

$$\omega(\xi) = \prod_{k=1}^{n+2} (\xi - \xi_k) \text{ and } \theta_0 = \arg \left[\sum_{k=1}^{n+2} f(\xi_k)/\omega'(\xi_k) \right].$$

The necessary portion of our theorem will then follow if it is shown that;

$$(2.1) \quad \arg\{[\omega'(\xi_i)/\Phi_i]/[\omega'(\xi_{i+1})/\Phi_{i+1}]\} = \pi \text{ for}$$

$i = 1, 2, \dots, n + 1$. Now substituting $z_j + a/z_j$ for ξ_j and using the definition of the Φ_j 's we can show the (2.1) is equivalent to;

$$(2.2) \quad \arg\{(z_{i+1}^{n/2}/z_i^{n/2}) \prod_{\substack{j=i, i+1 \\ j \neq i, i+1}}^{n+2} [(z_i - z_j)/(z_{i+1} - z_j)]\} = 0.$$

But, (2.2) follows since z_i and z_{i+1} are by virtue of their definition adjacent on the unit circle U (i.e., z_i and z_{i+1} are on a connected arc in U containing none of the other z_j 's) and since; $\arg (z_{i+1}/z_i) = -2 \arg [z_i - z_j]/(z_{i+1} - z_j)$ for $j \neq i, i + 1$.

In order to prove the converse of our theorem we simply work backwards and show that; $\arg [f(\xi_k) - P_n(\xi_k)] = \arg \omega'(\xi_k) + \theta_0$ for some θ_0 and $k = 1, 2, \dots, n + 2$ and apply the aforementioned result of Motzkin and Walsh.

3. Special cases and applications. Chebyshev's Alternation Theorem follows as a special case of Theorem 1, when $a = 1$, since it is known [1, p. 22] that all real functions, not themselves polynomials of degree n , continuous on $[-2, 2]$ are in the class $\mathcal{F}_n([-2, 2])$.

Also of interest because of its simple form is the case where $a = 0$ or $E_a = U$ is the unit circle and where n is even. In this case our main theorem appears to provide us with a valuable tool in determining if a given function f is in $\mathcal{F}_{2m}(U)$ and if it is, in finding $p_{2m}(f, U)$.

COROLLARY 1. *If f is continuous on U and p_{2m} is a polynomial of degree $2m$, $m \in \mathbb{Z}^+$, then $f \in \mathcal{F}_{2m}(U)$ and $p_{2m} = p_{2m}(f, U)$ if and only if there exists $2m + 2$ points, $\{z_k\}_{k=1}^{2m+2}$, with $0 \leq \text{Arg } z_1 < \dots < \text{Arg } z_{2m+2} < 2\pi$ where $|f(z) - p_{2m}(z)|$ attains its maximum on U and where $[f(z_k) - p_{2m}(z_k)]/z_k^m = -[f(z_{k+1}) - p_{2m}(z_{k+1})]/z_{k+1}^m$, for $k = 1, 2, \dots, 2m + 1$.*

Corollary 1 can be used to obtain a recently discovered example of best approximation [3], namely, if $f(z) = (\alpha z + \beta)/(z - a)(1 - \bar{a}z)$, $|a| > 1$, then;

$$p_{2m}(f, U)(z) = [\alpha z + \beta - K_1 z^{2m}(1 - \bar{a}z)^2 - K_2(z - a)^2]/(z - a)(1 - \bar{a}z),$$

where

$$K_1 = (\alpha a + \beta)/a^{2m}(1 - |a|^2)^2$$

and,

$$K_2 = \bar{a}(\alpha + \beta \bar{a})/(1 - |a|^2)^2.$$

REFERENCES

1. G. G. Lorentz, *Approximation of Functions*, Holt, Rinehart and Winston, 1966.
2. T. S. Motzkin and J. L. Walsh, *On the derivative of a polynomial and Chebyshev approximation*, Proc. Amer. Math. Soc., **4** (1953), 76-87.
3. S. J. Poreda, *An example of best uniform approximation in the complex plane*, Notices Amer. Math. Soc., **17** (1970), 569.
4. T. J. Rivlin, *Some explicit polynomial approximations in the complex domain*, Bull. Amer. Math. Soc., **73** (1967), 467-469.
5. T. J. Rivlin and H. S. Weiss, *Some best polynomial approximations in the plane*, Duke Math. J., **35** (1968), 475-482.

Received December 4, 1970. This work represents part of the author's doctoral dissertation that was directed by Professor Mishael Zedek at the University of Maryland.

CLARK UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

| | |
|---|-----|
| Alex Bacopoulos and Athanassios G. Kartsatos, <i>On polynomials approximating the solutions of nonlinear differential equations</i> | 1 |
| Monte Boisen and Max Dean Larsen, <i>Prüfer and valuation rings with zero divisors</i> | 7 |
| James J. Bowe, <i>Neat homomorphisms</i> | 13 |
| David W. Boyd and Hershy Kisilevsky, <i>The Diophantine equation $u(u + 1)(u + 2)(u + 3) = v(v + 1)(v + 2)$</i> | 23 |
| George Ulrich Brauer, <i>Summability and Fourier analysis</i> | 33 |
| Robin B. S. Brooks, <i>On removing coincidences of two maps when only one, rather than both, of them may be deformed by a homotopy</i> | 45 |
| Frank Castagna and Geert Caleb Ernst Prins, <i>Every generalized Petersen graph has a Tait coloring</i> | 53 |
| Micheal Neal Dyer, <i>Rational homology and Whitehead products</i> | 59 |
| John Fuelberth and Mark Lawrence Teply, <i>The singular submodule of a finitely generated module splits off</i> | 73 |
| Robert Gold, <i>Γ-extensions of imaginary quadratic fields</i> | 83 |
| Myron Goldberg and John W. Moon, <i>Cycles in k-strong tournaments</i> | 89 |
| Darald Joe Hartfiel and J. W. Spellmann, <i>Diagonal similarity of irreducible matrices to row stochastic matrices</i> | 97 |
| Wayland M. Hubbart, <i>Some results on blocks over local fields</i> | 101 |
| Alan Loeb Kostinsky, <i>Projective lattices and bounded homomorphisms</i> | 111 |
| Kenneth O. Leland, <i>Maximum modulus theorems for algebras of operator valued functions</i> | 121 |
| Jerome Irving Malitz and William Nelson Reinhardt, <i>Maximal models in the language with quantifier "there exist uncountably many"</i> | 139 |
| John Douglas Moore, <i>Isometric immersions of space forms in space forms</i> | 157 |
| Ronald C. Mullin and Ralph Gordon Stanton, <i>A map-theoretic approach to Davenport-Schinzel sequences</i> | 167 |
| Chull Park, <i>On Fredholm transformations in Yeh-Wiener space</i> | 173 |
| Stanley Poreda, <i>Complex Chebyshev alterations</i> | 197 |
| Ray C. Shiflett, <i>Extreme Markov operators and the orbits of Ryff</i> | 201 |
| Robert L. Snider, <i>Lattices of radicals</i> | 207 |
| Ralph Richard Summerhill, <i>Unknotting cones in the topological category</i> | 221 |
| Charles Irvin Vinsonhaler, <i>A note on two generalizations of QF-3</i> | 229 |
| William Patterson Wardlaw, <i>Defining relations for certain integrally parameterized Chevalley groups</i> | 235 |
| William Jennings Wickless, <i>Abelian groups which admit only nilpotent multiplications</i> | 251 |