

Pacific Journal of Mathematics

BAXTER'S THEOREM AND VARBERG'S CONJECTURE

BENNETT EISENBERG

BAXTER'S THEOREM AND VARBERG'S CONJECTURE

BENNETT EISENBERG

It is shown that Baxter's strong law for Gaussian processes provides necessary and sufficient conditions for equivalence for a wide class of Gaussian processes.

Let X be a Gaussian process with zero mean and covariance $R(s, t) = E(X(s)X(t))$. Baxter [1] proved that if R is continuous for $0 \leq s, t \leq 1$ with uniformly bounded second derivatives for $s \neq t$, then

$$D^+(t) = \lim_{s \downarrow t} \frac{R(s, t) - R(t, t)}{s - t}$$

and

$$D^-(t) = \lim_{s \uparrow t} \frac{R(t, t) - R(s, t)}{t - s}$$

exist and are continuous and with probability one

$$\lim_{n \rightarrow \infty} \Sigma \left[x\left(\frac{k}{2^n}\right) - x\left(\frac{k-1}{2^n}\right) \right]^2 = \int_0^1 D^-(t) - D^+(t) dt .$$

It follows that if Y is another Gaussian process with mean zero and covariance S continuous with bounded second derivatives for $s \neq t$ and if there exists a t with

$$D_{\bar{R}}^-(t) - D_{\bar{R}}^+(t) \neq D_{\bar{S}}^-(t) - D_{\bar{S}}^+(t)$$

then the measures μ_x and μ_y for the processes X are singular.

In the case where R and S are triangular covariances Varberg [8] has obtained a converse to this result. Varberg's Theorem:

$$\text{Let } R_i(s, t) = \begin{cases} u_i(s)v_i(t) & s \leq t \\ v_i(s)u_i(t) & s \geq t \end{cases},$$

where $i = 1$ or 2 .

Assume furthermore

- (A) $u_i(0) = 0$
- (B) $v_i(t) > 0$ on $[0, T]$
- (C) u_i' and v_i' exist and are continuous on $[0, T]$
- (D) $v_i(t)u_i'(t) - u_i(t)v_i'(t) > 0$ on $[0, T]$

then μ_x and μ_y are mutually absolutely continuous if and only if

$$v_1(t)u_1'(t) - u_1(t)v_1'(t) = v_2(t)u_2'(t) - u_2(t)v_2'(t)$$

for $t \in [0, T]$

Otherwise $\mu_x \perp \mu_y$.

Condition A says that both processes start at 0. This is no loss in generality since Baxter's theorem involves only the increments of the processes. For a triangular covariance R ,

$$D_R^-(t) - D_R^+(t) = v(t)u'(t) - u(t)v'(t)$$

so that Condition D means $D_R^-(t) - D_R^+(t) > 0$. Condition C is slightly stronger than Baxter's regularity condition on R .

Varberg conjectured that for two arbitrary Gaussian processes with covariances R and S satisfying regularity and boundary conditions of the type $A - D$, a necessary and sufficient condition for equivalence is that $D_R^-(t) - D_R^+(t) = D_S^-(t) - D_S^+(t)$.

In this paper we answer Varberg's conjecture. We also extend his theorem by requiring only one continuous derivative for each of the functions u and v .

1. **Varberg's conjecture.** Throughout this paper X and Y are Gaussian processes with mean zero and covariances R and S , respectively. μ_x and μ_y denote the corresponding measures on the sample paths.

Write $X \sim Y$ if μ_x and μ_y are mutually absolutely continuous over a given sigma field. Write $X \perp Y$ if μ_x and μ_y are singular.

Gaussian processes are known to be either equivalent or singular over the sigma-fields generated by x_t , for t in some parameter set. The Segal-Feldman [4, 6] equivalence conditions take the form:

Let H_x be the Hilbert space spanned by X_t in $L^2(d\mu_x)$ for t in A . Let H_y be the Hilbert space spanned by Y_t in $L^2(d\mu_y)$, for t in A . Then $X \sim Y$ if and only if the map $T: X_t \rightarrow Y_t$ extends to a linear homeomorphism between H_x and H_y (In this case we write $H_x \approx H_y$) and $I - T^*T$ is Hilbert Schmidt from H_x to H_x .

Otherwise $X \perp Y$.

In terms of covariance functions the conditions become [3] $X \sim Y$, if and only if

$R(s, t) - S(s, t) = E(HX_s X_t)$, where H is a Hilbert Schmidt operator on H_x with $I - H$ invertible.

In place of the invertibility condition we also have [3] $X \sim Y$ if and only if $H_x \approx H_y$ and

$$R(s, t) - S(s, t) = E(HX_s X_t), \text{ where } H \text{ is Hilbert Schmidt on } H_x.$$

For a quick application let X be an additive Gaussian process.

PROPOSITION 1. *Assume X has independent increments with $E(X_t^2) = F(t)$. Then $X \sim Y$ on $[0, T]$ if and only if*

$F(\min(s, t)) - S(s, t) = \int_{0^-}^s \int_{0^-}^t H(u, v) dF(d) dF(v)$, where H is a Hilbert Schmidt kernel on $L^2([0, T], dF)$ with $I - H$ invertible. Notation: \int_{0^-} signifies that the mass $F(0)$ at 0 is included in the integration.

Proof. $X_t \rightarrow c_{0t}$ in $L^2(dF)$ is an isometry of H_x to $L^2([0, t], dF)$; where C_{0t} is the characteristic function of $[0, T]$.

Thus $X \sim Y$ if and only if

$$F(\min(s, t)) - S(s, t) = E(HX_s X_t) \\ = \int_{0^-}^T \int_{0^-}^T \tilde{H}(u, v) C_{0s}(u) C_{0s}(v) dF(u) dF(v) = \int_{0^-}^s \int_{0^-}^t \tilde{H}(u, v) dF(u) dF(v)$$

where H is unitarily equivalent to \tilde{H} .

This generalizes Shepp's theorem [7] where X is Brownian motion and dF is Lebesgue measure.

COROLLARY. *If X is an additive process with $x(0) \equiv 0$, and $F(\min(s, t)) - S(s, t) = \int_0^s \int_0^t H(u, v) dF(u) dF(v)$ for $0 \leq s, t \leq T$, then there exists an interval where $X \sim Y$.*

Proof. If we consider a smaller interval $T' \subset T$

$$F(\min(s, t)) - S(s, t) = \int_0^s \int_0^t H(u, v) dF(u) dF(v)$$

for $0 \leq s, t \leq T'$ and the same function H . Moreover, as $T' \rightarrow 0$, the Hilbert-Schmidt norm of H acting on $L^2([0, T'], dF)$ approaches zero. Hence the operator norm of H approaches zero. But $\|H\| < 1 \Rightarrow I - H$ is invertible.

After this corollary the question might be raised, "Are there weaker conditions which imply that Gaussian processes are equivalent over the sigma field $\bigcap_{n=1}^\infty B_{1/n}$ where $B_{1/n}$ is generated by $x_t, 0 \leq t \leq 1/n$?" The answer is negative.

PROPOSITION 2. *Gaussian processes are singular or equivalent locally. They are equivalent locally if and only if there exists an interval of equivalence.*

Proof. Assume there exists an interval of equivalence. Then a fortiori the processes are locally equivalent.

Assume there does not exist an interval of equivalence. Then

the processes are singular over $B_{1/n}$ for every n . We may thus choose A_n in $B_{1/n}$ with $\mu_x(A_n) = 1$ and $\mu_y(A_n) = 0$. $A = \bigcap_{m=1}^\infty \bigcup_{n=m}^\infty A_n$ lies in $\bigcap B_{1/n}$ and $\mu_x(A) = 1$ while $\mu_y(A) = 0$. Hence the processes are singular locally.

This proposition implies that two processes share the same strong laws locally if and only if there exists an interval of equivalence.

Next assume both X and Y are additive processes with $E(X_t^2) = F(t)$ and $E(Y_t^2) = G(t)$. To keep the discussion focused on Baxter's theorem assume $F(0) = G(0) = 0$ and F and G are each absolutely continuous with $F' > 0$ and $G' > 0$. Baxter's theorem would say that if F'' and G'' are bounded, then unless $F' \equiv G'$ (which implies $F \equiv G$) the processes would be singular. A theorem of Feldman extends this idea to general F and G .

PROPOSITION 3 (Feldman) [5]: *Assume X and Y are additive processes with $F(0) = G(0) = 0$. Assume F and G are each absolutely continuous with $F' > 0$ and $G' > 0$.*

Then $X \sim Y$ if and only if $F(t) \equiv G(t)$.

Feldman used general F and G in his original theorem.

Proof. For completeness we prove this result, this time using the original Segal-Feldman equivalence conditions. (R. M. Dudley suggested this.)

$x_t \rightarrow C_{0t}$ is an isometry of H_x and $L^2(dF)$

$y_t \rightarrow C_{0t}(dG/dF)^{1/2}$ is an isometry of H_y and $L^2(dF)$.

$T: x_t \rightarrow y_t$ is then unitarily equivalent to $\tilde{T}: C_{0t} \rightarrow C_{0t}(dG/dF)^{1/2}$ in $L^2(dF)$.

$I - \tilde{T}^* \tilde{T}$ is then multiplication by $1 - (dG/dF)$

which is Hilbert Schmidt if and only if $(dG/dF) \equiv 1$ (and in this case $I - H = I$ is invertible).

Thus Baxter's theorem determines equivalence conditions for some processes without bounded second derivatives.

Next we answer Varberg's conjecture.

PROPOSITION 4. *Let X have covariance R , where*

(A') $R(0, 0) = 0$

(C') $(\partial^2 R / \partial s \partial t)$ exists and is continuous for $s \neq t$ and $\lim_{s=t} (\partial^2 R / \partial s \partial t)$ exists for all s .

(D') $D_R^-(t) - D_R^+(t) > 0$.

Then there is an additive process Y with X equivalent to Y

on some interval. Moreover, $D_R^- - D_R^+ = D_S^- - D_S^+$.

Before proving the proposition we need some real variable lemmas.

LEMMA. Assume $R(s, t)$ defined and continuous for $0 \leq s \leq t$. Assume $(\partial^2 R / \partial t \partial s)$ is continuous for $s < t$ and $\lim_{s=t} (\partial^2 R / \partial t \partial s)$ exists. Then

$$\frac{\partial}{\partial t^+} \frac{\partial R}{\partial s^-} \Big|_{s=t} = \lim_{s=t} \frac{\partial^2 R}{\partial t \partial s}.$$

Proof.

$$\begin{aligned} \int_T^{t'} \int_0^{s'} \frac{\partial^2 R}{\partial s \partial t} ds dt &= \int_T^{t'} \frac{\partial R}{\partial t}(s', t) - \frac{\partial R}{\partial t}(0, t) dt \\ &= R(s', t') - R(s', T) - R(0, t') + R(0, T). \end{aligned}$$

Hence

$$R(s, t) = \int_T^t \int_0^s (\partial^2 R / \partial s \partial t) ds dt + R(s, T) + R(0, t) - R(0, T).$$

And by continuity of R and the assumption this holds for $s = t$ as well as $s < t$. Hence $(\partial^2 R / \partial t^+ \partial s^-) = \lim (\partial^2 R / \partial s \partial t)$.

COROLLARY. If R is continuous for $0 \leq S, t \leq T$ and if $(\partial^2 R / \partial s \partial t)$ is continuous for $s \neq t$ and has limits at $s = t$, then

$$\frac{\partial}{\partial t^+} \frac{\partial R}{\partial s^-} \Big|_{s=t} = \frac{\partial}{\partial t^-} \frac{\partial R}{\partial s^+} \Big|_{s=t} = \lim_{s=t} \frac{\partial^2 R}{\partial s \partial t}.$$

In particular, if $(\partial R / \partial s^-) |_{s=t} = (\partial R / \partial s^+) |_{s=t}$ then $(\partial^2 R / \partial s \partial t)$ exists at $s = t$ and is continuous for $0 \leq s, t \leq T$.

Note that if R is symmetric and $\lim_{s=t} (\partial^2 R / \partial s \partial t)$ exists from each side of the diagonal, then the limits must be the same. Proof of Proposition 4:

Let $f(t) = D_R^-(t) - D_R^+(t)$. Then f is continuous. Let $E(Y_t^2) = F(t) = \int_0^t f(u) du$. Then by (A'), and the corollary to Proposition 1, $X \sim Y$ on some interval if and only if $F(\min(s, t)) - R(s, t) = \int_0^s \int_0^t H(u, v) f(u) f(v) dudv$, where H is in L^2 .

For $s \neq t$ we have $(\partial^2 (F(\min(s, t)) - R(s, t)) / \partial s \partial t)$ exists and is continuous.

On the diagonal $s = t$,

$$\frac{\partial}{\partial s^+} (F(\min(s, t)) - R(s, t)) \Big|_{s=t} = - \frac{\partial R}{\partial s}(s, t) \Big|_{s=t}$$

and

$$\frac{\partial}{\partial s^-} (F(\min(s, t)) - R(s, t)) \Big|_{s=t} = f(s) - \frac{\partial R}{\partial s^-} \Big|_{s=t} .$$

By definition $f(t) = (\partial R/\partial s^-)(t, t) - (\partial R/\partial s^+)(t, t)$.

Thus $(\partial(F(\min(s, t)) - R(s, t))/\partial s)$ is continuous everywhere. For $s \neq t$, $(\partial^2(F(\min(s, t)) - R(s, t))/\partial t \partial s) = -(\partial^2 R/\partial t \partial s)$. Using hypothesis C' and applying the corollary of the lemmas $(\partial^2(F(\min(s, t)) - R(s, t))/\partial t \partial s)$ exists at $s = t$ and equals $\lim_{s \rightarrow t} (\partial^2 R/\partial t \partial s)$. Hence, it is a continuous function everywhere. Since $f(u) \neq 0$ by (D') we may write

$$H(u, v) = \frac{\partial^2(F(\min(u, v)) - R(u, v))}{\partial u \partial v} (f(u)f(v)).$$

By the corollary to Proposition 1 there is an interval of equivalence of X and Y .

COROLLARY. *If R and S satisfy A', C' and D' then there exists an interval where $X \sim Y$ if and only if there is an interval with $D_R^- - D_R^+ = D_S^- - D_S^+$.*

Proof. They are both equivalent to the same additive process on some interval.

The strict positivity of $D_R^- - D_R^+$ is essential since these conditions are not sufficient for the equivalence of differentiable processes, for example. It would be nice to remove the phrase "there exists an interval", but this cannot be done without complicated positive definiteness conditions on R and S . Consider the case of the Brownian motion with covariance $\min(s, t)$ and the Brownian bridge with $S(s, t) = \begin{cases} s(1-t) & s \leq t \\ t(1-s) & s > t \end{cases}$. Both covariances start at zero, are smooth and satisfy $D^-(t) - D^+(t) \equiv 1$. Still the processes are singular over $[0, T]$, $T \geq 1$. Varberg handles this situation by Condition B, that $v_i(t) > 0$. Unfortunately there is no corresponding condition for general covariance functions.

2. Equivalent processes with triangular covariance functions. Next we extend Varberg's theorem. We assume u and v each have but one bounded continuous derivative.

PROPOSITION 5. *Let X have covariance function*

$$R(s, t) = \begin{cases} u(s)v(t) & s \leq t \\ v(s)u(t) & t < s \end{cases}$$

where $v > 0$, (u/v) is right continuous and increasing. Then $X \sim Y$ over $[0, T]$ if and only if

$$R(s, t) - S(s, t) = v(s)v(t) \int_0^s \int_0^t H(u, v) d\left(\frac{u}{v}\right) d\left(\frac{u}{v}\right),$$

where H is a Hilbert Schmidt kernel on $L^2([0, T], d(u/v))$ with $I - H$ invertible.

Proof. Since $v > 0$, $X \sim Y$ if and only if $X/v \sim Y/v$. But X/v is an additive process with covariance (u/v) ($\min(s, t)$). The result then follows from Proposition 1.

COROLLARY. Assume $R_i(s, t) = \begin{cases} u_i(s) v_i(t) & s \leq t \\ v_i(s) u_i(t) & t < s \end{cases}, i = 1, 2.$

Assume A, B , and D , but in place of C assume that u'_i and v'_i are continuous.

Then $X_1 \sim X_2$ on $[0, T]$ if and only if

$$(*) \quad u'_1(t)v_1(t) - u_1(t)v'_1(t) \equiv u'_2(t)v_2(t) - u_2(t)v'_2(t).$$

Proof. Assume $(*)$ holds. Then for $s \neq t$, $(\partial(R_1 - R_2)/\partial s)$ exists and is continuous.

At $s = t$ we have $(\partial/\partial s^+) (R_1 - R_2)|_{s=t} = u_1(t)v'_1(t) - u_2(t)v'_2(t)$

$$\frac{\partial(R_1 - R_2)}{\partial s^-} \Big|_{s=t} = u'_1(t)v_1(t) - u'_2(t)v_2(t).$$

By assumption these are equal.

This time the factorability makes checking the continuity of $(\partial^2(R_1 - R_2)/\partial t \partial s)$ easy.

Thus $R_1(s, t) - R_2(s, t) = E(H_1 X_1(S) X_1(t))$ and $R_1(s, t) - R_2(s, t) = E(H_2 X_2(S) X_2(t))$.

Both relations together imply $H_x \approx H_y$ (see the lemma which follows) so that $X \sim Y$.

Unlike the case in Varberg's proof the more difficult half is to prove that if the processes are equivalent the Baxter condition holds. (Baxter's theorem does not apply here.)

However, we now have

$$\frac{R_1(s, t) - R_2(s, t)}{v(s)v(t)} = \int_0^t \int_0^s H(x, y) d\left(\frac{u}{v}\right) d\left(\frac{u}{v}\right) = \int_0^t \int_0^s \hat{H} \, dx dy.$$

By conditions on R_1 and R_2

$(\partial^2(R_1 - R_2)/\partial s \partial t)$ is continuous and bounded for $s \neq t$.

Hence the same holds for $R_1(s, t) - R_2(s, t)/v(s)v(t)$. Since

$$\hat{H} = \frac{\partial^2}{\partial s \partial t} \left(\frac{R_1(s, t) - R_2(s, t)}{v(s)v(t)} \right) \text{ a.e.,}$$

we may assume \hat{H} is continuous and bounded for $s \neq t$. It is then an easy argument to show that $\int_0^t \hat{H}(x, y) dx$ is a continuous function of y for each t .

Hence $(\partial/\partial s) \int_0^s \int_0^t H(x, y) dx dy$ is a continuous function of s . But

$$R_1(s, t) - R_2(s, t) = v(s)v(t) \int_0^s \int_0^t H(x, y) dx dy$$

so $\partial R_1(s, t) - R_2(s, t)/\partial s|_{s=t}$ must exist. Hence (*) holds.

LEMMA. If $R(s, t) - S(s, t) = E(H_1 X_s X_t)$ and $S(s, t) - R(s, t) = E(H_2 Y_s Y_t)$, where H_1 and H_2 are Hilbert-Schmidt then $H_x \approx H_y$.

Proof.

$$S(s, t) = E(Y_s Y_t) = E((I - H) X_s X_t)$$

$$\|\Sigma C_k Y_{t_k}\|^2 = E((I - H)(\Sigma C_k X_{t_k}) \Sigma C_k X_{t_k}) \leq k_1 \|\Sigma C_k X_{t_k}\|^2.$$

Similarly,

$$\|\Sigma C_k X_{t_k}\|^2 \leq k_2 \|\Sigma C_k Y_{t_k}\|^2.$$

Recently, Yeh [9] has found singularity conditions of the Baxter type using different assumptions on u and v .

In the case where both processes are stationary Belayev has proved a generalized Baxter result: If $E((x_i(h) - x_i(0))^2) = \phi_i(h)$ and if $\lim_{h \rightarrow 0} h^2/\phi_i(h) = 0$, then $X_1 \perp X_2$ unless $\lim_{h \rightarrow 0} (\phi_1(h)/\phi_2(h)) = 1$.

The idea of relating a process to an equivalent additive or otherwise simple process could be useful in understanding these results as well as in extending many strong laws more easily, proved for the simpler processes.

Acknowledgement. I would like to thank Professors Ray and Dudley for their advice and criticism while I was at M.I.T.

REFERENCES

1. Glen Baxter, *A Strong Limit Theorem for Gaussian Processes*, Proc. Amer. Math. Soc., **7** (1956), 522-527.
2. Yu K. Belyayev, *Continuity and Holders Conditions for Sample Functions of Stationary Gaussian Processes*, Fourth Berkeley Symposium on Probability and Statistics, Vol 2, 23-32.
3. Bennett Eisenberg, *The relation of the equivalence conditions for the Brownian motion to the equivalence conditions for certain stationary processes*, Ann. Math. Stat., **40** (1969), 2143-2148.
4. Jacob Feldman, *Equivalence and perpendicularity of Gaussian processes*, Pacific J. Math., **8** (1958), 699-708.
5. ———, *Some classes of equivalent Gaussian processes, on an interval*, Pacific J. Math., **10** (1960), 1211-1220.
6. I. E. Segal, *Distributions in Hilbert space and canonical systems of operators*, Trans. Amer. Math. Soc., **88** (1958), 12-41.
7. Lawrence Shepp, *Radon-Nikodym derivatives of Gaussian measures*, Ann. Math. Stat., **37** (1966), (321-354).
8. Dale E. Varberg, *On equivalence of Gaussian measures*, Pacific J. Math., **11** (1961), 751-762.
9. J. Yeh, *Singularity of Gaussian measures on function spaces with factorable covariance functions*, Pacific J. Math., **31** (1969), 547-554.

Received November 30, 1970. Part of this work is in a Ph. D. thesis written while the author was supported by a National Science Foundation Fellowship at the Massachusetts Institute of Technology. The rest was done on a National Science Foundation Post-Doctoral Fellowship at the University of New Mexico.

UNIVERSITY OF NEW MEXICO

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Louis I. Alpert and L. V. Toralballa, <i>An elementary definition of surface area in E^{n+1} for smooth surfaces</i>	261
Eamon Boyd Barrett, <i>A three point condition for surfaces of constant mean curvature</i>	269
Jan-Erik Björk, <i>On the spectral radius formula in Banach algebras</i>	279
Peter Botta, <i>Matrix inequalities and kernels of linear transformations</i>	285
Bennett Eisenberg, <i>Baxter's theorem and Varberg's conjecture</i>	291
Heinrich W. Guggenheimer, <i>Approximation of curves</i>	301
A. Hedayat, <i>An algebraic property of the totally symmetric loops associated with Kirkman-Steiner triple systems</i>	305
Richard Howard Herman and Michael Charles Reed, <i>Covariant representations of infinite tensor product algebras</i>	311
Domingo Antonio Herrero, <i>Analytic continuation of inner function-operators</i>	327
Franklin Lowenthal, <i>Uniform finite generation of the affine group</i>	341
Stephen H. McCleary, <i>0-primitive ordered permutation groups</i>	349
Malcolm Jay Sherman, <i>Disjoint maximal invariant subspaces</i>	373
Mitsuru Nakai, <i>Radon-Nikodým densities and Jacobians</i>	375
Mitsuru Nakai, <i>Royden algebras and quasi-isometries of Riemannian manifolds</i>	397
Russell Daniel Rupp, Jr., <i>A new type of variational theory sufficiency theorem</i>	415
Helga Schirmer, <i>Fixed point and coincidence sets of biconnected multifunctions on trees</i>	445
Murray Silver, <i>On extremal figures admissible relative to rectangular lattices</i>	451
James DeWitt Stein, <i>The open mapping theorem for spaces with unique segments</i>	459
Arne Stray, <i>Approximation and interpolation</i>	463
Donald Curtis Taylor, <i>A general Phillips theorem for C^*-algebras and some applications</i>	477
Florian Vasilescu, <i>On the operator $M(Y) = TYS^{-1}$ in locally convex algebras</i>	489
Philip William Walker, <i>Asymptotics for a class of weighted eigenvalue problems</i>	501
Kenneth S. Williams, <i>Exponential sums over $GF(2^n)$</i>	511