

# Pacific Journal of Mathematics

**APPROXIMATION OF CURVES**

HEINRICH W. GUGGENHEIMER

## APPROXIMATION OF CURVES

H. GUGGENHEIMER

**Generalizing recent results of J. M. Sloss we show: A curve in  $n$ -space that admits a continuously differentiable first order frame can be  $C^1$  approximated to any desired accuracy by a continuous, piecewise  $C^{r+2}$  curve for which the curvature functions are prescribed  $C^r$  ( $r = 0, 1, \dots, \infty, \omega$ ) functions of the arc length. The result can be extended to riemannian geometry.**

The theorem published by James M. Sloss [4] in this journal to the effect that a regular  $C^3$  curve can be approximated by a piecewise helix that either is circular or whose curvature or torsion are those of the given curve, can be generalized. The proof can be simplified by an explicit use of Gronwall's inequality [2] which is one of the strongest tools available in the treatment of ordinary differential equations. Gronwall's inequality says that

$$u(t) \leq C + \int_a^t u(\tau)v(\tau)d\tau, \quad t \geq a, \quad u(t) \geq 0, \quad v(t) \geq 0,$$

implies

$$u(t) \leq C \exp \int_a^t v(\tau)d\tau.$$

We fix a coordinate system in  $E^n$ . Let  $a_i$  be the unit vector on the  $i$ th coordinate axis and

$$a = \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix}$$

the frame of these vectors. We are given a curve  $X(s)$ ,  $0 \leq s \leq L$ , as a function of its arc length. We say that  $X$  admits a continuously differentiable frame of first order [1] if there exists a continuously differentiable orthogonal matrix function  $\Theta(s)$  so that the unit vector  $e_1(s) = X'(s)$  is the first vector of the frame  $e = \Theta a$ . Then  $X$  must be a  $C^2$  curve. Conversely, if  $X$  is  $C^2$ , we may define  $e$  as the frame on  $S^{n-1}$  for which  $e_1$  is the unit tangent  $e_1(s) \in S^{n-1}$   $e_k$  ( $1 \leq k \leq n-1$ ) is tangent to the equatorial  $S^{k-1}$  in the  $S^k$  defined by  $a_1, \dots, a_{k+1}$  and  $e_n$  is tangent to the meridian. Naturally, if  $X$  is  $C^{n+1}$ , the Frenet frame may be taken as  $e$ . We put  $A(s) = \Theta'(s)\Theta(s)^{-1}$ .

For matrices we use the sup-norm of the corresponding linear

transformation. The norm of the frame  $e = \Theta a$  shall be the norm of  $\Theta$ .

**THEOREM.** *A  $C^2$  curve of finite length can be approximated in the  $C^1$  topology by a continuous, piecewise differentiable curve for which the curvature functions are prescribed  $C^r$  functions of the arc length,  $r \geq 0$ .*

The structure equations of the curve  $X$  and the frame  $e$  are

$$\begin{aligned} X'(s) &= e_1(s) , \\ e'(s) &= A(s)e(s) , \end{aligned}$$

with a continuous matrix  $A(s)$ ,  $0 \leq s \leq L$ . We want to approximate  $X(s)$  by a continuous curve  $Y(s)$  so that

$$\begin{aligned} Y'(s) &= e_1^*(s) , \\ e^{*'}(s) &= A^*(s)e^*(s) , \end{aligned}$$

except at finitely many points where the equations hold only in the sense of forward one-sided derivatives and where  $A^*(s)$  is a given (continuous) matrix function of  $s$ . We shall not use the fact that  $A^*(s)$  is skew-symmetric and has nonzero entries  $a_{ij}^*$  only for  $j = i \pm 1$  [3].

We put

$$M = \max \|A(s)\|, N = \max \|A^*(s)\| ,$$

and choose  $\Delta s$  subject to

$$\Delta s \leq \min (N^{-1}, \varepsilon(e(1 + L)(M + N))^{-1}), \frac{L}{\Delta s} \text{ an integer .}$$

For  $k\Delta s \leq s \leq (k + 1)\Delta s$ ,  $0 \leq k < L/\Delta s$ , define  $Y(s)$  by

$$\begin{aligned} e^*(s) &= e(k\Delta s) + \int_{k\Delta s}^s A^*(\sigma)e^*(\sigma)d\sigma , \\ Y(s) &= Y(k\Delta s) + \int_{k\Delta s}^s e_1^*(\sigma)d\sigma , \\ Y(0) &= X(0) . \end{aligned}$$

Then

$$\begin{aligned} \|e^*(s) - e(s)\| &= \left\| \int_{k\Delta s}^s (A^*e^* - Ae)d\sigma \right\| \\ &\leq \int_{k\Delta s}^s \|A^* - A\| \|e\| d\sigma + \int_{k\Delta s}^s \|A^*\| \|e^* - e\| d\sigma \end{aligned}$$

$$\leq (M + N)\Delta s + N \int_{k\Delta s}^s \|e^* - e\| d\sigma .$$

By Gronwall's inequality,

$$\begin{aligned} \|e^*(s) - e(s)\| &\leq (M + N)\Delta s \exp(N(s - k\Delta s)) \\ &\leq (M + N)e\Delta s \leq \varepsilon/(1 + L). \end{aligned}$$

Hence,

$$\begin{aligned} |Y(s) - X(s)| &\leq |Y(k\Delta s) - X(k\Delta s)| + \int_{k\Delta s}^s |e_1^*(\sigma) - e_1(\sigma)| d\sigma \\ &\leq |Y(k\Delta s) - X(k\Delta s)| + \int_{k\Delta s}^s \|e^*(\sigma) - e(\sigma)\| d\sigma \\ &\leq |Y(k\Delta s) - X(k\Delta s)| + \varepsilon\Delta s(L + 1)^{-1}. \end{aligned}$$

By induction,  $|Y(k\Delta s) - X(k\Delta s)| \leq \varepsilon k\Delta s(1 + L)^{-1}$  and

$$\begin{aligned} |Y(s) - X(s)| &\leq \varepsilon L(1 + L)^{-1}, \\ |Y(s) - X(s)| + |Y'(s) - X'(s)| &\leq \varepsilon \frac{L}{1 + L} + \varepsilon \frac{1}{1 + L} = \varepsilon. \end{aligned}$$

The same argument would hold for a matrix  $A^*(s)$  with bounded, integrable entries if the differential equations are supposed to hold a.e., in the sense of Caratheodory. For  $C^r$  entries,  $Y$  is a  $C^{r+2}$  curve. Since the structure of the matrices was not used, the same argument holds for riemannian or Minkowski geometry.

#### REFERENCES

1. E. Cartan, *La méthode du repère mobile, la théorie des groupes continus et les espaces généralisés*, ASI 194, Paris, Hermann 1935.
2. T. H. Gronwall, *Note on the derivatives with respect to a parameter of the solution of a system of differential equation*, Ann. of Math., **20** (1918), 292-296.
3. H. Guggenheimer, *Differential Geometry*, New York, McGraw-Hill, 1963.
4. J. M. Sloss, *The bending of space curves into piecewise helical curves*, Pacific J. Math., **32** (1970), 231-239.

Received October 23, 1970. Research partially supported by NSF Grant GP-19133.

POLYTECHNIC INSTITUTE OF BROOKLYN



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

C. R. HOBBY  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

|   |     |
|---|-----|
| Louis I. Alpert and L. V. Toralballa, <i>An elementary definition of surface area in <math>E^{n+1}</math> for smooth surfaces</i> ..... | 261 |
| Eamon Boyd Barrett, <i>A three point condition for surfaces of constant mean curvature</i> .....  | 269 |
| Jan-Erik Björk, <i>On the spectral radius formula in Banach algebras</i> .....  | 279 |
| Peter Botta, <i>Matrix inequalities and kernels of linear transformations</i> .....   | 285 |
| Bennett Eisenberg, <i>Baxter's theorem and Varberg's conjecture</i> .....   | 291 |
| Heinrich W. Guggenheimer, <i>Approximation of curves</i> .....  | 301 |
| A. Hedayat, <i>An algebraic property of the totally symmetric loops associated with Kirkman-Steiner triple systems</i> .....            | 305 |
| Richard Howard Herman and Michael Charles Reed, <i>Covariant representations of infinite tensor product algebras</i> .....              | 311 |
| Domingo Antonio Herrero, <i>Analytic continuation of inner function-operators</i> .....   | 327 |
| Franklin Lowenthal, <i>Uniform finite generation of the affine group</i> .....  | 341 |
| Stephen H. McCleary, <i>0-primitive ordered permutation groups</i> .....  | 349 |
| Malcolm Jay Sherman, <i>Disjoint maximal invariant subspaces</i> .....  | 373 |
| Mitsuru Nakai, <i>Radon-Nikodým densities and Jacobians</i> .....   | 375 |
| Mitsuru Nakai, <i>Royden algebras and quasi-isometries of Riemannian manifolds</i> .....  | 397 |
| Russell Daniel Rupp, Jr., <i>A new type of variational theory sufficiency theorem</i> .....   | 415 |
| Helga Schirmer, <i>Fixed point and coincidence sets of biconnected multifunctions on trees</i> .....                                    | 445 |
| Murray Silver, <i>On extremal figures admissible relative to rectangular lattices</i> .....   | 451 |
| James DeWitt Stein, <i>The open mapping theorem for spaces with unique segments</i> .....   | 459 |
| Arne Stray, <i>Approximation and interpolation</i> .....  | 463 |
| Donald Curtis Taylor, <i>A general Phillips theorem for <math>C^*</math>-algebras and some applications</i> .....                       | 477 |
| Florian Vasilescu, <i>On the operator <math>M(Y) = TYS^{-1}</math> in locally convex algebras</i> .....                                 | 489 |
| Philip William Walker, <i>Asymptotics for a class of weighted eigenvalue problems</i> .....   | 501 |
| Kenneth S. Williams, <i>Exponential sums over <math>GF(2^n)</math></i> .....  | 511 |