

Pacific Journal of Mathematics

THE OPEN MAPPING THEOREM FOR SPACES WITH UNIQUE SEGMENTS

JAMES DEWITT STEIN

THE OPEN MAPPING THEOREM FOR SPACES WITH UNIQUE SEGMENTS

JAMES D. STEIN, JR.

If X and Y are spaces with unique segments, an affine map from X into Y is a map which takes segments into segments. The purpose of this paper is to give conditions on spaces X and Y such that we can prove the following versions of the Open Mapping and Closed Graph Theorems: (1) a continuous affine map of X onto Y is open, and (2) an affine map of X onto Y with closed graph is continuous.

Let X and Y be spaces with unique segments, with $\Phi(x, y, t)$ ($0 \leq t \leq 1$) denoting the intrinsically parametrized segment from x to y . A map T between two such spaces is said to be affine if $T(\Phi(x, y, t)) = \Phi(Tx, Ty, t)$, and a subset A is convex if $x, y \in A, 0 \leq t \leq 1 \Rightarrow \Phi(x, y, t) \in A$. The purpose of this paper is to prove versions of the Open Mapping and Closed Graph theorems for classes of spaces with unique segments.

Throughout this paper, all metric spaces will be spaces with unique segments (unique curves of minimal, realizing the distance, length between any two points). The open sphere with center x and radius ε will be denoted by $S(x, \varepsilon)$.

DEFINITION 1. (X, d) is said to be regular if it is complete, the closure of convex sets is convex, open spheres are convex, and $x_n \rightarrow x_0 \Rightarrow \Phi(z, x_n, \alpha) \rightarrow \Phi(z, x_0, \alpha)$ for $z \in X, \alpha \in [0, 1]$.

DEFINITION 2. A sphere $S(x_0, \varepsilon)$ is said to be thick if for any $y \in S(x_0, \varepsilon)$ and $x \in X, x \neq y$, there is a $z \in S(x_0, \varepsilon)$ and $\alpha \in (0, 1]$ such that $y = \Phi(z, x, \alpha)$.

It is always possible to extend geodesics into thick spheres.

DEFINITION 3. A sphere $S(x_0, \varepsilon)$ is said to be round if, given $x \in S(x_0, \varepsilon), y \in S(x, \varepsilon - d(x_0, x))$, and λ such that $d(x, y) < \lambda < \varepsilon - d(x_0, x) \Rightarrow$ there is a $z \in S(x, \varepsilon - d(x_0, x))$ and $\alpha \in (0, 1)$ such that $d(x, z) = \lambda, y = \Phi(x, z, \alpha)$.

Given any sphere $S(x, \delta)$ contained in a round sphere $S(x_0, \varepsilon)$ and any $y \in S(x, \delta), y$ lies on a geodesic connecting x and a point in $S(x, \delta)$ whose distance from x is arbitrarily close to δ . It should be noted that both thickness and roundness are hereditary properties; that is, if $S(x_0, \varepsilon)$ is thick (round) and $S(x, \delta) \subseteq S(x_0, \varepsilon)$ then $S(x, \delta)$ is thick (round).

We now prove an Open Mapping theorem.

THEOREM 1. *Let (X, d) and (X, d') be regular, and assume that each point in X is the center of a thick sphere, and each point in Y is the center of a round sphere. Assume further that, for each open $U \subseteq Y$, $y \in Y$ and $\alpha \in (0, 1]$, the set $\{\Phi(y, u, \alpha) \mid u \in U\}$ is open. Let T be continuous affine map of X onto Y . Then T is open.*

Proof. Let U be an open subset of X , let $x_0 \in U$, and let $\varepsilon > 0$ such that $S(x_0, \varepsilon)$ is thick and a subset of U . Let $U_k = S(x_0, k)$, $k = 1, 2, \dots$; since T is into, $Y = \bigcup_{k=1}^{\infty} \overline{T(U_k)}$. By the Baire Category Theorem, some $\overline{T(U_k)}$ contains an $S(Tx', \varepsilon')$. Since U_k is convex and T is affine, $T(U_k)$ is convex, and so $\overline{T(U_k)}$ is convex. Since thickness is hereditary, we can assume $\varepsilon < k$. Choose $x \in S(x_0, \varepsilon)$ and $\alpha \in (0, 1]$ with $x_0 = \Phi(x, x', \alpha)$; letting $z = Tx$ we have $Tx_0 = \Phi(z, Tx', \alpha)$, and $z \in \overline{T(U_k)}$. Now $Tx_0 \in \{\Phi(z, u, \alpha) \mid u \in S(Tx', \varepsilon')\}$, which is open by hypothesis and also a subset of $\overline{T(U_k)}$. We can therefore find a round sphere $S(Tx_0, \varepsilon'') \subset \overline{T(U_k)}$.

If $d'(y, Tx_0) < \varepsilon''$, let $\gamma = 1/2(\varepsilon'' + d'(y, Tx_0))$; since $S(Tx_0, \varepsilon'')$ is round and $\gamma < \varepsilon''$, there is a $z \in S(Tx_0, \varepsilon'')$ with $d'(z, Tx_0) = \gamma$ and $y = \Phi(Tx_0, z, \alpha)$, clearly $\alpha = \gamma^{-1}d'(y, Tx_0)$. Now $z \in \overline{T(U_k)}$, so choose $\{x_n \mid n = 1, 2, \dots\}$ with $d(x_0, x_n) < k$ and $Tx_n \rightarrow z$; let $z_n = \Phi(x_0, x_n, \alpha)$. Now $Tz_n = T(\Phi(x_0, x_n, \alpha)) = \Phi(Tx_0, Tx_n, \alpha) \rightarrow \Phi(Tx_0, z, \alpha) = y$, and $d(x_0, z_n) = \alpha d(x_0, x_n) = \gamma^{-1}d'(y, Tx_0)d(x_0, x_n) = 2d'(y, Tx_0)d(x_0, x_n)/(\varepsilon'' + d'(y, Tx_0)) \leq (2k/\varepsilon'')d'(y, Tx_0)$.

We now pause for a brief recapitulation. Given a point x_0 , if there is an integer n and a $\lambda > 0$ such that $S(Tx_0, \lambda)$ is round and $S(Tx_0, \lambda) \subset \overline{T(S(x_0, n))}$, then given $z \in S(Tx_0, \lambda)$ and $\eta > 0$ there is an $x \in X$ with $d(x, x_0) \leq (2n/\lambda)d'(z, Tx_0)$ and $d'(Tx, z) < \eta$. We now show that a similar approximation can be performed uniformly in a neighbourhood of x_0 .

Since T is continuous at x_0 , there is a $\delta > 0$ such that $d(x_0, y) < \delta \Rightarrow d'(Tx_0, Ty) < \lambda/2$. Let $y \in S(x_0, \min(\delta, n))$. If $z \in S(Ty, \lambda/2)$, then $d'(z, Tx_0) \leq d'(z, Ty) + d'(Ty, Tx_0) < \lambda$, so

$$S(Ty, \lambda/2) \subset S(Tx_0, \lambda) \subset \overline{T(S(x_0, n))}.$$

Now $z \in S(x_0, n) \Rightarrow d(z, y) \leq d(z, x_0) + d(x_0, y) < 2n$, so $S(x_0, n) \subset S(y, 2n)$, and so $S(Ty, \lambda/2) \subset \overline{T(S(y, 2n))}$. Since $S(Ty, \lambda/2) \subset S(Tx_0, \lambda)$, $S(Ty, \lambda/2)$ is round, and by repeating the previous computation, given $z \in S(Ty, \lambda/2)$ and $\eta > 0$, there is an $x \in X$ with

$$d(x, y) \leq [2(2n)/(\lambda/2)]d'(z, Ty) = \left(\frac{8n}{\lambda}\right)d'(z, Ty)$$

and $d'(Tx, z) < \eta$.

Therefore, for any $x_0 \in X$, there are constants δ , ε and M such that $d(x_0, y) < \delta$, $d'(z, Ty) < \varepsilon$ and $\eta > 0 \Rightarrow$ there is an $x \in X$ such that $d(x, y) \leq Md'(z, Ty)$ and $d'(Tx, z) < \eta$.

Let δ , ε , and M be defined as above. We now show T is open by showing $T(S(x_0, \rho))$ contains the sphere $S(Tx_0, \alpha)$, where $\alpha = 1/2 \min(\varepsilon, \delta/2M, \rho/2M)$. For $n = 1, 2, \dots$ let

$$\eta_n = \frac{1}{2} \min(\varepsilon, \delta/2^{n+1}M, \rho/2^{n+1}M) .$$

Let $z \in S(Tx_0, \alpha)$.

Since $d'(z, Tx_0) < \varepsilon$, choose $x_1 \in X$ with

$$d(x_0, x_1) \leq Md'(Tx_0, z), d'(Tx_1, z) < \eta_1 .$$

Now $d(x_0, x_1) \leq Md'(Tx_0, z) < \alpha M < \delta$ and $d'(Tx_1, z) < \eta_1 < \varepsilon$, so we can choose $x_2 \in X$ with $d(x_1, x_2) \leq Md'(z, Tx_1)$ and $d'(Tx_2, z) < \eta_2$. Inductively, assume we have chosen x_2, \dots, x_n such that, for $2 \leq j \leq n$, $d(x_{j-1}, x_j) \leq Md'(z, Tx_{j-1})$ and $d'(Tx_j, z) < \eta_j$. Now $d(x_0, x_n) \leq d(x_0, x_1) + d(x_1, x_2) + \dots + d(x_{n-1}, x_n) \leq Md'(Tx_0, z) + M \sum_{j=1}^{n-1} d'(z, Tx_j) < M\alpha + M \sum_{j=1}^{n-1} \eta_j < M(\delta/2M) + M \sum_{j=1}^{n-1} (\delta/2^{j+1}M) = \sum_{j=0}^{n-1} \delta/2^{j+1} < \delta$, and also $d'(Tx_n, z) < \eta_n < \varepsilon$, so we can choose $x_{n+1} \in X$ with $d(x_n, x_{n+1}) \leq Md'(z, Tx_n)$ and $d'(Tx_{n+1}, z) < \eta_{n+1}$.

If $1 \leq n < m$, then

$$\begin{aligned} d(x_n, x_m) &\leq d(x_n, x_{n+1}) + \dots + d(x_{m-1}, x_m) \\ &\leq M \sum_{j=n}^{m-1} d'(z, Tx_j) < M \sum_{j=n}^{m-1} \eta_j < \delta/2^n ; \end{aligned}$$

by completeness there is an $x \in X$ such that $x_n \rightarrow x$. Since $\eta_n \rightarrow 0$, $Tx_n \rightarrow z$, by continuity $Tx_n \rightarrow Tx$, so $Tx = z$. We are done if we can show that $d(x_0, x) < \rho$. For $n \geq 1$, $d(x_0, x) \leq d(x_0, x_1) + \dots + d(x_{n+1}, x_n) + d(x_n, x) \leq Md'(Tx_0, z) + M \sum_{j=1}^{n-1} d'(z, Tx_j) + d(x_n, x) < M\alpha + M \sum_{j=1}^{n-1} \eta_j + d(x_n, x) < M(\rho/3M) + M \sum_{j=1}^{n-1} (\rho/3 \cdot 2_jM) + d(x_n, x)$, so, since $x_n \rightarrow x$, we have $d(x_0, x) \leq \rho/3 + \sum_{j=1}^{\infty} (\rho/3 \cdot 2^j) = 2\rho/3 < \rho$, completing the proof.

If the hypotheses seem somewhat cumbersome, it should be remembered that the Uniform Boundedness and Open Mapping Theorems in linear topological spaces also place heavy restrictions on the topologies involved.

We conclude by making some remarks relevant to the Closed Graph Theorem, one of the most useful consequences of the Open Mapping Theorem. Call a space (X, d) which satisfies all the hypothesis listed in Theorem 1 a normal space. Given two normal spaces (X, d) and (Y, d') , if we can show that the product $(X \times Y, d'')$ (where $d''((x_1, y_1), (x_2, y_2)) = d((x_1, x_2) + d'(y_1, y_2))$ is normal, standard techniques (for example, [1] p. 100 and p. 116) will prove that, if $T: X \rightarrow Y$ is affine with closed graph, then T is continuous.

The verification that the product $(X \times Y, d'')$ of normal spaces (X, d) and (Y, d') follows quickly from the following lemma.

LEMMA 2. $(X \times Y, d'')$ is a space with unique segments.

Proof. Given a path λ in X and a path γ in Y , define $\lambda \times \gamma(t) = (\lambda(t), \gamma(t))$, and if μ is a path in $X \times Y$, define $\pi_X(\mu)$ and $\pi_Y(\mu)$ by $\mu(t) = (\pi_X(\mu)(t), \pi_Y(\mu)(t))$. Given a path ρ of finite length, its length will be denoted by $l(\rho)$, and the variation of ρ over a partition P of $[0, 1]$ will be denoted by $l(\rho; P)$. We assert that, if $\pi_X(\mu)$ and $\pi_Y(\mu)$ have finite length, then $l(\mu) = l(\pi_X(\mu)) + l(\pi_Y(\mu))$. Since $l(\mu; P) = l(\pi_X(\mu); P) + l(\pi_Y(\mu); P)$, clearly $l(\mu) \leq l(\pi_X(\mu)) + l(\pi_Y(\mu))$. The reverse inequality is proved by choosing $\varepsilon > 0$, finding partitions P_X and P_Y which approximate $l(\pi_X(\mu))$ and $l(\pi_Y(\mu))$ to within $\varepsilon/2$, letting P denote the common refinement of P_X and P_Y will yield $l(\mu) + \varepsilon > l(\pi_X(\mu)) + l(\pi_Y(\mu))$. If λ and γ are geodesics in X and Y respectively, then $l(\lambda \times \gamma) = l(\lambda) + l(\gamma)$. If μ is a path in $X \times Y$ with $l(\mu) < l(\lambda \times \gamma)$, then $l(\pi_X(\mu)) + l(\pi_Y(\mu)) < l(\lambda) + l(\gamma) \Rightarrow$ either $l(\pi_X(\mu)) < l(\lambda)$ or $l(\pi_Y(\mu)) > l(\gamma)$, a contradiction. Similarly, if $l(\mu) = l(\lambda \times \gamma)$, then $l(\pi_X(\mu)) = l(\lambda)$ and $l(\pi_Y(\mu)) = l(\gamma)$, which shows that $\pi_X(\mu) = \lambda$ and $\pi_Y(\mu) = \gamma$, so $\mu = \lambda \times \gamma$ and geodesics in $(X \times Y, d'')$ are unique.

In conclusion, it is interesting to note that the hypothesis that $S(x_0, \varepsilon)$ is convex is clearly analogous to the assumption that a linear topological space is locally convex. It is clear that many standard concepts in linear topological space theory are geodesically defined (such as an absorbing neighbourhood), and consequently are easily translated into the theory of spaces with unique segments. In addition to the question of which theorems from normed linear spaces hold in suitably restricted spaces with unique segments, the following problem arises quite naturally: is there a category whose objects include both linear topological spaces and spaces with unique segments, and whose morphisms are the "affine" maps? A reasonable candidate would be topological spaces X with a geodesic-like structure such as a map $\Phi: X \times X \times I \rightarrow X$ (where $I = [0, 1]$) satisfying such conditions as (1) for fixed $x, y \in X$, $\Phi(x, y, t)$ is a continuous function of t , (2) $\Phi(x, x, t) = x$, and (3) $\Phi(\Phi(x, y, s), \Phi(x, y, t), u) = \Phi(x, y, (1 - u)s + ut)$. Some work has been done on spaces with convexity structures, but from a purely algebraic standpoint. It would be extremely interesting to find out to what extent such theorems as the Open Mapping Theorem and the Uniform Boundedness Theorem can be generalized.

REFERENCES

1. W. Rudin, *Real and Complex Analysis*, New York, McGraw-Hill, 1966.
2. J. Stein, *Two uniform boundedness theorems*, Pacific J. Math., **38** (1971), 251-260.

Received October 28, 1970.

UNIVERSITY OF CALIFORNIA, LOS ANGELES

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Louis I. Alpert and L. V. Toralballa, <i>An elementary definition of surface area in E^{n+1} for smooth surfaces</i>	261
Eamon Boyd Barrett, <i>A three point condition for surfaces of constant mean curvature</i>	269
Jan-Erik Björk, <i>On the spectral radius formula in Banach algebras</i>	279
Peter Botta, <i>Matrix inequalities and kernels of linear transformations</i>	285
Bennett Eisenberg, <i>Baxter's theorem and Varberg's conjecture</i>	291
Heinrich W. Guggenheimer, <i>Approximation of curves</i>	301
A. Hedayat, <i>An algebraic property of the totally symmetric loops associated with Kirkman-Steiner triple systems</i>	305
Richard Howard Herman and Michael Charles Reed, <i>Covariant representations of infinite tensor product algebras</i>	311
Domingo Antonio Herrero, <i>Analytic continuation of inner function-operators</i>	327
Franklin Lowenthal, <i>Uniform finite generation of the affine group</i>	341
Stephen H. McCleary, <i>0-primitive ordered permutation groups</i>	349
Malcolm Jay Sherman, <i>Disjoint maximal invariant subspaces</i>	373
Mitsuru Nakai, <i>Radon-Nikodým densities and Jacobians</i>	375
Mitsuru Nakai, <i>Royden algebras and quasi-isometries of Riemannian manifolds</i>	397
Russell Daniel Rupp, Jr., <i>A new type of variational theory sufficiency theorem</i>	415
Helga Schirmer, <i>Fixed point and coincidence sets of biconnected multifunctions on trees</i>	445
Murray Silver, <i>On extremal figures admissible relative to rectangular lattices</i>	451
James DeWitt Stein, <i>The open mapping theorem for spaces with unique segments</i>	459
Arne Stray, <i>Approximation and interpolation</i>	463
Donald Curtis Taylor, <i>A general Phillips theorem for C^*-algebras and some applications</i>	477
Florian Vasilescu, <i>On the operator $M(Y) = TYS^{-1}$ in locally convex algebras</i>	489
Philip William Walker, <i>Asymptotics for a class of weighted eigenvalue problems</i>	501
Kenneth S. Williams, <i>Exponential sums over $GF(2^n)$</i>	511