

Pacific Journal of Mathematics

CARDINALITY OF k -COMPLETE BOOLEAN ALGEBRAS

W. WISTAR (WILLIAM) COMFORT AND ANTHONY WOOD HAGER

CARDINALITY OF \aleph -COMPLETE BOOLEAN ALGEBRAS

W. W. COMFORT AND ANTHONY W. HAGER

An infinite complete Boolean algebra satisfies $|B|^{\aleph_0} = |B|$ (where $| \cdot |$ denotes cardinality). This is a theorem of R. S. Pierce, derived in consequence of his general decomposition theorem [9]. It is here shown (directly) that $|B|^{\aleph_0} = |B|$ for B merely countably complete; this has the corollary (actually, equivalent) that if A is an algebra of measurable functions modulo null functions, and D is a subset of A which is dense in the uniform topology, then $|D| = |A|$. The relation $|B|^{\aleph} = |B|$ for \aleph -complete Boolean algebras B is considered; the main result is a structure theorem for the nontrivial counterexamples (which are shown to exist abundantly).

The contribution of the referee deserves special mention. In detail, he translated our original paper from topology into Boolean algebras, simplifying the results and their proofs, and he removed our use of the Generalized Continuum Hypothesis in the theorem on countably complete algebras. (We announced the latter theorem, with GCH, in [3]. Subsequently, Monk and Sparks announced the result, with no mention of GCH, in [8]; this was our first knowledge that the result was obtainable with GCH. We do not know how our methods and those of Monk and Sparks compare.)

Following the referee's advice, our setting is Boolean algebras, but we indicate the translation to topology. The means is Stone duality, of course, whereby the Boolean algebra B is isomorphic to the algebra of open-and-closed subsets of the Stone space $S(B)$ [7, 11]. It results that $|B| = wS(B)$ (where w is the weight, or least cardinal of an open basis), and that B is \aleph -complete if and only if $S(B)$ has the property that the closure of $\bigcup \mathcal{U}$ is open whenever \mathcal{U} is a family of clopen sets with $|\mathcal{U}| \leq \aleph$. Thus, the dual form of the theorem on countably complete algebras is that $(wX)^{\aleph_0} = wX$ whenever X is infinite, compact, and has the above property, commonly called basic disconnectivity. (The Stone spaces of *complete* algebras are said to be extremally disconnected.)

For B a Boolean algebra and $b \in B$, we write $(b) = \{a \in B : a \leq b\}$ and we set $|b| = |(b)|$. A subset D of B is *disjointed* if $d_1 \neq d_2$ in D implies $d_1 \wedge d_2 = 0$.

1. LEMMA. *Let \aleph be infinite, B a \aleph -complete Boolean algebra and D a disjointed subset of B with $|D| \geq \aleph$. Let $b \equiv \text{lub } D$. Then*

$$(b) \cong \prod_{a \in D} (a) \quad \text{and} \quad |b| = \prod_{a \in D} |a|.$$

Proof. Define the Boolean homomorphism $\varphi: (b) \rightarrow \prod_{a \in D} (a)$ by the rule $(\varphi(c))_a = a \wedge c$. Clearly, φ is one-to-one. If $p \in \prod_{a \in D} (a)$, then $p_a \leq a$ for each $a \in D$, and $p = \varphi(\text{lub}\{p_a: a \in D\})$ results from \mathfrak{k} -completeness.

2. THEOREM. *If B is an infinite, countably complete Boolean algebra, then $|B|^{\aleph_0} = |B|$.*

Proof. Suppose there is a counterexample, and choose one, B , of minimal cardinal m . Let $J \equiv \{b \in B: |b| < m\}$; if $a \in J$, and $|a|$ is infinite, then $|a|^{\aleph_0} = |a|$ by minimality of m . We assert that

- (1) J is a σ -ideal of B ; and
- (2) B/J is finite, so that $|J| = m$.

Surely J is an ideal. Given a countably infinite subset D of J let $a = \text{lub } D$ and let D' be a countably infinite, disjointed subset of J with $b = \text{lub } D'$. Then

$$|b| = \prod_{a \in D'} |a| \geq 2^{\aleph_0}$$

by the lemma, so

$$|b|^{\aleph_0} = \prod_{a \in D'} |a|^{\aleph_0} = \prod_{a \in D'} |a| = |b|;$$

thus $|b| \neq m$, so $|b| < m$ and $b \in J$ and (1) is proved. If (2) fails there is a disjointed sequence $\{b_n\}_{n < \omega}$ of elements of $B \setminus J$, so that (again from the lemma) one has

$$m = |B| \geq |\text{lub}\{b_n: n < \omega\}| = \prod_{n < \omega} |b_n| = m^{\aleph_0}.$$

To complete the proof let

$$Q = \{S \subset J: |S| \leq \aleph_0\}$$

and for $S \in Q$ define $\varphi(S) = \text{lub } S$. Then $\varphi(S) \in J$ whenever $S \in Q$ (by (1)), and for $b \in J$ one has

$$\varphi^{-1}(b) \subset \{S \in Q: S \subset (b)\}.$$

Thus for each b in J either (b) is finite or

$$|\varphi^{-1}(b)| \leq |b|^{\aleph_0} = |b|,$$

so from (2) we have

$$m < m^{\aleph_0} = |J|^{\aleph_0} = |Q| \leq \sum_{b \in J} \aleph_0 \cdot |b| < m \cdot \aleph_0 \cdot m = m.$$

This contradiction completes the proof.

We derive as a corollary the result mentioned earlier on algebras of measurable functions perhaps modulo an ideal of null functions. Let T be a set and $\mathcal{A} \subset 2^T$ a σ -field; let M be the real functions f with $f^{-1}(\theta) \in \mathcal{A}$ if θ is open. Let \mathcal{N} be a σ -ideal of \mathcal{A} and N those $f \in M$ with support in \mathcal{N} . Endow M/N with the metric of uniform convergence except on a member of \mathcal{N} . (In this metric, M/N is complete.)

3. COROLLARY. *Any dense subset of M/N has cardinality $|M/N|$.*

Proof. If X is a topological space, let $D(X)$ be the almost-finite extended real-valued functions on X . Taking X the space of maximal ideals in M/N , X is basically disconnected and M/N is isomorphic and isometric to $D(X)$ [6] (where $D(X)$ has the metric of uniform convergence on all of X). With δ denoting minimum cardinal of a dense subset, $\delta D(X) = \delta C(X)$ [4]. Since X is compact, $wX = \delta C(X)$ [11]. By the dual form of the preceding theorem, $(wX)^{\aleph_0} = wX$. But of course, $\delta C(X) \leq |C(X)| \leq (\delta C(X))^{\aleph_0}$, since sequences from any dense subset of $C(X)$ determine $C(X)$ (as with any metric space). The proof is complete.

The corollary applies, of course, to Lebesgue measurable functions on the reals, modulo or not the usual null functions, and to Baire and Borel functions on any space modulo or not various ideals of null functions, e.g., the ones vanishing except on meager sets. See [7, 11].

We next consider the possibility of generalizing the result on countably complete Boolean algebras. As a point of reference, consider the statement: if B is infinite and \mathfrak{k} -complete, and contains a disjointed family of cardinal \mathfrak{k} , then $|B|^{\mathfrak{k}} = |B|$. (The last hypothesis prevents the choice of complete B and relatively huge \mathfrak{k} .) We present a positive result under additional hypotheses, a class of counterexamples, and a theorem describing all counterexamples. For these, recall that the cofinality of the cardinal m , $cf(m)$, is the least \mathfrak{k} for which a set of m points is the union of \mathfrak{k} sets each of $< m$ points. m is called regular if $cf(m) = m$, and otherwise singular. We encounter $cf(m)$ in these considerations because $m^{cf(m)} > m$ always, and with *GCH*, $cf(m)$ is the least such exponent [1]. Thus, we are forced, essentially, to consider algebras B which are $cf(|B|)$ -complete, and the structure theorem below (6.) is for these.

4. THEOREM. *Suppose that B is infinite, \mathfrak{k} -complete, has a disjointed family of cardinal \mathfrak{k} , and that $\mathfrak{p} < |B|$ implies $2^{\mathfrak{p}} \leq |B|$. If either $|B|$ is regular, or for each $b \in B$ there is $a \leq b$ with $|a| < |B|$, then $|B|^{\mathfrak{k}} = |B|$.*

This is from the original version of the present paper titled the relation $m^t = m$ in Stonian spaces. The proof can be found in the first author's survey [2] (in topological dual).

5. **EXAMPLES.** Let m be singular with $m^{\aleph_0} = m$. Let \mathfrak{f} and \mathfrak{l} satisfy $cf(m) \leq \mathfrak{f} \leq \mathfrak{l} \leq \mathfrak{l}' < m$. Let A_1 be the completion by cuts of the free Boolean algebra on m generators (or equivalently, the algebra of open-and-closed subsets of the projective cover [5] of the topological space 2^m); so $|A_1| = m$. Let A_2 be the algebra of \mathfrak{f} -sets (i.e., sets of cardinal at most \mathfrak{f}) and co- \mathfrak{f} -sets in a set of cardinal \mathfrak{l} ; evidently, A_2 is \mathfrak{f} -complete, but not \mathfrak{f}^+ -complete unless $\mathfrak{f} = \mathfrak{l}$ (in which case A_2 is complete).

The algebra $B \equiv A_1 \times A_2$ is \mathfrak{f} -complete and has a disjointed family of power \mathfrak{f} (indeed, \mathfrak{l}), but

$$|B| = m < m^{cf(m)} \leq m^{\mathfrak{f}}.$$

And all examples are like this.

6. **THEOREM.** *Let m be infinite and singular, with the property that for each $\mathfrak{p} < m$ either $2^{\mathfrak{p}} < m$ or $2^{\mathfrak{p}} = \mathfrak{p}^+$. If B is a $cf(m)$ -complete Boolean algebra with $|B| = m$, then $B = A_1 \times A_2$, where*

- (a) $|A_1| = m$ and $|A_2| < m$;
- (b) if $0 \neq b \in A_1$, then $|b| = m$;
- (c) if $D \subset A_1$ and D is disjointed, then $|D| < cf(m)$;
- (d) A_1 is complete.

Proof. Define

$$J_1 = \{a \in B : |c| = m \text{ whenever } 0 \neq c \in (a)\}$$

and $J_2 = \{b \in B : |b| < m\}$, so that J_1 and J_2 are disjoint ideals in B .

Given a disjointed subset D of J_1 with $|D| \leq cf(m)$ we have from the lemma

$$m = |B| = |\text{lub } D| = \prod_{a \in D} |a| = m^{|D|},$$

so that $|D| < cf(m)$. It follows that $J_1 = (a_1)$ for some element a_1 of J_1 : given a maximal disjointed subset D of J_1 we have $|D| < cf(m)$ from the computation above, so that $\text{lub } D = a_1$ exists in B ; evidently $a_1 \in J_1$, and $J_1 = (a_1)$ by maximality. Another consequence is that (a_1) is a complete Boolean algebra: given a subset S of (a_1) some disjointed subset D of (a_1) is maximal with respect to the property that it refines S (i.e., for each d in D , $d \in (s)$ for some s in S), so that $\text{lub } D$ exists in (a_1) .

Let a_2 be the complement in B of (a_1) , and set $A_1 = (a_1)$ and $A_2 = (a_2)$. The other assertions being obvious, it remains only to show that $|A_2| < m$.

We claim first:

(*) if E is a disjointed subset of J_2 and $p = \sum_{b \in E} |b|$, then $p < m$. If $p \geq m$ then either $|E| = m$ or $\sup\{|b| : b \in E\} = m$. In the former case we would have

$$m = |B| = |\{S \subset E : |S| = cf(m)\}| = m^{cf(m)} > m,$$

and in the latter, replacing E if necessary by a subset E' for which $|E'| = cf(m)$ and $\sup\{|b| : b \in E'\} = m$,

$$m \geq |B| \geq |\text{lub } E'| = \prod_{b \in E'} |b| > m.$$

In either case a contradiction is achieved and (*) is proved.

Now let E be a maximal, disjointed subset of (a_2) , let $|E| = p < m$, and set $K = \bigcup_{b \in E} (b)$. Because E is maximal the map $c \rightarrow K \cap (c)$ is one-to-one from (a_2) to the power set of K . Thus $|a_2| \leq 2^p$ (and $p < m$ by (*)). The assertion $|a_2| < m$ now follows from the cardinality hypothesis of this theorem together with the fact that p^+ is a regular cardinal.

REFERENCES

1. Heinz Bachmann, *Transfinite Zahlen*, Springer-Verlag, Berlin, 1955.
2. W. W. Comfort, *A survey of cardinal invariants*, General topology and its applications **1** (1971), 163-199.
3. W. W. Comfort and Anthony W. Hager, *The relation $m^k = m$ in Stonian spaces*, Notices Amer. Math. Soc., **17** (1970), 464 (abstract).
4. W. W. Comfort and Anthony W. Hager, *Dense subspaces of some spaces of continuous functions*, Math. Zeitschrift **114** (1970), 373-389.
5. Andrew M. Gleason, *Projective topological spaces*, Illinois J. Math., **2** (1958), 482-489.
6. Anthony W. Hager, *Algebras of measurable functions*, Duke Math. J., **38** (1971), 21-27.
7. Paul R. Halmos, *Lectures on Boolean Algebras*, D. Van Nostrand Co., Inc., Princeton, 1963.
8. J. Donald Monk and Paul R. Sparks, *Counting Boolean algebras*, Notices Amer. Math. Soc., **18** (1971), 551 (abstract).
9. R. S. Pierce, *A note on complete Boolean algebras*, Proc. Amer. Math. Soc., **9** (1958), 892-896.
10. Roman Sikorski, *Boolean Algebras*, Springer-Verlag, Berlin 1964.
11. Yu. M. Smirnov, *On the weight of the ring of bounded continuous functions over a normal space*, Mat. Sbornik N. S. **30** (72) (1952), 213-218 (Russian).

Received November 2, 1970. The authors gratefully acknowledge partial support received from the National Science Foundation (U.S.A.) under grants NSF-GP-8357 and NSF-GP-18825.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index. to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics

Vol. 40, No. 3

November, 1972

Wazir Husan Abdi, <i>A quasi-Kummer function</i>	521
Vasily Cateforis, <i>Minimal injective cogenerators for the class of modules of zero singular submodule</i>	527
W. Wistar (William) Comfort and Anthony Wood Hager, <i>Cardinality of k-complete Boolean algebras</i>	541
Richard Brian Darst and Gene Allen DeBoth, <i>Norm convergence of martingales of Radon-Nikodym derivatives given a σ-lattice</i>	547
M. Edelstein and Anthony Charles Thompson, <i>Some results on nearest points and support properties of convex sets in c_0</i>	553
Richard Goodrick, <i>Two bridge knots are alternating knots</i>	561
Jean-Pierre Gossez and Enrique José Lami Dozo, <i>Some geometric properties related to the fixed point theory for nonexpansive mappings</i>	565
Dang Xuan Hong, <i>Covering relations among lattice varieties</i>	575
Carl Groos Jockusch, Jr. and Robert Irving Soare, <i>Degrees of members of Π_1^0 classes</i>	605
Leroy Milton Kelly and R. Rottenberg, <i>Simple points in pseudoline arrangements</i>	617
Joe Eckley Kirk, Jr., <i>The uniformizing function for a class of Riemann surfaces</i>	623
Glenn Richard Luecke, <i>Operators satisfying condition (G_1) locally</i>	629
T. S. Motzkin, <i>On $L(S)$-tuples and l-pairs of matrices</i>	639
Charles Estep Murley, <i>The classification of certain classes of torsion free Abelian groups</i>	647
Louis D. Nel, <i>Lattices of lower semi-continuous functions and associated topological spaces</i>	667
David Emroy Penney, II, <i>Establishing isomorphism between tame prime knots in E^3</i>	675
Daniel Rider, <i>Functions which operate on $\mathbb{F}L_p(T)$, $1 < p < 2$</i>	681
Thomas Stephen Shores, <i>Injective modules over duo rings</i>	695
Stephen Simons, <i>A convergence theorem with boundary</i>	703
Stephen Simons, <i>Maximinimax, minimax, and antiminimax theorems and a result of R. C. James</i>	709
Stephen Simons, <i>On Ptak's combinatorial lemma</i>	719
Stuart A. Steinberg, <i>Finitely-valued f-modules</i>	723
Pui-kei Wong, <i>Integral inequalities of Wirtinger-type and fourth-order elliptic differential inequalities</i>	739
Yen-Yi Wu, <i>Completions of Boolean algebras with partially additive operators</i>	753
Phillip Lee Zenor, <i>On spaces with regular G_δ-diagonals</i>	759