Pacific Journal of Mathematics

SOME GEOMETRIC PROPERTIES RELATED TO THE FIXED POINT THEORY FOR NONEXPANSIVE MAPPINGS

JEAN-PIERRE GOSSEZ AND ENRIQUE JOSÉ LAMI DOZO

SOME GEOMETRIC PROPERTIES RELATED TO THE FIXED POINT THEORY FOR NONEXPANSIVE MAPPINGS

J.-P. Gossez and E. Lami Dozo

The main result of this paper asserts that if a Banach space admits a sequentially weakly continuous duality function, then a condition introduced by Opial to characterize weak limits by means of the norm is satisfied and the space has normal structure in the sense of Brodskii-Milman. This result of geometric nature allows some unification in the fixed point theory for both single-valued and multi-valued non-expansive mappings.

Let K be a nonempty weakly compact convex subset of a real Banach space X and let T be a nonexpansive mapping of K into its nonempty compact subsets (i.e., $D(Tx, Ty) \leq ||x - y||$ for all $x, y \in K$, where D(,) denotes the Hausdorff metric). While the question of the existence of a fixed point for T remains open, several positive results were proved recently under various conditions of geometric type on the norm of X. We list here the conditions we have in mind:

(I) (Browder [5]) X admits a sequentially continuous duality function $F_{\phi}: X, \sigma(X, X^*) \to X^*, \sigma(X^*, X)$ (i.e., a function F_{ϕ} such that $\langle x, F_{\phi}(x) \rangle = ||x|| ||F_{\phi}(x)||$ and $||F_{\phi}(x)|| = \phi(||x||)$ for all $x \in X$, where $\phi: \mathbf{R}^+ \to \mathbf{R}^+$ is continuous strictly increasing with $\phi(0) = 0$ and $\phi(+\infty) = +\infty$).

(II) (Opial [17]) If a sequence $\{x_n\}$ converges weakly in X to x_0 , then $\liminf ||x_n - x|| > \liminf ||x_n - x_0||$ for all $x \neq x_0$.

(III) (Brodskii-Milman [4]) Every weakly compact convex subset H of X has normal structure (i.e., for each convex subset L of H which contains more than one point there exists $x \in L$ such that $\sup\{||x - y||; y \in L\} < \sup\{||u - v||; u, v \in L\}$).

When T is single-valued, the existence of a fixed point for T in K was proved by Browder [5] if X satisfies (I) and if T can be extended outside K in a nonexpansive way, and by Kirk [12] if X satisfies (III). A similar situation occurs in the multivalued case where one also encounters two different approaches: one by Browder [6] who proved a fixed point theorem under condition (I) and some additional assumptions, and another by the second author [14] who obtained the same conclusion under condition (II).

It is a consequence of our main theorem that in both cases the

second approach is more general than the first¹:

THEOREM 1. (I) implies (II) and (II) implies (III). No converse implication holds, not even when X and X^* are supposed to be uniformly convex.

Although (II) does not imply (I), there is some result in that direction, which supports the feeling that the gap between (II) and (III) is much deeper than that between (I) and (II). To state this result we define (I₀) as (I) except that F_{ϕ} is only required to be sequentially continuous at zero and (II₀) as (II) except that > is replaced by \geq .

THEOREM 2. (I_0) implies (II_0) . The converse implication holds when the norm of X is uniformly Gâteaux differentiable.

Theorems 1 and 2 are proved in §2 and §3 respectively. Some related results and several examples are presented in §4 which shed more light on the connections between these geometric properties.

In §5 we show that the space c_0 endowed with Day's norm, which is locally uniformly convex (cf. [18]), does not satisfy (III). This example should be connected with the well-known facts that all uniformly convex spaces, as well as the spaces showed by Day and Lovaglia to be locally uniformly convex but not isomorphic to any uniformly convex space, satisfy (III) (cf. [3, 6, 11]).

2. The main result. To prove Theorem 1 we need two lemmas about the duality map $J_{\phi}: X \to 2^{X^*}$ defined by

$$J_{\phi}(x) = \{x^* \in X^*; \langle x, x^* \rangle = ||x|| ||x^*|| \text{ and } ||x^*|| = \phi(||x||)\}$$

for all $x \in X$. In this context condition (I) asserts the existence for some gauge ϕ of a sequentially weakly continuous selection for J_{ϕ} . The first lemma follows from the monotonicity of J_{ϕ} ; it has been extended in [10] to general monotone operators.

LEMMA 1. If X satisfies (I) (for some gauge ϕ), then J_{ϕ} is single-valued (for any gauge ϕ).

Proof. The monotone operator F_{ϕ} is hemicontinuous by (I), thus maximal monotone by Minty's classical argument [15]. Since J_{ϕ} is

¹ It should be noticed that the existence theorems of [12] and [14] have been recently extended in [13] and [2] where it is only required that T maps the boundary of K into K.

also monotone we must have $J_{\phi} = F_{\phi}$. That J_{ψ} is single-valued for another gauge ψ follows from the equality $J_{\psi}(x) = \psi(||x||)/\phi(||x||)J_{\phi}(x)$.

The second lemma uses the observation of Asplund [1] that J_{ϕ} is the subdifferential of the convex function $\Phi(||x||)$ where

$$\Phi(t) = \int_0^t \phi(s) ds,$$

i.e.,

$$J_{\phi}(x) = \{x^* \in X^*; \, \varPhi(||y||) \ge \varPhi(||x||) + \langle y - x, \, x^*
angle \forall y \in X\}$$
 .

LEMMA 2. If J_{ϕ} is single-valued then

for all $x, y \in X$.

Proof. If J_{ϕ} is single-valued then J_{ϕ} is the Gâteaux gradient of $\varPhi(||x||)$; this follows from a general result in the theory of convex functions (see [16; p. 66]). But J_{ϕ} is easily verified to be hemicontinuous. Consequently the lemma just expresses the fact that a function of a real variable is the integral of its continuous derivative.

Proof of Theorem 1. Assume (I) and let $x_n \rightarrow x_0$ (\rightarrow will denote weak convergence, \rightarrow norm convergence). By Lemmas 1 and 2,

for all $x \in X$, so

$$\liminf arPhi(||x_n-x||) \ge \liminf arPhi(||x_n-x_0||) + \liminf \int_0^1 \cdots dt$$
.

The sequential weak continuity of J_{ϕ} and the dominated convergence theorem give

(1)
$$\lim \inf \mathscr{P}(||x_n - x||) \ge \lim \inf \mathscr{P}(||x_n - x_0||) + \int_0^1 ||x_0 - x|| \mathscr{P}(t||x_0 - x||) dt,$$

an inequality which clearly implies condition (II).

The proof that (II) implies (III) relies upon a characterization of normal structure given in [11]:X satisfies (III) if and only if X does not contain a diametral sequence $\{x_n\}$ weakly converging to zero (i.e., a nonconstant sequence with

(2)
$$d(x_n; \operatorname{co} \{x_1, \cdots, x_{n-1}\}) \longrightarrow \delta(\{x_n\}),$$

where $d(x_n; \text{ co } \{x_1, \dots, x_{n-1}\})$ denotes the distance of x_n to the convex hull of $\{x_1, \dots, x_{n-1}\}$ and $\delta(\{x_n\})$ the diameter of $\{x_n\}$.

Assume that (III) does not hold and take such a sequence. It follows from (2) that

$$\lim ||x_n - y|| = \delta(\{x_n\})$$

for all $y \in \operatorname{co} \{x_n\}$, hence for all $y \in \operatorname{co} \{x_n\}$. Taking y = 0, we get $\lim ||x_n|| = \delta(\{x_n\})$, but for each $y = x_{n_0}$ we obtain $\lim ||x_n - x_{n_0}|| = \delta(\{x_n\})$. This contradicts (II).

We now turn to the last part of Theorem 1. When 1 $<math>p \neq 2$, $L^{p}(0, 2\Pi)$ satisfies (III) since it is uniformly convex, but Opial [17] showed that even (II₀) does not hold. When 1 the $Hilbert product of <math>l^{p}$ and l^{q} satisfies (II) (cf. [14]; since it is easily verified that (I₀) holds, (II) also follows from Theorem 2 of §3 and Proposition 1 of §4), but Bruck [7] showed that (I) does not hold.

REMARKS 1. A finite dimensional space whose norm is not differentiable provides another example of a space satisfying (I_0) , (II) but not (I) by Lemma 1.

2. In the Hilbert space case when $\phi(t) = t$, estimation (1) reduces to an estimation obtained by Opial [17; p. 592].

3. A partial converse. The following simple lemma whose proof proceeds by taking subsequences will be needed.

LEMMA 3. Conditions (II) and (II₀) are respectively equivalent to the analogous conditions obtained by replacing lim inf by lim sup.

Proof of Theorem 2. Assume (I₀) and let $x_n \rightharpoonup x_0$. As J_{ϕ} is the subdifferential of $\Phi(||x||)$ we have

$$arPsi_{n}(||x_{n}-x||) \ge arPsi_{n}(||x_{n}-x_{0}||) + \langle x_{0}-x, F_{\phi}(x_{n}-x_{0}) \rangle$$

for all $x \in X$, so

$$\liminf \Phi(||x_n - x||) \ge \liminf \Phi(||x_n - x_0||) ,$$

an inequality which clearly implies condition (II_0) .

In the second part of Theorem 2, the assumption of uniform Gâteaux differentiability is equivalent to the condition that J_{ϕ} be singlevalued and continuous on X, $|| \quad ||$ into X^* , $\sigma(X^*, X)$, uniformly on bounded sets; this follows easily from a result of Cudia [8; p. 302]. Under this assumption we will show that if (II₀) holds then for any ϕ , J_{ϕ} is sequentially weakly continuous at zero.

Let $x_n \rightarrow 0$ and suppose that $J_{\phi}(x_n)$ does not converge to zero for

 $\sigma(X^*, X)$. Then there exist $z \in X$, a subsequence $\{x_m\}$ and $x^* \in X^*$ such that $\langle z, J_{\phi}(x_m) \rangle \rightarrow \langle z, x^* \rangle \neq 0$. Define

$$f(x) = \limsup \Phi(||x_m - x||);$$

f is a continuous convex function on X which assumes its minimum at x = 0 by condition (II₀) and Lemma 3. Hence

$$\frac{1}{\lambda}(f(-\lambda y) - f(0) \ge 0 \qquad \forall \ y \in x, \ \forall \ \lambda > 0 \ ,$$

and thus

$$\limsup rac{1}{\lambda} (arPhi(||x_m+\lambda y||) - arPhi(||x_m||)) \ge 0 ;$$

as J_{ϕ} is the subdifferential of $\mathcal{O}(||x||)$,

$$\limsup \langle y, J_{\phi}(x_m + \lambda y) \rangle \geq 0$$
.

Letting $\lambda \downarrow 0$, limits can be interchanged by the uniform continuity of J_{ϕ} , and consequently

$$\limsup < y, \, J_{\phi}(x_m) > \geqq 0 \qquad \forall y \in X$$
 .

In particular $\langle z, x^* \rangle = 0$, a contradiction.

REMARK. We do not know whether the differentiability hypothesis in Theorem 2 is essential.

4. Further connections. We begin this section by showing that the following situation holds in general:



where $(A) \rightarrow (B)[(A) \rightarrow (B)]$ means that (A) implies [does not imply] (B). Taking into account §2 and 3, it suffices to exhibit a space satisfying (I₀) but not (III). Consider the space l^2 endowed with the norm

$$||x|| = \max \left\{ \frac{1}{2} ||x||_{l^2}; ||x||_{l^{\infty}} \right\}.$$

James [3] proved that (III) does not hold; however the application

$$x = (x(1), x(2), \dots) \longrightarrow egin{cases} iggl(rac{x}{4} ext{ if } rac{1}{2} ||x||_{l^2} \ge ||x||_{l^\infty}, \ (0, \ \dots, \ 0, \ x(m_0), \ 0, \ \dots) ext{ if } \ rac{1}{2} ||x||_{l^2} < ||x||_{l^\infty}, \end{cases}$$

where $m_0 = \inf \{m; x(m) = ||x||_{l^{\infty}}\}$ and $x(m_0)$ stands at the m_0 -th place, defines a duality function which is sequentially weakly continuous at zero. Another example of a space satisfying (I₀) but not (III) will be given in §5.

Although (II_0) does not generally imply (II), one can prove the following proposition which includes Opial's result of [17; lemma 3].

PROPOSITION 1. (II₀) implies (II) when X is uniformly convex.

This proposition as well as Opial's result are no longer true if the assumption of uniform convexity is weakened to that of local uniform convexity (see the example in §5).

Proof of Proposition 1. Let $x_n \rightarrow x_0$ and suppose that

 $\liminf ||x_n - x|| = \liminf ||x_n - x_0||$

with $x \neq x_0$. Then

$$\liminf \left\|x_n - rac{x+x_0}{2}
ight\| < \liminf \|x_n - x_0\|$$

by uniform convexity, which contradicts (II_0) .

We conclude this section with two results connecting (I) with two classical rotundity conditions.

PROPOSITION 2. If (I) holds then X has property (A), i.e., $x_n \rightarrow x$ whenever $x_n \rightarrow x$ and $||x_n|| \rightarrow ||x||$.

Proof. By Lemmas 1 and 2,

$$arPsi_{n}(||x_{n}-x||) = arPsi_{n}(||x_{n}||) + \int_{0}^{1} \langle -x, J_{\phi}(x_{n}-tx) \rangle dt$$

$$\mathbf{SO}$$

$$\lim \Phi(||x_n - x||) = \Phi(||x||) + \int_0^1 \langle -x, J_{\phi}(x - tx) \rangle dt .$$

Applying Lemma 2 to the last integral, we get $\lim \Phi(||x_n - x||) = 0$.

COROLLARY. If (I) holds and if X is strictly convex, then X has property (H), i.e., X has property (A) and is strictly convex.

PROPOSITION 3. If (I) holds and if X is strictly convex and reflexive, then X is locally uniformly convex.

Proof. We have to prove that $x_n \to x$ whenever $||x_n|| \to ||x||$ and $||x + x_n|| \to 2 ||x||$. Since the balls in X are weakly sequentially compact, it suffices to see that any weakly convergent subsequence $\{x_m\}$ converges in norm to x. Let $x_m \to y$. We will show that y = x, and the proof will follow from Proposition 2.

By Lemmas 1 and 2,

$$arPhi(||x_m+x||) - arPhi(||x_m||) = \int_0^1 < x, \, J_{\phi}(x_m+tx) > dt$$
 ;

going to the limit and applying Lemma 2 to the first member, we get

(3)
$$\int_0^1 \langle x, J_{\phi}(x + tx) \rangle dt = \int_0^1 \langle x, J_{\phi}(y + tx) \rangle dt.$$

The first integral equals $\int_0^1 ||x|| \phi(||x + tx||) dt$ and the second is majorized by

$$\int_{0}^{1} ||x|| \phi(||y + tx||) dt \leq \int_{0}^{1} ||x|| \phi(||x + tx||) dt$$

since $||y|| \leq ||x||$; consequently

$$\langle x, J_{\phi}(y + tx) \rangle = ||x|| \phi(||y + tx||) \quad \forall t \in [0, 1].$$

But the strict convexity of X means that every nonzero $x^* \in X^*$ assumes its norm at most one point of the unit sphere of X. Thus, eliminating the trivial case x = 0,

$$rac{x}{||x||} = rac{y+tx}{||y+tx||}$$

for all $t \in [0, 1]$ with $y + tx \neq 0$, an equality which gives y = kx with $k \ge 0$. It then follows from (3) that k = 1, which completes the proof.

5. An example. Consider the space c_0 . The formula

$$||x|| = \sup \left[\sum_{i=1}^\infty 2^{-2i} x^2(lpha_i)
ight]^{1/2}$$
 ,

where the supremum is taken over all permutations α of N, defines

on c_0 an equivalent norm [9] which is known to be locally uniformly convex [18]. We will show that c_0 endowed with this norm satisfies (I_0) but not (III).

First recall an equivalent definition of || || by means of the decreasing rearrangement map D of c_0 into l^2 (cf. [9]). Given $x \in c_0$, N can be enumerated in a sequence $\{\beta_i\}$ in such a way that $|x(\beta_i)| \ge |x(\beta_{i+1})|$ for all i and that $\beta_i \le \beta_j$ if $|x(\beta_i)| = x(\beta_j)|$ and $i \le j$. Define $(Dx)(\beta_i) = 2^{-i}x(\beta_i)$. Then $||x|| = ||Dx||_{l^2}$.

We assert that $D^2: c_0 \to l^i$ is a duality function with gauge $\phi(r) = r$ which is sequentially continuous at zero for $\sigma(c_0, l^1)$ and $\sigma(l^1, c_0)$. Indeed D^2x is defined by $(D^2x)(\beta_i) = 2^{-2i}x(\beta_i)$, and we have

$$<\!x,\,D^2\!x\!>=\sum\limits_{i=1}^\infty 2^{-2i}\!x^2(eta_i)=||\,Dx\,||_{l^2}^{_2}=||\,x\,||^2$$
 ,

while for all $y \in c_0$

by definition of the norm. Thus D^2 is a duality function. The continuity requirement is clearly satisfied.

To prove that (III) does not hold, we construct a diametral sequence $x_n \rightarrow 0$ (see the characterization of normal structure mentioned in the proof of Theorem 1). Take $x_1 = 0$ and, for n > 1,

$$x_n(i) = egin{cases} 0 & ext{if} \ i \leqq 2^{-1}n(n-1), \ 1 & ext{if} \ 2^{-1}n(n-1) < i \leqq 2^{-1}n(n-1) + n, \ 0 & ext{if} \ i > 2^{-1}n(n-1) + n \ . \end{cases}$$

It is immediate that $\langle x_n, x^* \rangle \rightarrow 0$ for each $x^* \in l^1$, thus $x_n \rightarrow 0$. On the other hand

$$\delta(\{x_n\})^2 = \sup_{n,m} ||x_n - x_m||^2 = \sup_{n,m} \sum_{i=1}^{n+m} 2^{-2i} = \sum_{i=1}^{\infty} 2^{-2i}$$

but

$$d(x_{n+1}, \operatorname{co} \{x_1, \, \cdots, \, x_n\})^2 \ge \sum_{i=1}^{n+1} 2^{-2i}
ightarrow \sum_{i=1}^{\infty} 2^{-2}i \; ,$$

and consequently the sequence is diametral.

References

1. E. Asplund, Positivity of duality mappings, Bull. Amer. Math. Soc., 73 (1967), 200-203.

2. N. A. Assad and W. A. Kirk, Fixed point theorems for set-valued mappings of contractive type, to appear.

3. L. P. Belluce, W. A. Kirk and E. F. Steiner, Normal structure in Banach spaces, Pacific J. Math., 26 (1968), 433-440.

4. M. S. Brodskii and D. P. Milman, On the center of a convex set, Dokl. Akad. Nauk. SSSR, **59** (1948), 837-840.

5. F. E. Browder, Fixed point theorems for nonlinear semicontractive mappings in Banach spaces, Arch. Rat. Mech. Anal., **21** (1966), 259-269.

6. _____, Nonlinear operators and nonlinear equations of evolution in Banach Spaces, Proc. Symp. Pure Math., 18, Part 2, Amer. Math. Soc., to appear.

7. R. E. Bruck, Approximating fixed points and fixed point sets of nonexpansive mappings in Banach spaces, Ph. D. Thesis, University of Chicago, 1969.

8. D. F. Cudia, The geometry of Banach spaces, Smoothness, Trans. Amer. Math. Soc., **110** (1964), 284-314.

9. M. M. Day, Strict convexity and smoothness of normed spaces, Trans. Amer. Math. Soc., 78 (1955), 516-528.

10. J. P. Gossez, A note on multivalued monotone operators, Michigan Math. J., 17 (1970), 347-350.

11. J. P. Gossez and E. Lami Dozo, Structure normale et base de Schauder, Bull. Cl. Sc. Ac. R. Belgique, 55 (1969), 673-681.

12. W. A. Kirk, A fixed point theorem for mappings which do not increase distances, Amer. Math. Monthly, **72** (1965), 1004-1006.

13. _____, Fixed point theorems for nonlinear nonexpansive and generalized contractive mappings, to appear.

14. E. Lami Dozo, Multivalued nonexpansive mappings and Opial's condition, to appear.

15. G. J. Minty, Monotone (nonlinear) operators in Hilbert space, Duke Math. J., **29** (1962), 341-346.

16. J. J. Moreau, Fonctionnelles convexes, Sém. Eq. Dér. Part., Collège de France, Paris, 1967.

17. Z. Opial, Weak convergence of the sequence of successive approximations for nonexpansive mappings, Bull. Amer. Math. Soc., **73** (1967), 591-597.

18. J. Rainwater, Local uniform convexity of Day's norm on $c_0(\Gamma)$, Proc. Amer. Math. Soc., **22** (1969), 335-339.

Received May 12, 1971. The first author was supported by the F.N.R.S. in Belgium. Part of the results herein are contained in the second author's doctoral thesis (University of Brussels, 1970).

UNIVERSITY OF BRUSSELS AND UNIVERSITY OF CHICAGO, UNIVERSITY OF BRUSSELS.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University Stanford, California 94305 J. DUGUNDJI

Department of Mathematics University of Southern California Los Angeles, California 90007

RICHARD ARENS University of California Los Angeles, California 90024

ASSOCIATE EDITORS

F. WOLF

E.F. BECKENBACH

B.H. NEUMANN

K. Yoshida

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON * * * * AMERICAN MATHEMATICAL SOCIETY NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index. to Vol. **39**. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The Pacific Journal of Mathematics is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

C. R. HOBBY University of Washington Seattle, Washington 98105

Pacific Journal of Mathematics

Vol. 40, No. 3 November, 1972

Wazir Husan Abdi, A quasi-Kummer function	521
Vasily Cateforis, Minimal injective cogenerators for the class of modules of	
zero singular submodule	527
W. Wistar (William) Comfort and Anthony Wood Hager, Cardinality of	
k-complete Boolean algebras	541
Richard Brian Darst and Gene Allen DeBoth, Norm convergence of	
martingales of Radon-Nikodym derivatives given a σ -lattice	547
M. Edelstein and Anthony Charles Thompson, Some results on nearest	
points and support properties of convex sets in $c_0 \dots \dots \dots$	553
Richard Goodrick, Two bridge knots are alternating knots	561
Jean-Pierre Gossez and Enrique José Lami Dozo, Some geometric properties	
related to the fixed point theory for nonexpansive mappings	565
Dang Xuan Hong, Covering relations among lattice varieties	575
Carl Groos Jockusch, Jr. and Robert Irving Soare, Degrees of members of Π_1^0	
classes	605
Leroy Milton Kelly and R. Rottenberg, Simple points in pseudoline	
arrangements	617
Joe Eckley Kirk, Jr., The uniformizing function for a class of Riemann	
surfaces	623
Glenn Richard Luecke, <i>Operators satisfying condition</i> (G_1) <i>locally</i>	629
T. S. Motzkin, On L(S)-tuples and l-pairs of matrices	639
Charles Estep Murley, The classification of certain classes of torsion free	
Abelian groups	647
Louis D. Nel, Lattices of lower semi-continuous functions and associated	
topological spaces	667
David Emroy Penney, II, <i>Establishing isomorphism between tame prime</i>	
knots in E^3	675
Daniel Rider, <i>Functions which operate on</i> $\mathcal{F}L_p(T)$, 1	681
Thomas Stephen Shores, <i>Injective modules over duo rings</i>	695
Stephen Simons, A convergence theorem with boundary	703
Stephen Simons, Maximinimax, minimax, and antiminimax theorems and a	
result of R. C. James	709
Stephen Simons, On Ptak's combinatorial lemma	719
Stuart A. Steinberg, <i>Finitely-valued f-modules</i>	723
Pui-kei Wong, Integral inequalities of Wirtinger-type and fourth-order	
elliptic differential inequalities	739
Yen-Yi Wu, Completions of Boolean algebras with partially additive	
oparators	
	753