

# Pacific Journal of Mathematics

**CONDITIONAL EXPECTATIONS ASSOCIATED WITH  
STOCHASTIC PROCESSES**

R. A. BROOKS

## CONDITIONAL EXPECTATIONS ASSOCIATED WITH STOCHASTIC PROCESSES

R. A. BROOKS

**A stochastic process,  $E[x(t, \cdot) | \mathcal{F}_t](\omega)$ , on a probability space  $(\Omega, \mathcal{A}, P)$  and an interval  $D$ , where  $x \in L_1(D \times \Omega)$  and  $\{\mathcal{F}_t, t \in D\}$  is an increasing collection of sigma-fields in  $\mathcal{A}$ , is considered. Sufficient conditions for the joint measurability of  $E[x(t, \cdot) | \mathcal{F}_t](\omega)$  in  $(t, \omega)$  are given, and if  $x \in L_2(D \times \Omega)$ , it is shown that, under certain fairly general conditions,  $E[x(t, \cdot) | \mathcal{F}_t](\omega)$  can be identified with the projection of  $x$  onto a certain subspace of the Hilbert space associated with  $L_2(D \times \Omega)$ . The results obtained herein have application in certain classes of stochastic optimization problems.**

**1. Introduction.** Interest in the application of mathematical analysis to problems associated with optimizing control systems has been significant in recent years. [16], [6], [7], [9], [13]. A particular class of stochastic optimization problems [1] which are of interest in certain control system applications can be abstractly formulated in the following manner.

Let  $(\Omega, \mathcal{A}, P)$  be a probability space and let  $(D, \mathcal{B}, m)$  be a measure space, where  $D = [a, b] \subset R$ ,  $\mathcal{B}$  is the collection of Borel measurable subsets of  $D$ , and  $m$  is a finite measure defined on  $(D, \mathcal{B})$ . Consider the product measure space  $(D \times \Omega, \mathcal{B} \times \mathcal{A}, m \times P)$ , and denote by  $L_2(D \times \Omega)$  the collection of real valued  $\mathcal{B} \times \mathcal{A}$ -measurable functions defined on  $D \times \Omega$  which are square-integrable with respect to  $m \times P$ . Let  $y(t, \omega)$  be a stochastic process on  $(\Omega, \mathcal{A}, P)$  and  $R$ , and let

$$Y(t, \omega) = [y(\tau, \omega), \tau < t], (t, \omega) \in R \times \Omega .$$

For each  $t \in R$ , denote by  $\mathcal{Y}_t$  the minimal sigma-field of subsets of  $\Omega$  with respect to which every element of the random vector  $Y(t, \cdot)$  is measurable, and observe that  $t', t'' \in D$ ,  $t' < t''$  implies  $\mathcal{Y}_{t'} \subset \mathcal{Y}_{t''}$ .

In the stochastic optimization application under consideration, the random vector  $Y(t, \cdot)$  is the observation available to a controller at time  $t \in D$ , and the control problem is the following: For an arbitrary but fixed element  $v \in L_2(D \times \Omega)$  determine a control  $u_0 \in L_2(D \times \Omega)$  with the property that  $u_0(t, \cdot)$  is  $\mathcal{Y}_t$ -measurable for each  $t \in D$  and

$$\int_{D \times \Omega} [u_0(t, \omega) - v(t, \omega)]^2 (m \times P)(d(t, \omega))$$

$$= \inf \left\{ \int_{D \times \Omega} [u(t, \omega) - v(t, \omega)]^2 (m \times P)(d(t, \omega)); u \in L_2(D \times \Omega), \right.$$

$$\left. u(t, \cdot) \text{ is } \mathcal{Y}_t\text{-measurable for each } t \in D \right\}.$$

The  $\mathcal{Y}_t$ -measurability condition is imposed in order that the minimizing control  $u_0(t, \cdot)$  be a Baire function of the observation vector  $Y(t, \cdot)$  for each  $t \in D$ .

Now let

$$\mathcal{M} = \{G \in \mathcal{B} \times \mathcal{A}; G_t \in \mathcal{Y}_t \text{ for all } t \in D\},$$

where  $G_t \equiv \{\omega \in \Omega; (t, \omega) \in G\}$ ,  $t \in D$ , and observe that  $\mathcal{M} \subset \mathcal{B} \times \mathcal{A}$  is a sigma-field. Consider the Hilbert space  $H$  of  $m \times P$ -equivalence classes in  $L_2(D \times \Omega)$ , and let  $S \subset H$  be the collection of  $m \times P$ -equivalence classes generated by the  $\mathcal{M}$ -measurable elements in  $L_2(D \times \Omega)$ . It is readily verified that  $S$  is a subspace of  $H$ , and hence  $P_S$ , the projection operator which maps  $H$  onto  $S$ , exists. Consequently, for each  $v \in H$ , there exists a unique  $u_0 \in S$  such that

$$\|u_0 - v\| = \inf \{\|u - v\|; u \in S\},$$

and, by the Projection theorem,

$$u_0 = P_S v.$$

Thus, ascertaining a solution to the stochastic optimization problem stated above is tantamount to identifying the projection operator  $P_S$ , and this is done in the following theorem, the proof of which will be given later.

**THEOREM 1.1.** *Let  $v \in L_2(D \times \Omega)$ . Then there exists an  $\mathcal{M}$ -measurable real-valued function defined on  $D \times \Omega$ , denoted by  $E[v(t, \cdot) | \mathcal{Y}_t](\omega)$ , which is a conditional expectation of  $v(t, \cdot)$  given  $\mathcal{Y}_t$  for  $m$ -a.e.  $t \in D$ , and moreover,*

$$(P_S v)(t, \omega) = E[v(t, \cdot) | \mathcal{Y}_t](\omega)$$

for  $m \times P$ -a.e.  $(t, \omega) \in D \times \Omega$ .

The crux of the proof of the above theorem is to prove the existence of the  $\mathcal{M}$ -measurable function  $E[v(t, \cdot) | \mathcal{Y}_t](\omega)$ . The main result obtained in the sequel is that an  $\mathcal{M}$ -measurable function  $E[v(t, \cdot) | \mathcal{Y}_t](\omega)$  does indeed exist. Other results which make use of this measurability are also presented.

The utility of the conditional expectation representation of the

operator  $P_s$  is twofold. First, some interesting and useful facts regarding the solution of the stochastic optimization problem can be deduced using properties of conditional expectations [1]; and second, the problem of calculating conditional expectations of the type encountered above has received much attention in the literature, and efficient computational algorithms have been developed in certain cases. [8], [10], [11], [12], [3], [15], [17], [5], [4].

**2. Preliminaries.** Let  $\{\mathcal{F}_t, t \in D\}$  be a collection of increasing sigma-fields in  $\mathcal{A}$ , i.e., for each  $t \in D$ ,  $\mathcal{F}_t \subset \mathcal{A}$  and  $t', t'' \in D$ ,  $t' < t''$  implies  $\mathcal{F}_{t'} \subset \mathcal{F}_{t''}$ . Let  $x \in L_1(D \times \Omega)$ . Then  $x(t, \cdot) \in L_1(\Omega)$  for  $m$ -a.e.  $t \in D$ , and consequently, [14], a conditional expectation  $E[x(t, \cdot) | \mathcal{F}_t](\omega)$  exists for  $m$ -a.e.  $t \in D$ . By defining  $E[x(t, \cdot) | \mathcal{F}_t](\omega)$  to be an arbitrary random variable for each  $t$  in the exceptional set,  $E[x(t, \cdot) | \mathcal{F}_t](\omega)$  can be extended to be a stochastic process on  $(\Omega, \mathcal{A}, P)$  and  $D$ , and this is the basis of the following definition.

**DEFINITION 2.1.** Let  $x \in L_1(D \times \Omega)$ . A conditional expectation of  $x$  with respect to  $\{\mathcal{F}_t, t \in D\}$ , denoted by  $E[x(t, \cdot) | \mathcal{F}_t](\omega)$ , is a stochastic process on  $(\Omega, \mathcal{A}, P)$  and  $D$  with the property that for  $m$ -a.e.  $t \in D$ ,

- (i)  $E[x(t, \cdot) | \mathcal{F}_t]$  is  $\mathcal{F}_t$ -measurable,
- (ii)  $E[x(t, \cdot) | \mathcal{F}_t] \in L_1(\Omega)$ ,
- (iii)  $\int_F E[x(t, \cdot) | \mathcal{F}_t](\omega) P(d\omega) = \int_F x(t, \omega) P(d\omega)$  for all  $F \in \mathcal{F}_t$ .

It has already been mentioned that a conditional expectation of  $x$  with respect to  $\{\mathcal{F}_t, t \in D\}$  exists, although not necessarily uniquely, for each  $x \in L_1(D \times \Omega)$ , and the question which is now asked is whether there exists a  $\mathcal{B} \times \mathcal{A}$ -measurable representation for the conditional expectation of  $x$  with respect to  $\{\mathcal{F}_t, t \in D\}$ . Before addressing this fundamental question, several preliminary results will be given.

**LEMMA 2.1.** Let  $x \in L_p(D \times \Omega)$ ,  $1 \leq p < \infty$ . If  $E[x(t, \cdot) | \mathcal{F}_t](\omega)$  is  $\mathcal{B} \times \mathcal{A}$ -measurable, then  $E[x(t, \cdot) | \mathcal{F}_t](\omega) \in L_p(D \times \Omega)$ . Moreover, if  $E[x(t, \cdot) | \mathcal{F}_t]'(\omega)$  is any other  $\mathcal{B} \times \mathcal{A}$ -measurable conditional expectation of  $x$  with respect to  $\{\mathcal{F}_t, t \in D\}$ , then

$$E[x(t, \cdot) | \mathcal{F}_t]'(\omega) = E[x(t, \cdot) | \mathcal{F}_t](\omega) \text{ } m \times P\text{-a.e.}$$

*Proof.* The uniqueness of  $\mathcal{B} \times \mathcal{A}$ -measurable representations for the conditional expectation of  $x$  with respect to  $\{\mathcal{F}_t, t \in D\}$  up to  $m \times P$ -equivalence on  $D \times \Omega$  is immediate, and an application of Jensen's inequality for conditional expectation, together with Tonelli's

theorem, proves  $E[x(t, \cdot) | \mathcal{F}_t](\omega) \in L_p(D \times \Omega)$ .

**LEMMA 2.2.** *Let  $\mathcal{F} \subset \mathcal{A}$  be a sigma-field and let  $x \in L_1(D \times \Omega)$ . Then there exists a  $\mathcal{B} \times \mathcal{F}$ -measurable stochastic process on  $(\Omega, \mathcal{A}, P)$  and  $D$ , denoted by  $E[x(t, \cdot) | \mathcal{F}](\omega)$ , which is a conditional expectation of  $x(t, \cdot)$  given  $\mathcal{F}$  for each  $t \in D$  such that  $x(t, \cdot) \in L_1(\Omega)$ .*

*Proof.* It is sufficient to prove that the lemma holds on  $L'_1(D \times \Omega)$ , where  $L'_1(D \times \Omega) = \{x \in L_1(D \times \Omega); x(t, \cdot) \in L_1(\Omega) \text{ for all } t \in D\}$ , and to this end let  $S$  be the collection of elements in  $L'_1(D \times \Omega)$  for which the lemma holds.

(1) To show that  $S$  contains the characteristic functions of measurable rectangles, observe that if  $F = B \times A$ , where  $B \in \mathcal{B}$ ,  $A \in \mathcal{A}$ , then  $\chi_F(t, \omega) = \chi_B(t)\chi_A(\omega)$ ,  $(t, \omega) \in D \times \Omega$ , and, for each  $t \in D$ ,

$$(1) \quad E[\chi_F(t, \cdot) | \mathcal{F}](\omega) = \chi_B(t)E[\chi_A | \mathcal{F}](\omega)$$

for  $P$ -a.e.  $\omega \in \Omega$ , where  $E[\chi_A | \mathcal{F}](\omega)$  is a conditional expectation of  $\chi_A$  given  $\mathcal{F}$ . Thus the right-hand side of (1) is a  $\mathcal{B} \times \mathcal{F}$ -measurable stochastic process on  $(\Omega, \mathcal{A}, P)$  and  $D$  which is a conditional expectation of  $\chi_F(t, \cdot)$  given  $\mathcal{F}$  for each  $t \in D$ .

(2) Observe that  $S$  is closed with respect to linear operations in the sense that if  $x_1(t, \omega), \dots, x_n(t, \omega) \in S$ ,  $c_1, \dots, c_n \in R$  and  $x(t, \omega) = \sum_{i=1}^n c_i x_i(t, \omega)$ ,  $(t, \omega) \in D \times \Omega$ , then  $x(t, \omega) \in S$ .

(3)  $S$  is also closed with respect to dominated convergence in a certain sense. To show this let  $\{x_n(t, \omega), n = 1, 2, \dots\} \subset S$ ,  $x(t, \omega), w(t, \omega) \in L'_1(D \times \Omega)$ , and suppose that  $x(t, \omega) = \lim_{n \rightarrow \infty} x_n(t, \omega)$  for all  $(t, \omega) \in D \times \Omega$ , and also that  $|x_n(t, \omega)| \leq w(t, \omega)$  for all  $(t, \omega) \in D \times \Omega$ ,  $n = 1, 2, \dots$ . It is now shown that  $x \in S$ . For each  $n = 1, 2, \dots$ , let  $E[x_n(t, \cdot) | \mathcal{F}](\omega)$  be a  $\mathcal{B} \times \mathcal{F}$ -measurable stochastic process on  $(\Omega, \mathcal{A}, P)$  and  $D$  which is also a conditional expectation of  $x_n(t, \cdot)$  given  $\mathcal{F}$  for each  $t \in D$ . Now for each  $t \in D$ ,

$$E[x(t, \cdot) | \mathcal{F}](\omega) = \lim_{n \rightarrow \infty} E[x_n(t, \cdot) | \mathcal{F}](\omega)$$

for  $P$ -a.e.  $\omega \in \Omega$  [2, p. 23]. Let  $z(t, \omega)$  be defined on  $D \times \Omega$  by

$$z(t, \omega) = \begin{cases} \lim_{n \rightarrow \infty} E[x_n(t, \cdot) | \mathcal{F}](\omega), & \text{when this limit exists,} \\ 0 & \text{, otherwise.} \end{cases}$$

Then  $z(t, \omega)$  is a  $\mathcal{B} \times \mathcal{F}$ -measurable stochastic process on  $(\Omega, \mathcal{A}, P)$  and  $D$ , and, for each  $t \in D$ ,  $z(t, \omega)$  is a conditional expectation of  $x(t, \cdot)$  given  $\mathcal{F}$ . Thus  $x \in S$ .

(4) Let  $\mathcal{C} = \{F \in \mathcal{B} \times \mathcal{A}; \chi_F \in S\}$ . From (1), it follows that  $\mathcal{C}$  contains the measurable rectangles, in  $\mathcal{B} \times \mathcal{A}$ , and (2) then implies

that  $\mathcal{C}$  contains the field generated by the measurable rectangles in  $\mathcal{B} \times \mathcal{A}$ . From (3) it follows that  $\mathcal{C}$  is a monotone class, and consequently, by the monotone class lemma,  $\mathcal{C} = \mathcal{B} \times \mathcal{A}$ .

(5) Let  $x \in L'_1(D \times \Omega)$ . Then there exists a sequence of simple functions  $\{x_n, n = 1, 2, \dots\} \subset L'_1(D \times \Omega)$  such that  $x_n^+(t, \omega) \uparrow x^+(t, \omega)$ ,  $x_n^-(t, \omega) \uparrow x^-(t, \omega)$ , as  $n \rightarrow \infty$ , for all  $(t, \omega) \in D \times \Omega$ . From (2), (4) it follows that  $\{x_n, n = 1, 2, \dots\} \subset S$ , and (3) then implies  $x \in S$ , which proves the lemma.

The next result is a Fubini-type theorem for conditional expectation.

LEMMA 2.3. *Let  $\mathcal{F} \subset \mathcal{A}$  be a sigma-field, let  $x \in L_1(D \times \Omega)$ , and let  $E[x(t, \cdot) | \mathcal{F}](\omega)$  be a  $\mathcal{B} \times \mathcal{F}$ -measurable conditional expectation of  $x(t, \cdot)$  given  $\mathcal{F}$  for m-a.e.  $t \in D$ . Then*

$$E\left[\int_D x(t, \cdot) m(dt) \mid \mathcal{F}\right](\omega) = \int_D E[x(t, \cdot) \mid \mathcal{F}](\omega) m(dt)$$

for P-a.e.  $\omega \in \Omega$ .

*Proof.* Immediate consequence of Fubini's theorem.

LEMMA 2.4. *Let  $D_0 \subset D$  be countable, and suppose that  $\mathcal{F}_t = \sigma\{\mathbf{U}_{t' < t} \mathcal{F}_{t'}\}$  for each  $t \in D \setminus D_0$ . Then for each  $A \in \mathcal{A}$ , there exists a  $\mathcal{B} \times \mathcal{A}$ -measurable conditional expectation of  $\chi_A$  with respect to  $\{\mathcal{F}_t, t \in D\}$ .*

*Proof.* Let  $S' \subset D$  be countable and dense in  $D$  and let  $S = S' \cup D_0 \cup \{a\}$ . Then  $S \subset D$  is also countable and dense in  $D$ . Let  $S = \{s_1, s_2, \dots\}$  be an arbitrary denumeration of  $S$  for which  $s_1 = a$ , and for each  $t \in D$  put  $s(t, n) = \max\{s_i \in S; s_i \leq t, i = 1, 2, \dots, n\}$ ,  $n = 1, 2, \dots$ .

Let  $A \in \mathcal{A}$  and for each  $t \in D$  let  $u(t, \omega)$  be a conditional expectation of  $\chi_A$  given  $\mathcal{F}_t$ . Then for each  $t \in D$  it is clear that  $|u(t, \omega)| < 1$  for P-a.e.  $\omega \in \Omega$ .

Let  $u_n(t, \omega) = u(s(t, n), \omega)$ ,  $(t, \omega) \in D \times \Omega$ ,  $n = 1, 2, \dots$ , and observe that  $\{u_n, n = 1, 2, \dots\}$  is a sequence of  $\mathcal{B} \times \mathcal{A}$ -measurable functions on  $D \times \Omega$ . Furthermore, for each  $n = 1, 2, \dots$ ,  $t \in D$ ,  $u_n(t, \cdot)$  is  $\mathcal{F}_t$ -measurable since  $s(t, n) \leq t$ .

Now if  $t \in S$ , then  $s(t, n) = t$  for  $n$  sufficiently large, and hence  $\lim_{n \rightarrow \infty} u_n(t, \omega) = u(t, \omega)$  for all  $\omega \in \Omega$ . On the other hand, if  $t \in D \setminus S$ , then  $s(t, n) \uparrow t$  as  $n \rightarrow \infty$ , and thus

$$\bigcup_{n=1}^{\infty} \mathcal{F}_{s(t,n)} = \bigcup_{t' < t} \mathcal{F}_{t'}$$

Therefore, by hypothesis,

$$\mathcal{F}_t = \sigma\{\dot{\bigcup}_{n=1}^{\infty} \mathcal{F}_{s(t,n)}\},$$

and consequently, [2, Theorem 4.3, p. 331],

$$u(t, \omega) = \lim_{n \rightarrow \infty} u_n(t, \omega)$$

for  $P$ -a.e.  $\omega \in \Omega$ . Hence for each  $t \in D$

$$(2) \quad u(t, \omega) = \lim_{n \rightarrow \infty} u_n(t, \omega)$$

for  $P$ -a.e.  $\omega \in \Omega$ .

By the Bounded Convergence theorem, it follows that for each  $t \in D$

$$\lim_{n \rightarrow \infty} \int_{\Omega} |u(t, \omega) - u_n(t, \omega)| P(d\omega) = 0,$$

and hence

$$\lim_{k, n \rightarrow \infty} \int_{\Omega} |u_k(t, \omega) - u_n(t, \omega)| P(d\omega) = 0.$$

For each  $t \in D$ ,  $n = 1, 2, \dots$ , let

$$v_n(t) = \sup_{k \geq n} \int_{\Omega} |u_k(t, \omega) - u_n(t, \omega)| P(d\omega).$$

Then  $\{v_n, n = 1, 2, \dots\}$  is a sequence of  $\mathcal{B}$ -measurable functions on  $D$ , and  $|v_n(t)| \leq 2$ ,  $t \in D$ ,  $n = 1, 2, \dots$ . Thus, applying the Bounded Convergence theorem once more, it follows that

$$\lim_{n \rightarrow \infty} \int_D v_n(t) m(dt) = 0,$$

and consequently,

$$(3) \quad \lim_{k, n \rightarrow \infty} \int_{D \times \Omega} |u_k(t, \omega) - u_n(t, \omega)| (m \times P)(d(t, \omega)) = 0.$$

Now let  $\mathcal{S} = \{F \in \mathcal{B} \times \mathcal{A}; F_t \in \mathcal{F}_t, t \in D\}$ , where  $F_t \equiv \{\omega \in \Omega; (t, \omega) \in F\}$ ,  $t \in D$ . It is readily verified that  $\mathcal{S}$  is a sigma-field, and it is also clear that  $\{u_n, n = 1, 2, \dots\}$  is a sequence of  $\mathcal{S}$ -measurable functions on  $D \times \Omega$ . Thus, by the Riesz-Fischer theorem, it follows from (3) that there exists a  $\mathcal{S}$ -measurable function  $w \in L_1(D \times \Omega)$  such that

$$\lim_{n \rightarrow \infty} \int_{D \times \Omega} |w(t, \omega) - u_n(t, \omega)| (m \times P)(d(t, \omega)) = 0,$$

and hence there exists a subsequence  $\{n_k\} \subset \{n\}$  such that

$$w(t, \omega) = \lim_{k \rightarrow \infty} u_{n_k}(t, \omega)$$

for  $m \times P$ -a.e.  $(t, \omega) \in D \times \Omega$ . Therefore, for  $m$ -a.e.  $t \in D$ ,

$$w(t, \omega) = \lim_{k \rightarrow \infty} u_{n_k}(t, \omega)$$

for  $P$ -a.e.  $\omega \in \Omega$ , and from (2) it follows that, for  $m$ -a.e.  $t \in D$ ,  $w(t, \omega) = u(t, \omega)$  for  $P$ -a.e.  $\omega \in \Omega$ . The  $\mathcal{S}$ -measurability of  $w$  implies  $w(t, \cdot)$  is  $\mathcal{F}_t$ -measurable for each  $t \in D$ , and consequently  $w$  is a  $\mathcal{B} \times \mathcal{A}$ -measurable conditional expectation of  $\chi_A$  with respect to  $\{\mathcal{F}_t, t \in D\}$ .

3. Main results. Let  $y(t, \omega)$  be a stochastic process on  $(\Omega, \mathcal{A}, P)$  and  $R$ , let

$$Y(t, \omega) = [y(\tau, \omega), \tau < t], (t, \omega) \in R \times \Omega,$$

and for each  $t \in R$ , let

$$\mathcal{Y}_t = \sigma\{Y(t, \cdot)\}.$$

LEMMA 3.1.  $\mathcal{Y}_t = \sigma\{\bigcup_{t' < t} \mathcal{Y}_{t'}\}$ ,  $t \in R$ .

*Proof.* Since  $\mathcal{Y}_t$  is a sigma-field and  $(\bigcup_{t' < t} \mathcal{Y}_{t'}) \subset \mathcal{Y}_t$ , it follows that  $\sigma\{\bigcup_{t' < t} \mathcal{Y}_{t'}\} \subset \mathcal{Y}_t$  for each  $t \in R$ .

Let  $t \in R$  be fixed. Then for each  $\tau < t$ , the random variable  $y(\tau, \cdot)$  is measurable with respect to  $\mathcal{Y}_{t'}$  for each  $t' > \tau$ , and hence  $y(\tau, \cdot)$  is measurable with respect to  $\sigma\{\bigcup_{t' < t} \mathcal{Y}_{t'}\}$ . Thus

$$\mathcal{Y}_t \equiv \sigma\{[y(\tau, \cdot), \tau < t]\} \subset \sigma\{\bigcup_{t' < t} \mathcal{Y}_{t'}\},$$

and this proves the lemma.

THEOREM 3.2. Let  $x \in L_1(D \times \Omega)$ . Then there exists a  $\mathcal{B} \times \mathcal{A}$ -measurable conditional expectation of  $x$  with respect to  $\{\mathcal{Y}_t, t \in D\}$ .

*Proof.* From Lemmas 2.4 and 3.1, it follows that the theorem holds for  $x = \chi_A$ ,  $A \in \mathcal{A}$ . By an argument which parallels the proof of Lemma 2.2, it can be shown that the theorem holds for all  $x \in L_1(D \times \Omega)$ .

Now let  $\mathcal{M} \subset \mathcal{B} \times \mathcal{A}$  be the sigma-field defined by

$$\mathcal{M} = \{G \in \mathcal{B} \times \mathcal{A}; G_t \in \mathcal{Y}_t \text{ for all } t \in D\},$$

and observe that Theorem 3.2 implies that for each  $x \in L_1(D \times \Omega)$  there exists an  $\mathcal{M}$ -measurable conditional expectation of  $x$  with respect to  $\{\mathcal{Y}_t, t \in D\}$ . Let  $H$  be the Hilbert space of  $m \times P$ -equivalence classes in  $L_2(D \times \Omega)$ , with innerproduct denoted by  $\langle \cdot, \cdot \rangle$ , let  $S \subset H$  be the collection of  $m \times P$ -equivalence classes generated by the  $\mathcal{M}$ -measurable elements in  $L_2(D \times \Omega)$ , and let  $P_S$  be the projection of  $H$  onto  $S$ .

**THEOREM 3.3.** *Let  $x \in L_2(D \times \Omega)$  and let  $E[x(t, \cdot) | \mathcal{Y}_t](\omega)$  be an  $\mathcal{M}$ -measurable conditional expectation of  $x$  with respect to  $\{\mathcal{Y}_t, t \in D\}$ . Then*

$$(P_S x)(t, \omega) = E[x(t, \cdot) | \mathcal{Y}_t](\omega)$$

for  $m \times P$ -a.e.  $(t, \omega) \in D \times \Omega$ .

*Proof.* The existence of an  $\mathcal{M}$ -measurable conditional expectation of  $x$  with respect to  $\{\mathcal{Y}_t, t \in D\}$  follows from Theorem 3.2, since  $L_2(D \times \Omega) \subset L_1(D \times \Omega)$ .

Let  $M \in \mathcal{M}$  and observe that

$$\begin{aligned} & \int_M [(P_S x)(t, \omega) - E[x(t, \cdot) | \mathcal{Y}_t](\omega)](m \times P)(d(t, \omega)) \\ &= \langle P_S x, \chi_M \rangle - \int_D \int_{M_t} E[x(t, \cdot) | \mathcal{Y}_t](\omega) P(d\omega) m(dt) \\ &= \langle x, \chi_M \rangle - \int_D \int_{M_t} x(t, \omega) P(d\omega) m(dt) \\ &= \int_M x(t, \omega)(m \times P)(d(t, \omega)) - \int_M x(t, \omega)(m \times P)(d(t, \omega)) \\ &= 0. \end{aligned}$$

From the arbitrariness of  $M \in \mathcal{M}$ , it follows that

$$(P_S x)(t, \omega) = E[x(t, \cdot) | \mathcal{Y}_t](\omega)$$

for  $m \times P$ -a.e.  $(t, \omega) \in D \times \Omega$ .

**4. Extensions.** The theorem which follows extends some of the results obtained in §§ 2, 3, and it has application in the study of the class of stochastic optimization problems defined in § 1.

**THEOREM 4.1.** *Let  $(D', \mathcal{B}')$  be a copy of  $(D, \mathcal{B})$ , let  $m'$  be an arbitrary finite measure on  $(D', \mathcal{B}')$ , and let  $x \in L_1(D' \times D \times \Omega)$ .*

Then there exists a  $\mathcal{B}' \times \mathcal{B} \times \mathcal{A}$ -measurable real-valued function defined on  $D' \times D \times \Omega$ , denoted by  $E[x(s, t, \cdot) | \mathcal{Y}_t](\omega)$ , which for m-a.e.  $t \in D$ , is a conditional expectation of  $x(s, t, \cdot)$  given  $\mathcal{Y}_t$  for every  $s \in D'$  such that  $x(s, t, \cdot) \in L_1(\Omega)$ . Furthermore, for every  $t \in D$ ,  $E[x(s, t, \cdot) | \mathcal{Y}_t](\omega)$  is  $\mathcal{B}' \times \mathcal{Y}_t$ -measurable on  $D' \times \Omega$ , and, for m-a.e.  $t \in D$ ,

$$E\left[\int_{D'} x(s, t, \cdot) m'(ds) | \mathcal{Y}_t\right](\omega) = \int_{D'} E[x(s, t, \cdot) | \mathcal{Y}_t](\omega) m'(ds)$$

for P-a.e.  $\omega \in \Omega$ .

*Proof.* The existence of  $E[x(s, t, \cdot) | \mathcal{Y}_t](\omega)$  with the stated conditional expectation and measurability properties is established by an extension of the proof of Lemma 2.2, and the validity of interchanging conditional expectation and integration is guaranteed by Lemma 2.3.

#### REFERENCES

1. R. A. Brooks, *Linear Stochastic Control: A Hilbert Space Measure Theoretic Approach*, Ph. D. Dissertation, University of California, Irvine, 1970.
2. J. L. Doob, *Stochastic Processes*, Wiley, New York, 1953.
3. J. R. Fisher and E. B. Stear, *Optimal nonlinear filtering for independent increment processes*, IEEE Trans. on Inform. Theory, IT-3 (October 1967), 558-578.
4. T. L. Gunkel III and G. F. Franklin, *A general solution for linear sampled-data control*, J. of Basic Engr., 85 (1963), 197-201.
5. P. D. Joseph and J. T. Tou, *On Linear Control Theory*, AIEE Trans. on Appl. and Indus., Part II, 80 (September 1961), 193-196.
6. R. E. Kalman, *On the General Theory of Control Systems*, Proc. First IFAC Congress, pp. 481-493.
7. ———, *Contributions to the theory of optimal control*, Bol. Soc. Mat. Mex., 5 (1960) 102-119.
8. ———, *A new approach to linear filtering and prediction problems*, J. of Basic Engr., 82 (March 1960), 35-45.
9. R. E. Kalman, et. al., *Controllability of Linear Dynamical Systems*, Contributions to Differential Equations, vol. 1, Wiley, New York, 1962.
10. H. J. Kushner, *On the dynamical equations of conditional probability functions, with applications to optimum stochastic control theory*, J. Math. Anal. & Appl., 8 (April 1964), 332-344.
11. ———, *On the differential equations satisfied by conditional probability densities of Markov processes, with applications*, SIAM J. of Control, 2 (1964), 106-119.
12. ———, *Dynamical equations for optimal nonlinear filtering*, J. of Diff. Eq., 3 (April 1967), 179-190.
13. J. P. LaSalle, *The Time Optimal Control Problem*, Contributions to Differential Equations, 5 (1960), 1-24. Princeton University Press, Princeton.
14. M. Loève, *Probability Theory*, 3rd ed., van Nostrand, Princeton, 1963.
15. R. E. Mortensen, *Optimal control of continuous-time stochastic systems*, Electronics, Research Lab. Tech. Report, ERL-66-1, Univ. of Calif., Berkeley, August 1966.
16. L. S. Pontryagin, et. al., *The Mathematical Theory of Optimal Processes*, Inter-

science, New York, 1962.

17. R. J. Stuart, *Continuous-time Nonlinear Filtering*, Ph. D. dissertation, UCLA, 1969.

Received January 28, 1971.

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

C. R. HOBBY  
University of Washington  
Seattle, Washington 98105

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

Anatole Beck and Peter Warren, <i>Weak orthogonality</i> .....	1
Jonnie Bee Bednar and Howard E. Lacey, <i>Concerning Banach spaces whose duals are abstract L-spaces</i> .....	13
Louis Harvey Blake, <i>Canonical extensions of measures and the extension of regularity of conditional probabilities</i> .....	25
R. A. Brooks, <i>Conditional expectations associated with stochastic processes</i> .....	33
Theodore Allen Burton and Ronald Calvin Grimmer, <i>On the asymptotic behavior of solutions of <math>x'' + a(t)f(x) = e(t)</math></i> .....	43
Stephen LaVern Campbell, <i>Operator-valued inner functions analytic on the closed disc</i> .....	57
Yuen-Kwok Chan, <i>A constructive study of measure theory</i> .....	63
Alexander Munro Davie and Bernt Karsten Oksendal, <i>Peak interpolation sets for some algebras of analytic functions</i> .....	81
H. P. Dikshit, <i>Absolute total-effective <math>(N, p_n)(c, 1)</math> method</i> .....	89
Robert E. Edwards, Edwin Hewitt and Kenneth Allen Ross, <i>Lacunarity for compact groups. II</i> .....	99
James Daniel Halpern, <i>On a question of Tarski and a maximal theorem of Kurepa</i> .....	111
Gerald L. Itzkowitz, <i>A characterization of a class of uniform spaces that admit an invariant integral</i> .....	123
Mo Tak Kiang, <i>Semigroups with diminishing orbital diameters</i> .....	143
Glenn Richard Luecke, <i>A class of operators on Hilbert space</i> .....	153
R. James Milgram, <i>Group representations and the Adams spectral sequence</i> .....	157
G. S. Monk, <i>On the endomorphism ring of an abelian p-group, and of a large subgroup</i> .....	183
Yasutoshi Nomura, <i>Homology of a group extension</i> .....	195
R. Michael Range, <i>Approximation to bounded holomorphic functions on strictly pseudoconvex domains</i> .....	203
Norman R. Reilly, <i>Inverse semigroups of partial transformations and <math>\theta</math>-classes</i> .....	215
Chris Rorres, <i>Strong concentration of the spectra of self-adjoint operators</i> .....	237
Saharon Shelah, <i>A combinatorial problem; stability and order for models and theories in infinitary languages</i> .....	247
George Gustave Weill, <i>Vector space decompositions and the abstract imitation problem</i> .....	263
Arthur Thomas White, <i>On the genus of the composition of two graphs</i> .....	275