

Pacific Journal of Mathematics

GROUP REPRESENTATIONS AND THE ADAMS SPECTRAL SEQUENCE

R. JAMES MILGRAM

GROUP REPRESENTATIONS AND THE ADAMS SPECTRAL SEQUENCE

R. JAMES MILGRAM

Homotopy groups admit primary operations analogous to the Steenrod operations in ordinary cohomology theory and a good understanding of them seems vital to interpreting patterns in the homotopy of spheres.

Also, it has been known for a long time that a type of Steenrod algebra acts in $\text{Ext}_A(Z_p, Z_p)$ if A is a cocommutative Hopf algebra. Recently, D. S. Kahn showed that in the E_2 term of the Adams spectral sequence $\text{Ext}_{\mathcal{A}(2)}^{**}(Z_2, Z_2)$, certain of these operations on infinite cycles converge to the graded elements associated to the actual homotopy operations. Also, on infinite cycles, he showed how this structure determined some differentials.

In this paper, we further explore the relations between the operations in $\text{Ext}_{\mathcal{A}(p)}^{**}(Z_p, Z_p)$ and differentials in the Adams spectral sequence. In particular, for elements which need not be infinite cycles, we prove

THEOREM 4.1.1. (a) There are operations Sq^i in $\text{Ext}_{\mathcal{A}(2)}(Z_2, Z_2)$ so that

$$\partial_2(Sq^i(a)) = \begin{cases} h_0 Sq^{i+1}(a), & i \equiv s(2) \\ 0 & \text{otherwise} \end{cases}$$

for $a \in \text{Ext}_{\mathcal{A}(2)}^{r,s}(Z_2, Z_2)$.

(b) There are operations $\mathcal{P}^i, \beta \mathcal{P}^i$ in $\text{Ext}_{\mathcal{A}(p)}(Z_p, Z_p)$ for p an odd prime so that

$$\partial_2(\mathcal{P}^i(a)) = \alpha_0 \beta \mathcal{P}^i(a),$$

for $a \in \text{Ext}_{\mathcal{A}(p)}^{r,s}(Z_p, Z_p)$. (Here, Sq^i takes $\text{Ext}^{s,r}$ homomorphically to $\text{Ext}^{s+i,2r}$ while \mathcal{P}^i takes $\text{Ext}^{s,r}$ to $\text{Ext}^{s+(2i-r)(p-1),pr}$, and $\beta \mathcal{P}^i(\text{Ext}^{s,r}) \subset \text{Ext}^{s+(2i-r)(p-1)+1,pr}$.)

These operations are readily computable in the Ext groups. (Methods for calculating them are given in §6 and [2], [19].) For example, $Sq^0(h_i) = h_{i+1}$, $Sq^1(h_i) = h_i^2$ where h_i is the nonzero element in $\text{Ext}^{1,2^i}$ dual to Sq^{2^i} . (Our notation for elements in Ext is that of [13].) Applying 4.1.1, $\partial_2(h_i) = h_0 h_{i-1}^2$, which is nonzero for i greater than 3. Consequently, we recover

COROLLARY (Adams). *An element of Hopf invariant one (mod 2) exists in $\pi_{n+s}(S^n)$ if and only if $n > s$ and $s = 1, 3, 7$.*

(The Hopf invariant of a homotopy class α is nonzero in Z_p -cohomology if and only if α is represented by an infinite cycle in

Ext^1 .)

Similarly, the mod p operations can be calculated. We have ([12])

$$\mathcal{P}^{i(p-1)}(h_i) = h_{i+1},$$

and $\alpha_0 \beta \mathcal{P}^{i(p-1)}(h_i) \neq 0$ for $i \geq 0$. Thus we recover the result of Liulevicius and Shimada on the elements of mod p Hopf invariant one.

In § 6, we calculate the complete action of the Sq^i operations in $\text{Ext}_{\mathcal{A}(2)}^{s,t}(Z_2, Z_2)$ for $t - s < 42$. It turns out that routine relations among the various classes, together with the differentials of 4.1.1(a), determine all ∂_2 differentials in this range. (I am indebted to M. Tangora for showing me how to obtain some of the more obscure ∂_2 differentials in this way.)

One surprising result is that $\partial_2(c_2) = h_0 f_1$, a differential in the 41 stem which was overlooked in [13]. This differential, in turn, implies $\nu \theta_4 \neq 0$ (which contradicts a result in [13]) where θ_4 is the class corresponding to the Kervaire invariant 1 manifold in dimension 30. This in turn forces a ∂_3 differential in the 34 stem. Once these two differentials are accounted for, there seem to be no further corrections necessary in [13], and we can thus assume the two primary components of $\pi_t^s(S^0)$ known for $t < 44$. For further details, see [5].

Perhaps equally surprising, the technique used to prove Theorem 4.1.1 is purely geometric in nature. We never need mention secondary cohomology operations or even primary ones.

Next, we study some of the ways in which the operations imply higher order differentials. Here the answers are not as satisfactory as before. However, we do succeed in characterizing all primary differentials through ∂_{20} on these elements. More exactly, Theorem 5.1.1 characterizes the first possible nonzero $\partial_k(Sq^i(a))$ for $k < 20$, provided $\partial_j(a) = 0$ for $j \leq k + 1$. However, there are no places in the first 40 stems where such differentials occur (except, indirectly, the differential $\partial_3(r) = h_1 d_3^2$), so in the absence of examples, it seemed fruitless to pursue the matter further.

Also, there are further applications of these geometric techniques. For example, in § 7 we give a very direct proof that θ_4 is nonzero. Moreover, by using similar techniques with the two-cell-complex $S^{15} \cup {}_2e^{16}$, together with the fact that $(\theta_3)^2 = (\sigma)^4 = 0$, one obtains a proof that θ_5 is nonzero. Similarly, using the result of Barratt-Mahowald that $(\theta_4)^2 = 0$, one obtains the existence of θ_6 (see [22] for details).

Finally, I would like to take this opportunity to express my thanks to D. S. Kahn for sharing his insights with me, and the Centro de Investigación for their support while this research was carried out.

1. Group representations and their Thom spaces.

1. We consider representations

$$r: G \longrightarrow L$$

where G is a finite group and L is the unitary group $U_{(n)}$ or the orthogonal group $O_{(n)}$. Any such representation gives rise to a map of classifying spaces

$$B_{(r)}: B_G \longrightarrow B_L$$

see e.g., [18]. By the Steenrod classification theorem $B_{(r)}$ is equivalent to specifying a principal L -bundle or its associated n -plane bundle over B_G . We denote this n -plane bundle by $B_{(r)}^*(\xi_n)$ where ξ_n is the universal bundle over B_L .

DEFINITION 1.1. The representation r is said to have L -filtration k if the associated map $B_{(r)}$ is homotopically trivial on the $k-1$ skeleton of B_G .

This is equivalent, of course, to saying $B_{(r)}^*(\xi_n)$ is trivial over the $k-1$ skeleton of B_G and may also be rephrased as

PROPOSITION 1.2. *Let E_G be the universal covering space of B_G ; then the representation r has filtration k if and only if there is an r -equivariant map*

$$\Psi: (E_G)_{k-1} \longrightarrow L$$

where $(E_G)_{k-1}$ is the $k-1$ skeleton of E_G and r -equivariant means $\Psi(gx) = r(g)\Psi(x)$ for all $x \in (E_G)_{k-1}$.

Proof. Let E_L be the universal principal L -bundle over B_L . Given a representation r this induces an r -equivariant map

$$\varphi: E_G \longrightarrow E_L;$$

moreover, by standard arguments (since E_L is contractible and E_G is G -free) any 2 such are equivariantly homotopic. Likewise, given any r -equivariant map $\Psi: (E_G)_{k-1} \rightarrow E_L$ it can be extended to an r -equivariant map φ . The induced map of classifying spaces $\bar{\varphi}$ will send the $k-1$ skeleton of B_G to $*$ in B_L . To obtain the converse note that $B_{(r)}$ is covered by an equivariant map $E_G \rightarrow E_L$ and use the covering homotopy property.

2. By use of the Whitney sum maps

$$\mu_{ij}: L_i \times L_j \longrightarrow L_{i+j}$$

we can define Whitney sums of representations ($r \oplus r'$ is the composition

$$G \xrightarrow{\Delta} G \times G \xrightarrow{r \times r'} L_i \times L_j \rightarrow L_{i+j} .$$

In particular, the positive integral multiples of r are defined and we have

THEOREM 1.2.1. *Let $L_i = U_{(i)}$ and suppose the order of G is s . Suppose $r: G \rightarrow U_{(i)}$ has filtration $k < 2i$, then sr has filtration $k + 2$.*

Proof. The obstruction to compression of the k -skeleton is an element σ in $H^k(B_G, \pi_k(B_U))$. By naturality the obstruction to compressing sr is $s\sigma$. However, $|G| \cdot H^*(B_G, Z) = 0$ so $s\sigma = 0$ and the theorem follows.

Similarly in the case $L_{i'} = O_i$ we have

THEOREM 1.2.2. *Let $m = (2|G|)^{2s} 2^{\delta_1} |G|^{\delta_2}$ with $0 \leq \delta_2 \leq \delta_1 - 1$, $0 \leq \delta_1 \leq 2$ and let $r: G \rightarrow O_n$ be any representation, then mr has filtration at least $8s + 2^{\delta_1} + 4^{\delta_2} - 1$ in B_0 .*

3. We now consider the Thom complex of the associated n -plane bundle $B_{(r)}^*(\xi_n)$. It admits a very simple description.

DEFINITION 1.3.1. Let B be a space with base point $*$, and A be any space, then the half smash product

$$A \bowtie B$$

is $A \times B/A \times *$.

A representation $r: G \rightarrow L$ induces an action of G on S^{2k_n} ($k = 0$ if $L = 0_{(n)}$, $k = 1$ if $L = U_{(n)}$) induced from the action of L on $S^{2k_n} = R^{2k_n} \cup \infty$. In terms of this action we have

THEOREM 1.3.2. *The Thom complex of $B_{(r)}^*(\xi_n)$ is homeomorphic to*

$$E_G \bowtie_G S^{2k_n} .$$

Proof. The associated bundle $B_{(r)}^*(\xi_n)$ can also be described as

$$E_G \times_G R^{2k_n}$$

where the action of G is that given above. Also, the associated sphere bundle admits a similar description. Hence, the theorem follows on equating the sphere bundle to the base point and using the fact that L acts linearly in R^{2k_n} .

As an immediate corollary we have

THEOREM 1.3.3. *If r has filtration p there is a G -equivariant map*

$$\varphi: (E_G)_{p-1} \times S^{2k_n} \longrightarrow S^{2k_n}$$

which is of degree 1 on the bottom cell (i.e., $\varphi(xg, y) = \varphi(x, gy)$).

Proof. If r has filtration p then $B_{(r)}^*(\xi_n)$ is trivial on the $p-1$ skeleton of B_G , so there is a bundle mapping

$$\rho: B_{(r)}^*(\xi_n) | (B_G)_{p-1} \longrightarrow R^{2k_n}$$

and an induced map Thom spaces $T(\rho): (E_G)_{p-1} \times_G S^{2k_n} \rightarrow S^{2k_n}$. Now φ is obtained by composing with the map

$$\pi: E_G \times S^{2k_n} \longrightarrow E_G \times S^{2k_n}.$$

4. In order to obtain alternate descriptions of these spaces we need to observe that two bundles will have homeomorphic Thom spaces if the bundles are isomorphic. In particular for representations r_1, r_2 this is equivalent to requiring that $B_{(r_1)} \simeq B_{(r_2)}$.

PROPOSITION 1.4.1. *$B_{(r_1)} \simeq B_{(r_2)}$ if there is an element a in the identity component of L so $r_1(g) = a^{-1}r_2(g)a$ for all $g \in G$.*

Proof. Let P be a path from the identity to a , then $P(t)^{-1}r_2(g)P(t)$ provides an equivariant homotopy from r_1 to r_2 and hence a homotopy of $B_{(r_1)}$ to $B_{(r_2)}$.

Since U_n is connected we can apply 1.4.1 without difficulty and we find that the Thom complex of $B_{(r)}^*(\xi_n)$ is homeomorphic to the Thom complex of a sum of irreducible representations which can usually be calculated.

Over O_n , however, we must be careful to check that the a used to make r_1 equivalent to r_2 has determinant $+1$ and not -1 .

2. Some explicit representations and their Thom complexes.

1. Let Z_q be a finite cyclic group with generator T and let R^i be the representation sending T to $\rho = e^{2\pi i/q}$. Then R^i is the i^{th} tensor product of R^1 with itself and sends T to ρ^i . These R^i form a complete set of irreducible representations of Z_q and any unitary representation is uniquely equivalent to a sum of these.

Let Z_q act on S^{2n-1} by $T(z_1 \cdots z_n) = (\rho^{i_1} z_1, \dots, \rho^{i_n} z_n)$. The resulting quotient space is the Lens space

$$L_q(i_1, \dots, i_n).$$

We define a classifying space construction for Z_q by letting B_{Z_q} be the limit of the $L_q(1, \dots, 1)$.

THEOREM 2.1.1. *The Thom space of $m_0(I) + m_1R^1 + \dots + m_{q-1}R^{q-1}$ over B_{Z_q} is homeomorphic to*

$$\lim_{s \rightarrow \infty} \Sigma^{m_0} \left(\frac{L_q(\underbrace{1 \dots 1}_{m_1} \underbrace{2 \dots 2 \dots}_{m_2} \underbrace{(q-1, \dots, q-1)}_{m_{q-1}}, \underbrace{1 \dots 1}_s)}{L_q(\underbrace{1 \dots 1}_{m_1} \dots \underbrace{q-1, \dots, q-1}_{m_{q-1}})} \right)$$

(Here I is the trivial 1-dimensional representation.)

Proof. Embed S^{2k-1} in S^{2n-1} as the points $(0 \dots 0, z_1 \dots z_k)$. Define an action of Z_q on S^{2m-1} as $T(z_1 \dots z_{n-k}, \hat{z}_1 \dots \hat{z}_k) = (\rho^{i_1} z_1 \dots \rho^{i_{n-k}} z_{n-k}, \rho \hat{z}_1 \dots \rho \hat{z}_k)$. The embedding induces an inclusion $L_q(1 \dots 1) \subset L_q(i_1, \dots, i_{n-k}, 1 \dots 1)$ with normal bundle $R^{i_1} \oplus \dots \oplus R^{i_{n-k}}$. It is now evident that the Thom complex of $R^{i_1} \oplus \dots \oplus R^{i_{n-k}}$ is $L_q(i_1 \dots i_{n-k}, 1 \dots 1)/L_q(i_1 \dots i_{n-k})$ since the normal bundle maps homeomorphically till it arrives at $L_q(i_1 \dots i_{n-k})$, which is then the image of the sphere bundle.

For Z_2 and real representations we have 2 irreducible ones, the trivial one I and the identity $R: Z_2 \rightarrow Z_2 = 0(1)$, so we have

THEOREM 2.1.2. *Let RP^∞ be B_{Z_2} , then the Thom space of $m_0I + m_1R$ is homeomorphic to*

$$\Sigma^{m_0}(RP^\infty/RP^{m_1-1}).$$

(The proof does not differ essentially from that of 2.1.1.)

2. We recall the characteristic classes of these representations.

THEOREM 2.2.1. *Let $\iota \in H^2(B_{Z_q}, Z)$ be the first Chern class of R^1 , then the total Chern class of R^k is $1 + k\iota$. (This allows us to calculate, if necessary, the action of the Steenrod algebra $\mathcal{A}(p)$ in the Thom spaces above.)*

3. Consider the representation $r_n: Z_2 \rightarrow O_{2n}$ defined by $r_n(T)(x, y) = (y, x)$. From 1.3.2 the Thom complex of r_n is $E_{Z_2} \times_T S^n \wedge S^n = \lim_{m \rightarrow \infty} S^m \times_T S^n \wedge S^n$, where T acts as the antipodal map on S^m and as the interchange map on $S^n \wedge S^n$.

PROPOSITION 2.3.1. *r_n is equivalent to $nR \oplus nI$. Thus $\Sigma^n RP^\infty/RP^{n-1}$ is homeomorphic to $E_{Z_2} \times_T S^n \wedge S^n$.*

Proof. By changing coordinates we can assume $r_n(T)$ has the form

$$\begin{pmatrix} M & 0 & 0 & 0 & 0 \\ 0 & M & 0 & \cdot & \cdot \\ \cdot & \cdot & M & \cdot & 0 \\ 0 & \cdot & \cdot & \cdot & M \end{pmatrix} \text{ where } M \text{ is the matrix } \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

But $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and 2.3.1 follows.

Similarly we can consider the complex representation $r_n: Z_q \rightarrow U_{q^n}$ where $r(T)(x_1 \cdots x_q) = (x_q, x_1, x_2 \cdots x_{q-1})$. Its Thom complex is $E_{Z_q} \times_{Z_q} S^n \wedge \cdots \wedge S^n = \lim_{m \rightarrow \infty} S^{2m-1} \times_{Z_q} S^n \wedge \cdots \wedge S^n$ where Z_q acts on S^{2m-1} by $T(z_1 \cdots z_m) = (\rho z_1, \dots, \rho z_m)$ and by the shift map in $S^n \wedge \cdots \wedge S^n$.

PROPOSITION 2.3.2. r_n is equivalent to $n[I \oplus R^1 \oplus R^2 \oplus \cdots \oplus R^{q-1}]$ so $E_{Z_q} \times_{Z_q} S^n \wedge \cdots \wedge S^n$ is homeomorphic to

$$\frac{\Sigma^{2n} L_p(\underbrace{1 \cdots 1}_n, \cdots, \underbrace{q-1, \cdots, q-1}_n, 1, 1 \cdots)}{L_p(1, \cdots, 1, \cdots, q-1, \cdots, q-1)}.$$

Proof. The characteristic polynomial of $r(T)$ is $(1 - \lambda^q)^n$. Moreover, by changing coordinates T becomes $\begin{pmatrix} M & 0 & \cdot & 0 \\ 0 & M & 0 & 0 \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 0 & M \end{pmatrix}$ where M

is the matrix $\begin{pmatrix} 0 & 1 & 0 & \cdot & \cdot \\ 0 & 0 & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 \\ 1 & 0 & \cdot & \cdot & 0 \end{pmatrix}$. Thus, since $1 - \lambda^q$ has q distinct roots,

M can be diagonalized to $\begin{pmatrix} 1 & 0 & \cdot & 0 \\ 0 & \rho & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot \\ 0 & \cdot & \cdot & \rho^{q-1} \end{pmatrix}$ and 2.3.1 follows.

4. As a final example we consider the symmetric group \mathcal{S}_4 and the representation $r_n: \mathcal{S}_4 \rightarrow O_{4n}$ defined by

$$r(\alpha)(x_1 \cdots x_4) = (x_{\alpha(1)} \cdots x_{\alpha(4)}).$$

The Z_2 cohomology of $B_{\mathcal{S}_4}$ is $P(A, B, C)/AC = 0$ where $P(\)$ is a polynomial ring on three generators with A of dimension 3, B of dimension 2 and C of dimension 1. Let K be the Klein group. The inclusion $i: K \rightarrow \mathcal{S}_4$ induces the cohomology map $i^*: H^*(\mathcal{S}_4) \rightarrow H^*(K) = P(e_1, e_2)$ which sends $A \rightarrow e_1^2 e_2 + e_1 e_2^2$, $B \rightarrow e_1^2 + e_1 e_2 + e_2^2$, $C \rightarrow 0$. Next let M be the group $Z_2 \oplus Z_2$ and $j: Z_2 \oplus Z_2 \rightarrow \mathcal{S}_4$ be given by $j(T_1) = (12)$, $j(T_2) = (34)$. Then $j^*(A) = 0$, $j^*(B) = e_1 e_2$, $j^*(C) = e_1 + e_2$.

THEOREM 2.4.1. The total Stiefel-Whitney class of r_n is $(1 + C + B + A)^n$.

Proof. Clearly r_n is equivalent to nr_1 . Hence we show the Stiefel-Whitney class of r_1 is $(1 + C + B + A)$.

From the character tables we find

$$\begin{aligned} W(i^*r_1) &= (1 + e_1)(1 + e_2)(1 + e_1 + e_2) \\ &= 1 + i^*(B) + i^*(A) . \end{aligned}$$

Also $W(j^*(r_1)) = (1 + e_1)(1 + e_2) = 1 + j^*(C) + j^*(B)$. Now the result follows since

$$i^* \oplus j^*: H^*(\mathcal{S}_1) \longrightarrow H^*(K) \oplus H^*(Z_2 \oplus Z_2)$$

is a monomorphism.

Thus, for example, the first 7 Stiefel-Whitney classes of $r_{2'+7}$, $\iota > 3$ are $C, B + C^2, A + C^3, C^4 + C^2B, C^5 + CB^2, BC^4 + B^3 + C^6 + A^2, AB^2 + C^7$.

By generalizing slightly the techniques of [6] we can use these results to obtain a complete description of $H^*(\mathcal{S}_n, Z_2)$ for all n (see e.g. [20, §9]) as well as the structure of the corresponding Thom complexes

$$E_{\mathcal{S}_n} \bowtie_{\mathcal{S}_n} \underbrace{S^m \wedge \cdots \wedge S^m}_{n\text{-times}} .$$

We omit the details.

3. The geometry of the action of the Steenrod algebra on $\text{Ext}_{\mathcal{N}(p)}(Z_p, Z_p)$.

1. We start by recalling some well known facts from homological algebra. Throughout this paragraph we assume given a fixed augmented Hopf algebra A over a ground ring Γ (Z or Z_p for some prime p) and we require that the coproduct

$$\psi: A \longrightarrow A \otimes A$$

be coassociative and cocommutative. We have

THEOREM 3.1.1. *Suppose \mathcal{C} is an A -free resolution of the augmentation, then for each subgroup G of the symmetric group \mathcal{S}_n there is a chain map*

$$\varphi_G: \mathcal{E}_G \otimes \mathcal{C} \longrightarrow \underbrace{\mathcal{C} \otimes \cdots \otimes \mathcal{C}}_{n\text{-times}}$$

(where \mathcal{E}_G is a $\Gamma(G)$ free resolution of the augmentation $\Gamma(G) \rightarrow \Gamma$), and

$$\begin{aligned} \varphi_G(\omega \otimes ac) &= \psi^{(n)}(a)\varphi_G(\omega \otimes c) \\ * \quad \varphi_G(g\omega \otimes c) &= g\varphi_G(\omega \otimes c) . \end{aligned}$$

Moreover, given any other map φ'_G satisfying $*$ there is an equivariant

chain homotopy H between φ_a and φ'_a so that $H(\omega \otimes ac) = \Psi^{(n)}(a)H(\omega \otimes c)$.

Proof. (Compare [12]). Since $\Psi^{(n)}: A \rightarrow A^n$ is a map of algebras it induces a chain map

$$\varphi_0: \mathcal{C} \longrightarrow \mathcal{C}^{(n)}$$

and $\varphi_0(ac) = \Psi^{(n)}(a)\varphi_0(c)$. Now $*$ tells how to extend φ_0 over $(\mathcal{E}_G)_0 \otimes \mathcal{C}$. Note that since Ψ is cocommutative and coassociative, $g\Psi^n = \Psi^{(n)}$ for each g in G . Thus the various $g\varphi_0$ are all homotopic, and $\varphi_0(\alpha \times \mathcal{C})$ is homotopic to 0 when regarded as a map $\varphi_\alpha: \mathcal{C} \rightarrow \mathcal{C}^{(n)}$ if α is in the kernel of the augmentation. Thus we can extend over the 1 skeleton of \mathcal{E}_G . The remaining arguments are equally direct.

Now, in the standard way (e.g. as in [25, Chapt. VII]) we can construct “cohomology operations” from the map $\bar{\varphi}_G: \bar{\mathcal{E}}_G \otimes \bar{\mathcal{C}} \rightarrow \bar{\mathcal{C}}^{(n)}$ when $\bar{\mathcal{C}}$ is the reduced resolution of A . In particular this gives us operations coming from the cohomology of the symmetric group which have formal properties analogous to those of the ordinary Steenrod operations. Explicitly

THEOREM 3.1.2. *Let $\Gamma = Z_p$ for p an odd prime, then the φ_a allow one to define operations $\mathcal{P}^i, \beta\mathcal{P}^i$ in $\text{Ext}^{**}_A(Z_p, Z_p)$ with the following properties.*

- (i) $\mathcal{P}^k: \text{Ext}^{i,j} \rightarrow \text{Ext}^{i+(2k-j)(p-1), pj}$
 $\beta\mathcal{P}^k: \text{Ext}^{i,j} \rightarrow \text{Ext}^{i+(2k-j)(p-1)+1, pj}$
- (ii) $\mathcal{P}^j, \beta\mathcal{P}^j$ are homomorphisms
- (iii) Cartan formulae
 $\mathcal{P}^i(ab) = \sum_{i \geq j \geq 0} \mathcal{P}^j(a) \mathcal{P}^{i-j}(b)$
 $\beta\mathcal{P}^i(ab) = \sum \beta\mathcal{P}^ja \mathcal{P}^{i-j}b + \sum (-1)^{(a)} \mathcal{P}^ja \beta\mathcal{P}^{i-j}u$
- (iv) Adem formulae

$$\begin{aligned} \mathcal{P}^a \mathcal{P}^b &= \sum_{t=0}^{[a/p]} (-1)^{a+t} \binom{(p-1)(b-t)-1}{a-pt} \mathcal{P}^{a+b-t} \mathcal{P}^t \\ \mathcal{P}^a(\beta\mathcal{P}^b) &= \sum_{t=0}^{[a/p]} (-1)^{a+t} \binom{(p-1)(b-t)}{a-pt} (\beta\mathcal{P}^{a+b-t}) \mathcal{P}^t \\ &\quad + \sum_{t=0}^{[(a-1)/p]} (-1)^{a+t+1} \binom{(p-1)(b-t)-1}{a-pt-1} \mathcal{P}^{a+b-t}(\beta\mathcal{P}^t) \end{aligned}$$

- (v) Normation $\mathcal{P}^k(a) = a^p$ if $a \in \text{Ext}^{i,j}$, and $2k = i + j$,
 $\mathcal{P}^k(a) = 0$ if $2k < j$.

THEOREM 3.1.3. *For $p = 2$ there are homomorphisms $Sq^i: \text{Ext}^{**}_A(Z_2, Z_2) \rightarrow \text{Ext}^{**}_A(Z_2, Z_2)$ and*

- (i) $Sq^i: \text{Ext}^{k,j} \rightarrow \text{Ext}^{i+k, 2j}$

$$\begin{aligned}
\text{(ii)} \quad Sq^i(ab) &= \sum Sq^i(a)Sq^{i-j}(b) \\
\text{(iii)} \quad Sq^a Sq^b &= \sum_{t=0}^{\lfloor a/2 \rfloor} \binom{b-t-1}{a-2t} Sq^{a+b-t} Sq^t .
\end{aligned}$$

(The properties listed in 3.1.2, 3.1.3 follow directly from the corresponding properties in the homology of the symmetric groups $\mathcal{S}_p, \mathcal{S}_{p^2}$ and the proofs are in no way different from the corresponding proofs in [25] for the ordinary Steenrod operations. The mod p case 3.1.2 appears in [12], however our grading of the mod 2 case differs from those to be found in [12], [2].)

Theorem 3.1.2 differs from the corresponding result for the ordinary Steenrod algebra in two ways. First, since there is no notion of a Bockstein homomorphism in $\text{Ext}_A(Z_p, Z_p)$ if A is a Z_p -module $\beta\mathcal{S}^i$ is not a Bockstein of \mathcal{S}^i , and, in fact is an entirely independent operation. Similarly, it is definitely not true that \mathcal{S}^0 is the identity operation (It is zero in general). Mod 2, Sq^{2i+1} is independent of Sq^{2i} . Also note that Sq^0 is neither zero nor the identity and both the Z_p and Z_2 Adem relations preserve the *number* of terms, thus two-fold iterations $Sq^a Sq^b$ or $\mathcal{S}^i \mathcal{S}^j$ can only be equal to other two-fold iterations — a considerable simplification of the topological case. Finally we point out the important special example of the Adem relations,

$$Sq^0 Sq^a = Sq^a Sq^0 .$$

2. Now our object is to imitate the construction of 3.1.1 geometrically. Thus, let

$$S^n \supset Y_n^1 \supset Y_n^2 \supset \dots \supset Y_n^r \supset \dots$$

be an Adams resolution of S^n valid for a large range of dimensions and filtrations, and suppose

$$S^{np} \supset Z_{np}^1 \supset Z_{np}^2 \supset \dots \supset Z_{np}^k \subset \dots$$

is another such, then we have

THEOREM 3.2.1. *Let $G \subset \mathcal{S}_p$ and suppose there is a G -equivariant map*

$$u: (E_G)_s \times \underbrace{S^n \wedge \dots \wedge S^n}_{p\text{-times}} \rightarrow S^{np}$$

then u is equivariantly homotopic to a map \bar{u} so

$$\bar{u}(E_G)_t \times Y_n^{i_1} \wedge \dots \wedge Y_n^{i_p} \subset Z_{np}^{i_1 + \dots + i_p - t} ,$$

moreover any two such \bar{u}, \bar{u}' are equivariantly homotopic via a homotopy H with

$$H\{I \times (E_G)_t \times Y_n^{i_1} \wedge \dots \wedge Y_n^{i_p}\} \subset Z_{np}^{i_1 + \dots + i_p - t - 1} .$$

(The proof proceeds by induction on the dimension of the skeleta of E_G , extending, first over a G -basis for the i -cells and then by G -equivariance. Similarly for the homotopy. This uses essentially the same analysis of the obstruction to compression as contained in 3.4, 3.5 of [1].)

Thus \bar{u} induces maps

$$\begin{aligned} 3.2.2. \quad ((E_G)_t, (E_G)_{t-1}) \rtimes_G (Y_n^{i_1}, Y_n^{i_1+1}) \wedge \cdots \wedge (Y_n^{i_p}, Y_n^{i_p+1}) \\ \longrightarrow (Z_{n_p}^{i_1+\cdots+i_p-t}, Z_{n_p}^{i_1+i_2+\cdots+i_p-t+1}), \end{aligned}$$

which, since $Y^i/Y^{i+1}, Z^j/Z^{j+1}$ are just one point unions of $K(Z_p, n)$'s through large ranges of dimensions, induces an $\mathcal{A}(p)$ map

$$\mathcal{C} \longrightarrow \mathcal{E}_G^* \otimes_G \mathcal{C}^{(n)}.$$

This, in turn, defines a G -equivariant $\mathcal{A}(p)$ map

$$\mathcal{E}_G \otimes \mathcal{C} \longrightarrow \mathcal{C}^{(n)},$$

and thus is suitable for defining the action of the \mathcal{P}^i in $\text{Ext}_{\mathcal{A}(p)}(Z_p, Z_p)$. In particular, on the E^1 level the map \bar{u} defines operations from 3.2.2 for each t -cell of E_G , which depend at the E^2 level only on the homology classes in $H_*(G)$ and the classes in $\text{Ext}_{\mathcal{A}(p)}(Z_p, Z_p)$ involved. Thus, these are precisely the \mathcal{P}^i operations, and their iterates.

3. Hence the existence of geometric models for the $\mathcal{P}^i, \beta\mathcal{P}^i, Sq^i$ depends on the existence of appropriate maps

$$(E_G)_s \rtimes_G S^n \wedge \cdots \wedge S^n \longrightarrow S^{nq}.$$

But these are given by 1.2.1, 1.2.2 and 1.3.3 if we start with the representations

$$r^q: \mathcal{S}_q \longrightarrow O_q,$$

defined by $r^q(\alpha)(x_1 \cdots x_q) = (x_{\alpha(1)}, \cdots, x_{\alpha(q)})$.

We have

THEOREM 3.3.1. *For a given s there is an n and an \mathcal{S}_q -map*

$$\varphi_s: (E_{\mathcal{S}_q})_s \rtimes \underbrace{S^n \wedge \cdots \wedge S^n}_{q\text{-times}} \longrightarrow S^n \wedge \cdots \wedge S^n.$$

Moreover, let $G \subset \mathcal{S}_q$ and suppose there is an equivariant map

$$\varphi: (E_G)_s \rtimes S^n \wedge \cdots \wedge S^n \longrightarrow S^n \wedge \cdots \wedge S^n$$

then given any G -map

$$\omega: E_G \longrightarrow E_{\mathcal{S}_q}$$

there is an m so $\tilde{\varphi}_s(\omega \times id)$ is equivariantly homotopic to $\tilde{\varphi}$ on $(E_G)_{s-1} \times S^{nm} \wedge \dots \wedge S^{nm}$.

(Here $\tilde{\varphi}$ is defined as the composition

$$\begin{aligned} E_{\mathcal{S}_p} \times S^{nm} \wedge \dots \wedge S^{nm} &\xrightarrow{(\mathcal{A} \times \text{shuff})} (E_{\mathcal{S}_p} \times S^n \wedge \cdot \wedge S^n)^{(m)} \xrightarrow{\varphi^{(m)}} S^{nm} \wedge \\ &\wedge \dots \wedge S^{nm} \xrightarrow{(\text{shuff})} S^{nm} \wedge \dots \wedge S^{nm} \end{aligned}$$

and similarly for $\tilde{\varphi}$. The proof is entirely similar to the proof of 1.2.2 except applied to the obstruction to extending an equivariant homotopy.

3.3.1 shows that the operations obtained in this way are stably well defined. (For further details see [22, p. 205].)

4. The \mathcal{P}^i operations and the d_2 differential in the Adams spectral sequence.

1. We pointed out in 3.1 that Sq^{2i+1} is not the Z_2 Bockstein in any sense of Sq^{2i} ; nor is $\beta\mathcal{P}^i$ the Z_p Bockstein of \mathcal{P}^i , because there is no cohomological notion of Z_p Bockstein in $\text{Ext}_{\mathcal{S}(p)}(Z_p, Z_p)$. However, there is a homotopical notion of multication by p which, in some sense, takes its place; namely, multiplication by α_0 when p is odd, and by h_0 for $p = 2$. With this in mind, we state

THEOREM 4.1.1. *Let $a \in \text{Ext}_{\mathcal{S}(p)}^{r,s}(Z_p, Z_p)$ (the E_2 term of the Adams spectral sequence); then*

$$(a) \quad d_2 Sq^i(a) = \begin{cases} h_0 Sq^{i+1}(a), & i \equiv s(2) \\ 0 & \text{otherwise} \end{cases}$$

for $p = 2$.

$$(b) \quad d_2 \mathcal{P}^i(a) = \alpha_0 \beta \mathcal{P}^i(a)$$

for p odd.

[Warning: The proof that follows appears to follow the formal algebra which would prove a similar result for a spectral sequence associated to a filtration of ordinary chain complexes. However, we are dealing with homotopy groups, so there is an error term, and the main work in the proof is to show that this error is in fact zero in the E_2 term of the Adams sequence, though it may definitely represent a nonzero element at level E_3 or E_4 .]

Proof. (a) Consider the homotopy groups $\pi_*(Y^i, Y^{i+2})$ for $S^N \supset Y_N^1 \supset \dots \supset Y_N^i \supset \dots$, a Z_2 -Adams resolution of the sphere. There is an exact sequence

$$4.1.2 \quad \xrightarrow{\partial} \pi_j(Y^{i+1}, Y^{i+2}) \xrightarrow{I} \pi_j(Y^i, Y^{i+2}) \xrightarrow{J} \pi_j(Y^i, Y^{i+1}) \xrightarrow{\partial}$$

where $\pi_s(Y^k, Y^{k+1})$ is a Z_2 vector space isomorphic to $\bar{C}_{k, k+s}$ ([1]) for

the associated resolution of Z_2 over $\mathcal{A}(2)^1$. (Of course, the ∂ map in 4.1.2 is just the d_1 differential.)

Hence $\pi_j(Y^i, Y^{i+2})$ is a sum of Z_4 's and Z_2 's, and multiplication by h_0 corresponds to multiplication by 2 here. In particular if $J(B)$ represents b in E_2 then the class $I^{-1}(2B)$ which is well defined in E_2 represents $h_0 b$.

Turning now to the case at hand suppose a is represented by a specific map

$$A: (D^{s-r}, S^{s-r-1}) \longrightarrow (Y^r, Y^{r+2})$$

and we are in the case $i \equiv s(2)$.

Thus if $v = \bar{u}(id \times A \times A): E_{Z_2} \times D^{r-s} \wedge D^{s-r} \rightarrow S$, we find that $v(e^{r-i} \times D^{s-r} \wedge D^{s-r})$ represents $Sq^i(a)$. Then

$$\begin{aligned} \partial v(e^{r-i} \times D^{s-r} \wedge D^{s-r}) &= v[(T + (-1)^{r-i})e^{r-i-1} \times D^{s-r} \wedge D^{s-r}] \\ 4.13 \quad &+ v((-1)^{r-i}e^{r-i} \times \partial D^{s-r} \wedge D^{s-r}) \\ &+ v((-1)^{s-i}e^{r-i} \times D^{s-r} \wedge \partial D^{s-r}). \end{aligned}$$

(More precisely $\partial(e^{r-i} \times D^{s-r} \wedge D^{s-r})$ is the topological union of discs which we name in this way to stay as close as possible to the standard \mathbf{U}_i formulae for cochain operations [11].) The disc $(e^{r-i} \times \partial D^{s-r} \wedge D^{s-r}) = M$ satisfies $v(M) \subset Y^{i+r+2}$, $v(\partial M) \subset Y^{i+r+3}$, and similarly $N = (e^{r-i} \times D^{s-r} \wedge \partial D^{s-r})$ satisfies $v(N) \subset Y^{i+r+2}$, $(\partial N) \subset Y^{i+r+3}$. Thus each of these discs represents a term $\{M\}, \{N\}$ in $\pi_*(Y^{r+i+1}, Y^{r+i+3})$, however $v(e^{r-i-1} \times D^{s-r} \wedge D^{s-r}) \subset Y^{r+i+1}$ while part of its boundary, the two discs $e^{r-i-2} \times D^{s-r} \wedge D^{s-r}$, $Te^{r-i-2} \times D^{s-r} \wedge D^{s-r}$, is contained in Y^{r+i+2} , and similarly for $Te^{r-i-1} \times D^{s-r} \wedge D^{s-r}$. Hence the formal decomposition of 4.1.3 is not quite valid in $\pi_*(Y^{i+r+1}, Y^{i+r+3})$.

Now, from the Blakers-Massey theorem we can regard $\{v\partial e^{r-i} \times D^{s-r} \wedge D^{s-r}\}$ as contained in $\pi_*(Y^{i+r+1}/Y^{i+r+3})$. Here the first two terms of 4.1.3 give a sphere P as does each of M, N and the boundary is $\{P\} + \{M\} + \{N\}$.

Also, note that $\{\partial v(e^{r-i+1} \times \partial D^{s-r} \wedge D^{s-r})\} = -\{M\} + \{N\}$, thus $\{P\} + \{M\} + \{N\} \sim \{P\} + 2\{N\} \sim \{P\}$ since $\{N\} \in \text{im}(I)$. It remains to prove $\{P\} \sim 2\gamma$ where γ satisfies $J(\gamma) \sim Sq^{i+1}(a)$.

Since v is equivariant there is a homeomorphism

$$\begin{aligned} h: e^{r-i-1} \times D^{s-r} \wedge D^{s-r} &\rightarrow Te^{r-i-1} \times D^{s-r} \wedge D^{s-r} \text{ so} \\ vh &= v. \text{ Since } \partial e^{r-i-1} \times D^{s-r} \wedge D^{s-r} \\ &= Te^{r-i-2} \times D^{s-r} \wedge D^{s-r} - e^{r-i-2} \times D^{s-r} \wedge D^{s-r} \end{aligned}$$

¹ At this point, we indicate our stability conventions. We put

$$\pi_j(Y^k, Y^{k+s}) = \lim_{N \rightarrow \infty} \pi_{N+j}(Y_N^k, Y_N^{k+s}).$$

Indeed, the absence of subscripts on the filtering spaces will always imply that we are considering stable groups.

we have the homotopy factorization of v on the boundary

$$\partial(\quad) \xrightarrow{\pi} S^k \vee S^k \xrightarrow{(1 \vee -1)} S^k \xrightarrow{v} S^0 \text{ where } k = 2s - r - i - 2.$$

The iteration $(+1 \vee -1) \circ \pi$ is homotopically trivial. Let $I: Q \rightarrow S$ be such a homotopy. Now we can define a class $\gamma \in \pi_*(Y^{i+r+1}/Y^{i+r+3})$ as $v(id \times I)(e^{r-i-1} \times D^{s-r} \wedge D^{s-r} \bigcup_i Q)$. Note that $J(\gamma) \sim Sq^{i+1}(a)$. A homotopy class η in $\pi_{k+1}(S^k)$ is defined as $I \bigcup_k I: Q \bigcup_{k \partial} Q \rightarrow S^k$ and we have easily

LEMMA 4.1.4. $\{P\} = 2\gamma + \eta\{v \cdot S^k\}$.

Finally, note that $\{vS^k\}$ is $(I)(\beta)$ for some β . Hence $\eta I(\beta) = I(\eta\beta)$ and part a of 4.1.1 follows from

LEMMA 4.1.5. *Let $S \supset Y^1 \supset \dots \supset Y^i \dots$ be an Adams resolution (any prime p) and $a \in \pi_*(Y^i, Y^{i+1})$ then for any element τ in $\pi_i(S)$ $i > 0$, we have $\tau a = 0$ in $\pi_*(Y^i, Y^{i+1})$.*

(Since in the Adams resolution τ has positive filtration the proof of multiplicativity in [1] shows τa is represented by an element of filtration at least $i + 1$.)

The remaining case $i \neq s(2)$ is analogous. However, here the homeomorphism h is isotopic to the identity and the resulting error term is 0. Part *b* is similar to *a*. Again by standard arguments we arrive at $d_2 \mathcal{P}^i(a)$ represented by $p\gamma + \tau\{S^k\}$ and 4.1.5 now gives the conclusion.

2. By the methods of [8] or [17], together with the results of 6.1 it is easy to prove

THEOREM 4.2.1 (a). $\text{Ext}_{\mathcal{S}_p}^{1,*}(Z_p, Z_p)$ is generated by elements α_i , h_i $i \geq 0$, $\mathcal{P}^{p^i(p-1)}(h_i) = h_{i+1}(h_i \text{ is dual to } |\mathcal{P}^{p^i}|)$ and $\alpha_0 \beta \mathcal{P}^{p^i(p-1)}(h_i) \neq 0$ in $\text{Ext}_{\mathcal{S}_p}^{3,*}(Z_p, Z_p)$.

(b) $\text{Ext}_{\mathcal{S}_2}^{1,*}(Z_2, Z_2)$ is generated by elements h_i $i \geq 0$ (h_i dual to Sq^{2^i}) $Sq^0 h_i = h_{i+1}$, and $h_0 h_i^2 \neq 0$ in $\text{Ext}_{\mathcal{S}_2}^{3,*}(Z_2, Z_2)$ for $i \geq 3$.

(4.2.1 (a) was first proved in [12], and (b) is proved in [2].)

Thus $d_2 h_i = \alpha_0 \beta \mathcal{P}^{p^i(p-1)}(h_i)$ for $i > 0$ and p odd, and the only elements which live to E_∞ in Ext^1 are α_0, h_0 . Similarly $d_2(h_i) = h_0 Sq^1 h_{i-1} = h_0 h_{i-1}^2$ and the elements h_0, h_1, h_2, h_3 are the only survivors from Ext^1 in E_∞ , for $p = 2$.

Of course these differentials are well known [16] and it is also well known that their being nonzero is equivalent to the nonexistence of elements of Hopf invariant one in these dimensions, but 4.1.1 certainly provides the first purely geometric proof of these results.


3. It sometimes happens that there is a relation such as $h_0 h_2^2 = 0$ which holds in $\text{Ext}_{\mathcal{S}_{(2)}}(Z_2, Z_2)$. The relation then “propagates”.

For example, let $\partial_1 u = h_0 h_2^2$, then $\partial_1(u \cup_1 u + h_0 h_2^2 \cup_2 u) = h_2^2 h_3^2$, $\partial_1(u \cup_2 u + h_0 h_2^2 \cup_3 u) = h_1 h_3^2$. However, some of these cup- i operations on u ultimately become parts of classes in Ext , as with $(u^2 + h_0 h_2^2 \cup_1 u) = w$ since $\partial w = h_1 h_2^4$ which equals 0 for another reason. When this occurs and we look more closely at this ∂_1 differential—say in $\pi_*(Y^i, Y^{i+2})$ rather than $\pi_*(Y^i, Y^{i+1})$ it turns out that the “filtration 2” role of the differential is to make $\partial_1(v) = \text{some element in higher filtration}$. In the example above, we have in fact

$$\partial(u \cup_1 u + h_0 h_2^2 \cup_2 u) = h_0^2 h_3^2 + 2(W + K)$$

in $\pi_*(Y^4, Y^6)$ where $K = h_1 V$, $\partial_1 V = h_2^4$ say. This and relations similar to it sometimes account for higher differentials.

Consider, for example, the situation in dimensions 14 and 15



$d_2(h_4) = h_0 h_3^2$ hence $d_2(h_0 h_4) = h_0^2 h_3^2$, but from the above remarks $h_3^2 h_3^2 = h_0 d_0$, hence $d_2 h_0 h_4 = 0$ but $d_3 h_0 h_4$ should equal $h_0 d_0$. This would even be a proof if we had a better hold on $h_3^2 h_3^2$ represented as a boundary above as compared with $h_0^2 h_3^2$ represented as 4 times h_3^2 . Such a program is possible and if carried out implies all known differential through the 45 stem in a more or less direct manner.

5. The Sq^i operation and higher differentials.

1. In [10] some theorems were proved on the connection of the Sq^i operations with d_3, d_9 under very restrictive hypothesis. Our object here is to remove most of these restrictions and increase the range of differentials considered. We have

THEOREM 5.1.1. *Let $a \in \text{Ext}_{(2)}^{t,i}(Z_2, Z_2)$ and suppose $d_s(a) = 0$ for $s < r$. Let $N = t - i + k$ for a given k and define $\varphi(N) = 8m + 2^n$*

if $N + 1 = 2^{m+n}c$, c odd, $0 \leq n < 4$, then

$$\begin{aligned} d_{\varphi(N)+1}Sq_k(a) &= h_i Sq_{k-\varphi(N)}(a), \varphi(N) = 2^i, i \leq 3 \\ d_{\varphi(N)+2}Sq_k(a) &= h_1 h_3 Sq_{k-\varphi(N)}(a), \varphi(N) = 9 \\ d_{\varphi(N)+3}Sq_k(a) &= h_1^2 h_3 Sq_{k-\varphi(N)}(a), \varphi(N) = 10 \\ d_{\varphi(N)+5}Sq_k(a) &= P^1(h_2) Sq_{k-\varphi(N)}(a), \varphi(N) = 12 \\ d_{\varphi(N)+4}Sq_k(a) &= h_0^3 h_4 Sq_{k-\varphi(N)}(a), \varphi(N) = 16 \end{aligned}$$

provided $\varphi(N) + q < r$ is satisfied in each case for the differential occurring.

In particular we mean to imply by this that all lower differentials on the specified elements are trivial. Also, note that $Sq_k(a) = Sq^{i-k}(a)$. It is probable that the theorem can be strengthened so its conclusions hold even if $\varphi(N) \leq r - q$. It is also likely that even for greater $\varphi(N)$ the differentials could still be obtained but now in terms of $d_*(a)$ as well.

2. To start the proof we “relativize” the construction of the Sq^i operations. Thus, suppose $a \in \pi_j(Y^i, Y^{i+2})$ satisfies $d_k(a) = 0$ $k < r$ then a is the restriction of an element α in $\pi_j(Y^i, Y^{i+r})$ and we can assume a represented by a map $A: (D^i, S^{j-1}) \rightarrow (Y^i, Y^{i+r})$. We have the diagram

5.2.1

$$\begin{array}{ccccc} (W_{Z_2})_t \times (D^i, S^{j-1}) \wedge (D^j, S^{j-1}) & \xrightarrow{id \times A \wedge A} & W_t \times Y^i \wedge Y^i & \xrightarrow{\hat{u}} & S \\ \downarrow id \times i \otimes i & & \swarrow & & \downarrow \\ (W_{Z_2})_t \times S^j \wedge S^j & \xrightarrow{id \times \hat{A} \wedge \hat{A}} & W_t \times Y^i / Y^{i+r} \wedge Y^i / Y^{i+r} & \xrightarrow{\hat{u}} & S / Y^{2i+r-t} \end{array}$$

where $\hat{i}: (D^j, S^{j-1}) \rightarrow (S^j, *)$ is the evident collapsing map.

The filtered map

$$\begin{array}{ccccccc} S \supset Y^1 \supset \dots \supset Y^i \supset \dots \supset Y^{2i+r-t} \\ \downarrow \quad \quad \quad \downarrow \quad \quad \quad \downarrow \\ S/Y^{2i+r-t} \supset \dots \subset Y^i/Y^{2i+r-t} \supset * \end{array}$$

induces a map of spectral sequences which is clearly a monomorphism in all E_s^t for $t < 2i + r - t$. Thus, if we can prove the existence of the stated differentials in the spectral sequence of the bottom row of 5.2.2 the theorem will follow

3. From the bottom row of 5.2.1 there is a map

$$\hat{u}: \sum^j P_j^{i+t} \longrightarrow S/Y^{2i+r-t} \text{ so that } \sum^j e^{j+k} \subset Y^{2i-k}/Y^{2i+r-t}.$$

We now define a “skeletal Adams filtration” of $\Sigma^j P_j^{i+t}$ so the map \hat{u}

becomes filtration preserving and thus induces a map of spectral sequences.

LEMMA 5.3.1. *Let X be a finite CW complex, then there is a spectral sequence converging to $\pi_*(X, Q_2)$ with*

$$\begin{aligned}\tilde{E}_2 &\cong \text{Ext}_{\mathcal{A}(2)}(Z_2, Z_2) \otimes H_*(X, Z_2), \\ (\tilde{E}_2^{r,t} &= \sum_s \text{Ext}^{r+s,t}(Z_2, Z_2) \otimes H_s(X, Z_2)).\end{aligned}$$

It is multiplicative in the sense that the pairing

$$\text{Ext}_{\mathcal{A}(2)}(Z_2, Z_2) \otimes \tilde{E}_2 \rightarrow \tilde{E}_2$$

commutes with differentials and induces pairings

$$E_r(S^0) \otimes \tilde{E}_r(X) \rightarrow \tilde{E}_r(X)$$

which also commute with differentials.

(Here Q_2 is the ring of fractions a/b with b prime to 2.)²

Proof. Suppose X is N -dimensional. We start a resolution of $\Sigma^k X$ as the fiber in the evident map

$$5.3.2 \quad \varphi_1: \Sigma^k X \rightarrow K(H^N(X), N+k).$$

Let F_1 be the fiber of φ_1 , then F_1 looks like the $N+k-1$ skeleton of $\Sigma^k X$ attached to $F'_{1, N+k}$ where F'_1 is the fiber in the map similar to 5.3.2 for a boquet of spheres. Continue the resolution by killing all the cohomology which corresponds to $F'_{1, N+k}$ and *independently* killing the cohomology from $H^{N-1}(X)$. The new fiber looks now like $\Sigma^k X_{N-2} \cup F'_{1, N+k-1} \cup F'_{2, N+k}$. Kill all the cohomology of $F'_{2, N+k}$, independently that due to $F'_{1, N+k-1}$, and that due to $H^{N-2}(X)$. The cohomology splitting again occurs so the process continues. (Note that this splitting is valid over $\mathcal{A}(2)$.) The resulting filtration of $\Sigma^k(X)$ gives the spectral sequence of 5.3.1 on taking the homotopy exact couple. We omit the remaining details as they are standard.

COROLLARY 5.3.3. *Suppose X is a finite K -connected CW complex and there is a map*

$$f: X \rightarrow S^0/Y^{2i+r-t}$$

so that $f: X_{K+1} \subset Y^{2i}$, $f: X_{K+2} \subset Y^{2i-1}$, \dots , $f: X_{K+j} \subset Y^{2i-j+1}$ then f induces a map of the spectral sequence 5.3.1 into the Adams sequence

² We must distinguish the spectral sequence introduced here from that developed in [15, Chapter III]. The bottom edge of Mahowald's sequence lies on the $s=0$ axis, while the bottom edge of ours lies on a line at a -45° angle to the $s=0$ axis. It is this change that allows us to compare our sequence with the Adams sequence.

for Y which is natural with respect to the action of the Adams sequence for a sphere on each.

(This follows directly from the proof of 5.3.1.)

4. In the situation of 5.3.3 we assume X is $\Sigma^j P_j^{i+j}$ and the map $f: X \rightarrow S^0/Y$ is \hat{u} . Clearly $\hat{u}_*(\Sigma^j e_{j+s}) = Sq^{i-s}(a)$ in the spectral sequence of 5.3.1. Thus it will suffice to prove the differentials of 5.1.1 occur in this spectral sequence. We now need

LEMMA 5.4.1. *For $L \geq N - \varphi(N)$ we have*

$$\Sigma^L P_{N-\varphi(N)+1}^N \cong S^{N+L} \vee \Sigma^L P_{N-\varphi(N)+1}^{N-1}.$$

Moreover there is a map

$$h: D^{N+L} \cup_{\alpha} S^{N+L-\varphi(N)} \longrightarrow \Sigma^L P_{N-\varphi(N)}^N$$

so h_* is a monomorphism in mod 2 homology and α is an odd multiple of the generator of $\text{im}(J)$.

(Essentially 5.4.1 says the first nontrivial attaching map of any cell in a truncated projective space is in $\text{im}(J)$.)

Proof. $P_{N-\varphi(N)+1}^N$ is the Thom complex of $(N - \varphi(N) + 1)\xi_{\varphi(N)-1}$, hence the normal bundle to $P^{\varphi(N)-1}$ is $(2^q - \varphi(N))\xi$ but

$$2^q - \varphi(N) \equiv N - \varphi(N) + 1 \pmod{2^q}$$

so

$$P_{N-\varphi(N)+1}^N = \Sigma^M(P_{2^q-\varphi(N)}^{2^q-1})$$

and, since the top class of the Thom complex for the normal bundle is spherical the first statement follows.

To obtain the remaining statement note that by Atiyah's duality theorem (3.3 of [4]) the S -dual of $P_{N-\varphi(N)}^N$ is $P_{2^q+\varphi(N)}^{2^q}$. But this the Thom complex of $2^q\xi$ over $P^{\varphi(N)}$. Now, $2^q\xi$ is trivial over $P^{\varphi(N)-1}$ and it follows from [3] that using this trivialization there is a map

$$P_{2^q+\varphi(N)}^{2^q} \longrightarrow S^{2^q} \bigcup_{\alpha} e^{2^q+\varphi(N)}$$

where α is some odd multiple of the generator of $(\text{im } J)$. Taking S -duals the result follows.

The proof of 5.1.1 follows since the $h_i, h_1h_3, P^i(h_2), h_0^3h_4$ are all known to carry the generator of $\text{im}(J)$. Indeed this shows that in 5.3.1 the boundary of the relevant cell is the term desired plus an error term from lower cells. Checking the form of Ext in these early dimensions it is clear that the only possible term besides those in 5.1.1 is

$$\varepsilon(h_3^2 Sq_{k-\varphi(N)-3}(a)) \text{ for all } \varphi(N) = 12.$$

However, by checking secondary operations we can show this attach-

ing map is 0 so $\varepsilon = 0$ and the result follows.

5. It is easy to see how to generalize this to the mod p -case. In view of the results of § 2.3 it would seem that the hardest step in doing this is to calculate sufficiently far into $\text{Ext}_{\mathcal{A}(p)}(Z_p, Z_p)$ and still identify $\text{im}(J)$.

REMARK 5.5.1. Recent results of Mahowald [14] enable one to completely identify $\text{im}(J)$ in $\text{Ext}_{\mathcal{A}(2)}(Z_2, Z_2)$. It seems reasonable to suppose the mod(p) problem can be handled in a similar way.

6. Calculation of the action of the Steenrod Algebra in $\text{Ext}_{\mathcal{A}(p)}(Z_p, Z_p)$.

1. Some examples. We consider, mod p , the algebra $A = DP(\lambda)$ where $DP(\lambda)$ is a divided polynomial algebra. It is given a Hopf algebra structure by requiring that $\Delta(\lambda_i) = \sum \lambda_j \otimes \lambda_{i-j}$. Then for p an odd prime

$$\text{Ext}_A^{**}(Z_p, Z_p) = E(|\lambda|, |\lambda_p|, \dots, |\lambda_{p^i}|, \dots) \otimes P(\theta_1, \theta_p, \dots, \theta_{p^i} \dots)$$

where θ_{p^i} is dual to $|\lambda_{p^i}| \lambda_{p^i}^{p-1}|$.

We calculate the \mathcal{S}^i operations as

$$\begin{aligned} \text{THEOREM 6.1.1. } \mathcal{S}^{p^i(p-1)} |\lambda_{p^i}| &= c_1 |\lambda_{p^{i+1}}| \\ \beta \mathcal{S}^{p^i(p-1)} |\lambda_{p^i}| &= c_2 \theta_{p^i} \\ \mathcal{S}^{p^i(p-1)+1} |\lambda_{p^i}| &= 0 \\ \mathcal{S}^{(p-1)p^i+1+1} \theta &= (\theta_{p^i})^p \end{aligned}$$

where c_1, c_2 are nonzero constants, and this completely determines the action of $\tilde{\mathcal{A}}(p)$ here.

Proof. Consider in $\Delta^{(p)} |\lambda_{p^i}| \lambda_{p^i}^{p-1}|$ the term $|\lambda_{p^i}| \otimes |\lambda_{p^i}| \otimes \lambda_{p^i} \otimes \dots \otimes \lambda_{p^i}$. Clearly as we go through the successive constructions of the higher homotopies M^i which give the \mathcal{S}^i operations, we arrive at the term $|\lambda_{p^i}| \otimes \dots \otimes |\lambda_{p^i}|$ in $M^{p-2}(\bar{\theta}_{p^i})$. This proves the second statement, the first is virtually identical.


Similarly, for the same algebra mod 2 we have

$$\text{Ext}_A(Z_2, Z_2) = P(|\lambda|, |\lambda_2| \dots |\lambda_{2^i}| \dots)$$

and for the Sq^i operations we find

$$\begin{aligned} \text{THEOREM 6.1.2 } Sq^1 |\lambda_{2^i}| &= |\lambda_{2^i}|^2 \\ Sq^0 |\lambda_{2^i}| &= |\lambda_{2^{i+1}}| \end{aligned}$$

and these determine the action of $\tilde{\mathcal{A}}(2)$ in $\text{Ext}_A(Z_2, Z_2)$.
(The proof is unchanged from that of 6.1.1)



Massey product representations

$$\begin{aligned}
 c_0 &= \langle h_{11}, h_{10}, h_{12}^2 \rangle \\
 c_1 &= \langle h_{21}, h_{11}, h_{13}^2 \rangle \\
 c_2 &= \langle h_{31}, h_{21}, h_{14}^2 \rangle \\
 d_0 &= \langle h_{01}, h_{22}, h_{01}, h_{12}^2 \rangle \\
 d_1 &= \langle h_{11}, h_{22}, h_{11}, h_{13}^2 \rangle \\
 p &= \langle h_{01}, h_{22}, h_{14}^2 \rangle \\
 x &= \langle h_{31}, h_{11}, d_0 \rangle \\
 f_0 &= \langle h_{01}, h_{22}, h_{12}^2 \rangle \\
 f_1 &= \langle h_{11}, h_{22}, h_{13}^2 \rangle
 \end{aligned}$$

(The families $Sq^i(\omega)$ for ω fixed go along a line of slope -45° , with highest point ω^2 .)

FIGURE 1 (Continued)

Clearly $\omega(|\lambda_2|) = h_i$ in $\text{Ext}_{\mathcal{A}(2)}^{1,2^i}(Z_2, Z_2)$. Since ω is induced from a map of Hopf algebras $\omega(Sq^i) = Sq^i(\omega)$ and we can calculate the action of $\mathcal{A}(2)$ in the subalgebra of $\text{Ext}_{\mathcal{A}(2)}(Z_2, Z_2)$ generated by the (h_i) .

The first generator which is not a polynomial in the h_i is c_0 which has the Massey product representation $\langle h_1, h_0, h_2^2 \rangle$. (For further details and the “names” of various elements we refer to Figure 1.)

$$\begin{aligned}
\text{PROPOSITION 6.2.1. } Sq^3(c_0) &= c_0^3 = h_1^3 d_0 \\
Sq^2(c_0) &= h_0 e_0 \\
Sq^1(c_0) &= f_0 \\
Sq^0(c_0) &= c_1
\end{aligned}$$

(The first statement and the fourth are known. The first following from [17], and the fourth from [19]. The second and third are shown using the techniques of [19] rather than the main theorem and explicitly writing representatives for these elements.) Thus, in view of 4.1.1 we find $d_2(f_0) = h_0^2 e_0$, $d_2(h_0 f_0) = d_2(h_1 e_0) = h_0^3 e_0 = h_1^3 d_0$ so $d_2 e_0 = h_1^2 d_0$, and it is only the differential $d_3(h_0 h_4) = h_0 d_0$ which is not due to 4.1.1, at least through the 21 stem.

3. Now we consider the algebra A^2 more closely. $(A^2)^* = P(\xi_2, \xi_3 \cdots \xi_i \cdots)$ with $\varphi(\xi_i) = \sum_{2 \leq j \leq i-1} \xi_j^{2^{i-j}} \otimes \xi_{i-j}$. In particular (ξ_2, ξ_3) generate a sub-Hopf algebra B of $(A^2)^*$ and, if $A^{(4)}$ is the dual of B in A^2 , we have

$$A^2/A^4 = DP(\lambda_2) \otimes DP(\lambda_3) .$$

Consequently there is a map

$$u: \text{Ext}_{DP(\lambda_2) \otimes DP(\lambda_3)}^s(Z_2, Z_2) \longrightarrow \text{Ext}_{A^2}^{s,t}(Z_2, Z_2) .$$

PROPOSITION 6.3.1. $\text{Ext}_{DP(\lambda_2) \otimes DP(\lambda_3)}(Z_2, Z_2) = \text{Ext}_{DP(\lambda_2)} \otimes \text{Ext}_{DP(\lambda_3)}$ as algebras over $\mathscr{A}(2)$. Moreover for $t - s$ less than 50 u is onto and has kernel generated by

$$\lambda_{2,2^i} \cdot \lambda_{2,2^{i+2}}, \lambda_{2,2^{i+3}} \cdot \lambda_{3,2^i} + \lambda_{3,2^{i+2}} \cdot \lambda_{2,2^i}, \lambda_{2,2^i}^2 \cdot \lambda_{2,2^{3+i}} + \lambda_{2,2^{i+1}} \cdot \lambda_{2,2^{i+2}}^2$$

and

$$\lambda_{2,1}^4 \cdot \lambda_{2,16} + \lambda_{2,4}^5 .$$

Proof. The first part is evident. To prove the second we use the techniques of [8] and read off the differentials in the weighted augmentation filtration spectral sequence developed there.

This determines the action of $\mathscr{A}(2)$ in Ext_{A^2} and in this range. Consider finally the map $v: A^2 \rightarrow \mathscr{A}(2)$, this induces $v^*: \text{Ext}_{\mathscr{A}(2)} \rightarrow \text{Ext}_{A^2}$ and we have, again making use of [8]

$$\begin{aligned}
\text{PROPOSITION 6.3.2. } v^*(d_0) &= \{(\lambda_{22}\lambda_{22})^2\} \\
v^*(e_0) &= \{\lambda_{21}(\lambda_{22})^3\} \\
v^*(g) &= \{\lambda_{22}^4\} \\
v^*P^1(d_0) &= \{\lambda_{21}^6 \lambda_{22}^2\} \\
v^*P^1(e_0) &= \{\lambda_{21}^5 \lambda_{22}^3\} \\
v^*(d_1) &= \{\lambda_{22}\lambda_{24}\}^2 .
\end{aligned}$$

$$v^*(k) = \{\lambda_{21}^2 \lambda_{22}^5\}, v^*(n) = \{\lambda_{21} \lambda_{22}^4\}$$

4. From 6.3.2 we obtain $Sq^0(d_0) = d_1$ (this is clear anyway from [19]) $Sq^0(e_0) = e_1$, $Sq^2(d_0) = r$ since $v^*Sq^2(d_0) = \lambda_{22}^5 \neq 0$ and the only element in $t - s = 30$ and filtration 6 in $\text{Ext}_{\mathcal{A}(2)}$ is r . Similarly $Sq^3(e_0) = m$ since $v^*Sq^3(e_0) = \lambda_{22}^7 \neq 0$ and m is the only candidate. Also $Sq^2(e_0) = t$ since $Sq^2v^*(e_0) = \lambda_{22}^5 \lambda_{24} \neq 0$ and t is the only candidate. Putting these together we have

$$\begin{aligned} \text{THEOREM 6.4.1. } Sq^3(d_0) &= Sq^1(d_0) = 0 \\ Sq^4(d_0) &= d_0^2, Sq^2(d_0) = r, Sq^0(d_0) = d_1 \\ Sq^4(e_0) &= e_0^2, Sq^3(e_0) = m, Sq^2(e_0) = t, Sq^1(e_0) = x \\ Sq^0(e_0) &= e_1. \end{aligned}$$

Proof. We first show $Sq^1(e_0) = x$. From ([13, Proposition 5.1.3]) we have $h_2e_0^2 = h_0^4x$. But from

$$Sq^4(h_1e_0) = h_2e_0^2 = Sq^4(h_0f_0) = h_0^2Sq^2(f_0)$$

it follows that

$$6.4.2. \quad Sq^3(f_0) = \varepsilon(h_1t) + h_0^2x.$$

Also

$$Sq^3f_0 = Sq^3Sq^1c_0 = Sq^2Sq^2c_0 = Sq^2(h_0e_0)$$

by the Adém relation $Sq^2Sq^2 = Sq^3Sq^1$. But

$$6.4.3. \quad Sq^2(h_0e_0) = h_0^3Sq^1(e_0) + h_1t.$$

Now the result follows by comparing 6.4.2 with 6.4.3.

To verify $Sq^1(d_0) = 0$ note that the only candidate is n but $v^*(n) \neq 0$ while $Sq^1v^*(d_0) = 0$. Again, the only candidate for $Sq^3(d_0)$ is k but $v^*(k) \neq 0$ while $Sq^3(v^*(d_0)) = 0$. This completes the proof of 6.4.1.

$$\text{COROLLARY 6.4.4. } d_2m = h_0e_0^2.$$

$$\text{COROLLARY 6.4.5. } h_0r = s, h_3r = h_1t + h_0^2x.$$

Proof. From 6.4.2, 6.4.3, 6.4.1 $Sq^2(h_2d_0) = h_3r = h_0^2Sq^1(e_0) + h_1t$. Now, since $h_0h_3r \neq 0$ it must follow that $h_0r = s$. (This result is asserted and an outline proof given in [13, 7.4], but the proof given here is much more direct.)

5. We now have

$$\text{THEOREM 6.5.1. } Sq^0(c_1) = c_2, Sq^1(c_1) = f_1, Sq^2(c_1) = h_1e_1, Sq^3c_1 = h_2d_1, \\ Sq^0(f_0) = f_1, Sq^1f_0 = 0, Sq^2(f_0) = y, Sq^3(f_0) = h_1t + h_0^2x.$$

Proof. $Sq^0(c_1) = c_2$, $Sq^0(f_0) = f_1$ follow from [19], so $Sq^1(c_1) = f_1$ since $Sq^0Sq^1 = Sq^1Sq^0$. Now note that $h_1e_0^2 = h_0e_0f_0 = h_0^2m$. Thus $e_0f_0 = h_0m$ and $h_2m = h_0^2y \neq 0$ since $h_2e_0f_0 = h_1e_0g$. Hence $Sq^3(h_0f_0) = h_0^2Sq^2(f_0) + h_1Sq^3(f_0) = h_2m + h_1^2t = h_2m$ since $h_1^2t = 0$. We have previously shown $Sq^3(f_0)$. $Sq^1(f_0) = Sq^1Sq^1(c_0) = 0$ by the Adém relation. Similarly $Sq^0Sq^2(c_0) = h_1e_1 = Sq^2(Sq^0c_0) = Sq^2(c_1)$.

This essentially completes the description of the Sq^i operations in the range $t - s < 45$. In particular we have

COROLLARY 6.5.2. $d_2(c_2) = h_0f_1$

$$d_2(y) = h_0^3x.$$

This first differential was overlooked in [13] and to obtain the second differential they had to work considerably harder. The first differential also implies $h_2h_4^2$ is nonzero as a homotopy element [$\nu\theta_4 \neq 0$] since c_2 has Massey product representation $\langle h_3, h_2, h_4^2 \rangle$ which by [24] would converge to the Toda bracket $\{\sigma, \nu, \theta_4\}$ if this bracket were defined. This in turn can be shown to imply another differential overlooked in [13], namely $d_3(h_2h_5) = h_0p$. I should point out here that Barratt, Mahowald and Tangora independently discovered this error by investigating an apparent inconsistency in the 47 stem, and tracing it back [5]. The error was also found at about the same time by G.W. Whitehead using the semi-simplicial techniques of Kan-Curtis.

7. The use of the symmetric groups in homotopy theory.

1. So far the constructions $E_{Z_2} \rtimes X \wedge \cdots \wedge X$, respectively $E_{Z_2} \rtimes X \wedge X$, have been used by Toda [13], respectively Adams, Barratt and Mahowald, to obtain much information on homotopy groups. Basically they use the retractions supplied by § 3 or other maps implied by the E.H.P. sequence. However it is clear from §§ 1 and 2 that we can also use $E_{\mathcal{S}_n} \rtimes \underbrace{X \wedge \cdots \wedge X}_{n\text{-times}}$ to obtain information and it seems re-

asonable to assume that much remains to be done in this direction.

2. By way of an example we show how to use \mathcal{S}_4 to obtain a particularly simple proof that θ_4 exists and is nonzero.

Consider $\sigma: S^7 \rightarrow S^0$. This induces a map

$$\hat{u}(\sigma): E_{\mathcal{S}_4} \rtimes_{\mathcal{S}_4} S^7 \wedge \cdots \wedge S^7 \rightarrow E_{\mathcal{S}_4} \rtimes S^0 \wedge \cdots \wedge S^0 \xrightarrow{i} S^0.$$

Now by the results of § 2.4 we have $H^*(E_{\mathcal{S}_4} \rtimes_{\mathcal{S}_4} S^7 \wedge \cdots \wedge S^7)$ is $H^*(\mathcal{S}_4) \cup U$ and $Sq^1U = CU$, $Sq^2U = (B + C^2)U$. Consider now the map $Z_2 \rightarrow \mathcal{S}_4$ by $T \rightarrow (13)(24)$. This induces a map $f: E_{\mathcal{S}_2} \rtimes_{\mathcal{S}_2} S^{14} \wedge S^{14} \rightarrow E_{\mathcal{S}_4} \rtimes_{\mathcal{S}_4} S^7 \wedge \cdots \wedge S^7$ and it is easy to check $f^*(BU) = e^2U'$. Also

there is a map $Z_2 \oplus Z_2 \rightarrow \mathcal{S}_4(T_1 \rightarrow (12), T_2 \rightarrow (34))$ which induces a map

$$g: E_{Z_2 \oplus Z_2} \times_{Z_2 \oplus Z_2} S^7 \wedge \cdots \wedge S^7 \rightarrow E_{\mathcal{S}_4} \times S^7 \wedge \cdots \wedge S^7.$$

$$g^*(C) = (e \otimes 1 + 1 \otimes e)U' \quad g^*(B) = e \otimes eU'.$$

This last shows C^2 is dual to $(h_3 \cup_2 h_3)h_3^2$ while B is dual to h_4^2 . $h_3 \cup_2 h_3$ has filtration at least 2 so $(h_3 \cup_2 h_3)h_3^2$ has filtration 4 or more. Since $Sq^2 U = (B + C^2)U$ it follows that both these discs h_4^2 , $(h_3 \cup_2 h_3)h_3^2$ attach to U by η . Hence their difference represents a sphere which must be called h_4^2 in the Adams spectral sequence.

REFERENCES

1. J. F. Adams, *On the structure and applications of the Steenrod algebra*, Comm. Math. Helv., **32** (1958), 180-214.
2. ———, *On the non-existence of elements of Hopf invariant one*, Ann. of Math. **72** (1960), 20-104.
3. ———, *Vector fields on spheres*, Ann. of Math. **75** (1962), 603-632.
4. M. F. Atiyah, *Thom complexes*, Proc. London Math. Soc., **11** (1961), 291-310.
5. M. Barratt, M. Mahowald and M. Tangora, *Some differentials in the Adams spectral sequence II*, Topology **9** (1970), 309-317.
6. H. Cardenas, *El algebra de cohomologia del grupo simétrico de grado p^2* , Bol. Soc. Mat. Mexicana, **10** (1965), 1-30.
7. W. Feit, *Characters of Finite Groups*, Benjamin (1967).
8. V. K. A. M. Gugenheim and R. J. Milgram, *On successive approximations in homological algebra*, Trans. Amer. Math. Soc., **150** (1970), 157-182.
9. D. S. Kahn, *A differential in the Adams spectral sequence*, Proc. Amer. Math. Soc., **20** (1961), 188-191.
10. ———, *Cup-i products and the Adams spectral sequence*, Topology, **9** (1970), 1-10.
11. L. Kristensen, *On the cohomology of two stage Postnikov systems*, Acta Math., **107** (1962), 73-123.
12. A. Liulevicius, *The factorization of cyclic reduced powers by secondary cohomology operations*, Mem. Amer. Math. Soc., **42** (1962).
13. M. Mahowald and M. Tangora, *Some differentials in the Adams spectral sequence*, Topology **6**(1967), 349-369.
14. M. Mahowald, *The order of the image of the J-homomorphism*, Bull. Amer. Math. Soc., **76** (1970), 1310-1314.
15. ———, *The metastable homotopy of S^n* , Mem. Amer. Math. Soc., **72** (1967).
16. C. R. Maunder, *Some differentials in the Adams spectral sequence*, Proc. Cambridge Philos. Soc., **60**(1964), 409-420.
17. J. P. May, Ph. D. thesis, Princeton University, 1964.
18. R. J. Milgram, *The bar construction and abelian H-spaces*, Illinois J. Math., **11** (1967), 242-250.
19. ———, *Steenrod squares and higher Massey products*, Bol. Soc. Mat. Mexicana, (1968), 32-51.
20. ———, *The mod 2 spherical characteristic classes*, Ann. of Math., **92** (1970), 238-261.
21. ———, *A construction for s-parallelizable manifolds and primary homotopy operations*, Topology of Manifolds, Markham (1970), 483-489.
22. ———, *Symmetries and operations in homotopy theory*, Amer. Math. Soc. Proc. Symposia Pure Math., **22** (1971), 203-211.
23. J. Milnor, *The Steenrod algebra and its dual*, Ann. of Math., **67** (1958), 150-171.

- 24. M. Moss, *Secondary compositions and the Adams spectral sequence*, Topology, **10** (1971).
- 25. N. Steenrod and D. Epstein, *Cohomology operations*, Princeton University Press, Ann. of Math. Study **51** (1962).
- 26. H. Toda, *Extended pth powers of complexes and applications homotopy theory*, Proc. Japan Acad., **44**(1968), 198-203.

Received October 19, 1970.

UNIVERSITY OF ILLINOIS AT CHICAGO CIRCLE
AND
STANFORD UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 41, No. 1

November, 1972

Anatole Beck and Peter Warren, <i>Weak orthogonality</i>	1
Jonnie Bee Bednar and Howard E. Lacey, <i>Concerning Banach spaces whose duals are abstract L-spaces</i>	13
Louis Harvey Blake, <i>Canonical extensions of measures and the extension of regularity of conditional probabilities</i>	25
R. A. Brooks, <i>Conditional expectations associated with stochastic processes</i>	33
Theodore Allen Burton and Ronald Calvin Grimmer, <i>On the asymptotic behavior of solutions of $x'' + a(t)f(x) = e(t)$</i>	43
Stephen LaVern Campbell, <i>Operator-valued inner functions analytic on the closed disc</i>	57
Yuen-Kwok Chan, <i>A constructive study of measure theory</i>	63
Alexander Munro Davie and Bernt Karsten Oksendal, <i>Peak interpolation sets for some algebras of analytic functions</i>	81
H. P. Dikshit, <i>Absolute total-effective $(N, p_n)(c, 1)$ method</i>	89
Robert E. Edwards, Edwin Hewitt and Kenneth Allen Ross, <i>Lacunarity for compact groups. II</i>	99
James Daniel Halpern, <i>On a question of Tarski and a maximal theorem of Kurepa</i>	111
Gerald L. Itzkowitz, <i>A characterization of a class of uniform spaces that admit an invariant integral</i>	123
Mo Tak Kiang, <i>Semigroups with diminishing orbital diameters</i>	143
Glenn Richard Luecke, <i>A class of operators on Hilbert space</i>	153
R. James Milgram, <i>Group representations and the Adams spectral sequence</i>	157
G. S. Monk, <i>On the endomorphism ring of an abelian p-group, and of a large subgroup</i>	183
Yasutoshi Nomura, <i>Homology of a group extension</i>	195
R. Michael Range, <i>Approximation to bounded holomorphic functions on strictly pseudoconvex domains</i>	203
Norman R. Reilly, <i>Inverse semigroups of partial transformations and θ-classes</i>	215
Chris Rorres, <i>Strong concentration of the spectra of self-adjoint operators</i>	237
Saharon Shelah, <i>A combinatorial problem; stability and order for models and theories in infinitary languages</i>	247
George Gustave Weill, <i>Vector space decompositions and the abstract imitation problem</i>	263
Arthur Thomas White, <i>On the genus of the composition of two graphs</i>	275