

Pacific Journal of Mathematics

ON THE GENUS OF THE COMPOSITION OF TWO GRAPHS

ARTHUR THOMAS WHITE

ON THE GENUS OF THE COMPOSITION OF TWO GRAPHS

ARTHUR T. WHITE

Given two graphs G and H , a new graph $G(H)$, called the composition (or lexicographic product) of G and H , can be formed. In this paper, a formula is developed to give the genus for a large class of lexicographic products. In the simplest special case, the genus of the product is given by the first Betti number of one of the factors.

In the present context, a *graph* is a finite 0- or 1- complex. For terms not defined below, see [2] and [6].

The *genus*, $\gamma(G)$, of a graph G is the minimum genus among the genera of all closed orientable 2-manifolds M in which G can be imbedded. An imbedding of G in M is said to be *minimal* if M has genus $\gamma(G)$. The *first Betti number*, $\beta(G)$, of a graph G is given by $\beta(G) = q - p + k$, where G has q edges, p vertices, and k components; $\beta(G)$ counts the number of independent cycles in G . Given two graphs G and H with disjoint vertex sets $V(G)$, $V(H)$ and edge sets $E(G)$, $E(H)$ respectively, the *composition* (or *lexicographic product*) $G(H)$ has vertex set given by the cartesian product $V(G) \times V(H)$, with two vertices (u_i, v_j) and (u_k, v_m) adjacent in $G(H)$ if and only if either: (i) $u_i = u_k$ and $v_j v_m$ is in $E(H)$, or (ii) $u_i u_k$ is in $E(G)$. For example, the regular complete m -partite graph on mn vertices is just the composition $K_m(\overline{K_n})$, where K_s denotes the complete graph on s vertices, and $\overline{K_s}$ denotes the complement of K_s (a 0-complex).

We will also employ the following notions. If G is imbedded in M , the components of $M - G$ are called *regions*. A region bounded by a circuit of length 3(4) in G is said to be *triangular* (*quadrilateral*). The number of triangular (quadrilateral) regions in a given imbedding is denoted by $r_3(r_4)$. In general, r_k designates the number of regions having a connected boundary consisting of k edges of G , and r denotes the total number of regions. It is well known (see, for example, [6]) that, for a minimal imbedding of a connected graph G having p vertices and q edges, the Euler formula $p - q + r = 2 - 2\gamma(G)$ applies. Also, it is easy to show that $2q = \sum_{i \geq 3} i r_i$. We note that a 3-cycle in a graph G need not bound a triangular region in a given minimal imbedding of G . For example, there are 35 3-cycles in K_7 ; yet any minimal imbedding of K_7 has $r = r_3 = 14$.

The following result of Battle, Harary, Kodama, and Youngs [1] will be useful:

THEOREM. *The genus of a graph is the sum of the genera of its components.*

We are now prepared to state the main result.

THEOREM. *Let G have p vertices of positive degree, q edges, k nontrivial components, and no 3-cycles. Let H have $2n(n \geq 1)$ vertices and maximum degree less than two. Then $\gamma(G(H)) = k + n(nq - p)$.*

Proof. Let G have nontrivial components $C_i, i = 1, \dots, k$; then $G(H)$ has nontrivial components $C_i(H), i = 1, \dots, k$. It will suffice to prove the theorem for G connected, since then (by the result of Battle, Harary, Kodama and Youngs):

$$\begin{aligned} \gamma(G(H)) &= \sum_{i=1}^k \gamma(C_i(H)) \\ &= \sum_{i=1}^k (1 + n(nq_i - p_i)) \\ &= k + n(nq - p) . \end{aligned}$$

We therefore assume G to be connected. Let $V(G) = \{u_1, \dots, u_p\}$, and $V(H) = \{v_1, \dots, v_{2n}\}$.

Suppose the vertices $(u_i, v_j), (u_k, v_m)$, and (u_r, v_s) form a 3-cycle in $G(H)$. Since there are no 3-cycles in G , the vertices u_i, u_k , and u_r cannot be distinct in $V(G)$. Hence every 3-cycle in $G(H)$ must contain an edge of the form $(u_i, v_j)(u_i, v_m)$. There are exactly pe such edges in $G(H)$, where e designates the number of edges in $H(0 \leq e \leq n)$. Since each one of these edges can appear in the boundary of at most 2 triangular regions, it follows that $r_3^* \leq 2pe$ in any imbedding of $G^* = G(H)$. (A parameter with (without) an asterisk will apply to graph $G^*(G)$).

We will construct an imbedding of G^* so that $r_3^* = 2pe$ and $r_4^* = r^* - 2pe$; since $r^* = \sum_{i \geq 3} r_i^*$ and $2q^* = \sum_{i \geq 3} i r_i^*$, r^* will be maximal for such an imbedding. Then, by the Euler formula, the imbedding itself will be minimal. Now, for G^* , $p^* = 2np$, and $q^* = pe + 4n^2q$. Also, if $r_3^* = 2pe = r^* - r_4^*$, then $r^* = pe + 2n^2q$, since $2q^* = 2pe + 8n^2q = 3(2pe) + 4(r^* - 2pe)$. Then, from the Euler formula,

$$\begin{aligned} \gamma(G^*) &= 1 + 1/2(q^* - p^* - r^*) \\ &= 1 + 1/2(pe + 4n^2q - 2np - (pe + 2n^2q)) \\ &= 1 + n(nq - p) . \end{aligned}$$

We now construct such an imbedding. Let the edges of G be designated by x_1, \dots, x_p . For each edge there is a subgraph of $G(H)$ isomorphic to the complete bipartite graph $K_{2n, 2n}$. Imbed q copies of $K_{2n, 2n}$, minimally, in q closed orientable 2-manifolds M_1, \dots, M_q of

genus $(n - 1)^2$ each, in the fashion described by Ringel [3]. Select these 2-manifolds so that each is exterior to any other. Each imbedding has $r' = r'_i = 2n^2$, and it has been shown in [5] that the $2n^2$ quadrilateral regions can be partitioned into $2n$ mutually disjoint sets of n regions each, each set containing all $4n$ vertices of the graph $K_{2n, 2n}$. Furthermore, in any region, diagonally opposite vertices are in the same part of the vertex set partition for $K_{2n, 2n}$.

Suppose edges x_i and x_j are adjacent in G . We make $2n$ vertex identifications between M_i and M_j as follows. Select one set of n quadrilateral regions in M_i and the $2n$ vertices of one part of the vertex set partition for $K_{2n, 2n}$ from the boundaries of these regions (two diagonally opposite vertices are selected from the boundary of each region). Make similar selections in M_j . Now attach n tubes between M_i and M_j , one tube for each pair of regions (one from each 2-manifold) that we have selected. The first such tube may be attached as follows. Let region R^i in M_i correspond to region R^j in M_j . Let C^h be a simple closed curve bounding the open disk D^h interior to R^h , $h = i, j$. Let T be a topological cylinder, with bases C_i and C_j , such that $(M_i \cup M_j) \cap T = C_i \cup C_j$. Form $(M_i - D^i) \cup (M_j - D^j) \cup T$. It is clear how to add the remaining tubes. The result is a closed orientable 2-manifold M (of genus $2(n - 1)^2 + n - 1$).

We now make two vertex identifications per tube, as indicated by the sequence of operations in Figure 1.




FIGURE 1

This process destroys two quadrilaterals and creates two new quadrilaterals for each tube. Furthermore, the two identifications for each tube yield two vertices diagonally opposite in a common region R . If edge x_k is also adjacent to x_i (and to x_j) in G , there are now n regions available on the 2-manifold M with which to make the appropriate $2n$ identifications with the 2-manifold M_k . From these n regions on M , we select the diagonally opposite vertices that resulted from the first identification. It is clear that this process may be continued until a quadrilateral imbedding of $G(\overline{K}_{2n})$ results. We need only insure that, for 2-manifold M_i corresponding to edge $x_i = [u_{i1}, u_{i2}]$ in G , if we selected the $2n$ vertices of one part of the vertex-set

partition of $K_{2n,2n}$ with which to make the identifications at u_{i_1} in G , then we must select the $2n$ vertices of the second part of the vertex-set partition of $K_{2n,2n}$ with which to make the identifications at u_{i_2} in G .

Corresponding to each vertex of G , there is now a copy of \overline{K}_{2n} , within which the e edges of H may be added. Each such edge converts one quadrilateral region of the imbedding of $G(\overline{K}_{2n})$ into two triangular regions. The result is an imbedding of G^* having $r_3^* = 2pe$ and $r_4^* = r^* - 2pe$, as desired. This completes the proof.

We note that the value $r^* = 2qn^2 + pe$ may be verified by a direct count, since $r_4^* = q(2n^2) - pe$. Also, the genus of $G(H)$ may be computed directly, for this construction, without recourse to any Euler type formula. The contributions to the genus are of three types:

- (i) $q(n - 1)^2$, representing the collective genera of the q 2-manifolds with which we began our construction;
- (ii) $(2q - p)(n - 1)$, representing the contribution of the $2q - p$ sets of $2n$ vertex identifications each, each "bundle" of n tubes adding $n - 1$ to the genus;
- (iii) $\beta(G) = q - p + 1$, representing the contribution of the bundles of tubes taken collectively.

Adding, we find:

$$\begin{aligned} \gamma(G(H)) &= q(n - 1)^2 + (2q - p)(n - 1) + (q - p + 1) \\ &= 1 + n(nq - p) . \end{aligned}$$

It is no surprise that the formula $\gamma(K_{2n,2n}) = (n - 1)^2$ is included in the above theorem. For the case where G is the cycle C_s and $H = \overline{K}_{2n}$, we may combine the theorem with the result of Ringel and Youngs [4] that $\gamma(K_3(\overline{K}_m)) = ((m - 1)(m - 2))/2$ to establish the following:

COROLLARY 1.

$$\gamma(C_s(\overline{K}_{2n})) = \begin{cases} 1 + n(2n - 3), & \text{if } s = 3 \\ 1 + ns(n - 1), & \text{if } s \geq 4 . \end{cases}$$

In the situation where G is the complete bipartite graph $K_{r,s}$ and H is as in the statement of the theorem, we have:

COROLLARY 2. $\gamma(K_{r,s}(H)) = (nr - 1)(ns - 1)$.

We list here only one other result, for the special case $n = 1$ of the theorem:

COROLLARY 3. *Let G be a graph containing no 3-cycles. Then $\gamma(G(K_2)) = \gamma(G(\overline{K}_2)) = \beta(G)$.*

REFERENCES

1. J. Battle, F. Harary, Y. Kodama, and J. W. T. Youngs, *Additivity of the genus of a graph*, Bull. Amer. Math. Soc., **68** (1962), 565-568.
2. F. Harary, *Graph Theory*, Addison-Wesley, Reading (1969).
3. G. Ringel, *Das Geschlecht des Vollständigen Paaren Graphen*, Abh. Math. Sem. Univ. Hamburg, **28** (1965), 139-150.
4. G. Ringel and J. W. T. Youngs, *Das Geschlecht des Symmetrische Vollständige Drei-Farbaren Graphen*. Comment. Math. Helv., **45** (1970), 152-158.
5. A. T. White, *The genus of repeated cartesian products of bipartite graphs*. Trans. Amer. Math. Soc., **151** (1970), 393-404.
6. J. W. T. Youngs, *Minimal imbeddings and the genus of a graph*, J. Math. Mech., **12** (1963), 303-315.

Received May 22, 1970

WESTERN MICHIGAN UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

C. R. HOBBY
University of Washington
Seattle, Washington 98105

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Anatole Beck and Peter Warren, <i>Weak orthogonality</i>	1
Jonnie Bee Bednar and Howard E. Lacey, <i>Concerning Banach spaces whose duals are abstract L-spaces</i>	13
Louis Harvey Blake, <i>Canonical extensions of measures and the extension of regularity of conditional probabilities</i>	25
R. A. Brooks, <i>Conditional expectations associated with stochastic processes</i>	33
Theodore Allen Burton and Ronald Calvin Grimmer, <i>On the asymptotic behavior of solutions of $x'' + a(t)f(x) = e(t)$</i>	43
Stephen LaVern Campbell, <i>Operator-valued inner functions analytic on the closed disc</i>	57
Yuen-Kwok Chan, <i>A constructive study of measure theory</i>	63
Alexander Munro Davie and Bernt Karsten Oksendal, <i>Peak interpolation sets for some algebras of analytic functions</i>	81
H. P. Dikshit, <i>Absolute total-effective $(N, p_n)(c, 1)$ method</i>	89
Robert E. Edwards, Edwin Hewitt and Kenneth Allen Ross, <i>Lacunarity for compact groups. II</i>	99
James Daniel Halpern, <i>On a question of Tarski and a maximal theorem of Kurepa</i>	111
Gerald L. Itzkowitz, <i>A characterization of a class of uniform spaces that admit an invariant integral</i>	123
Mo Tak Kiang, <i>Semigroups with diminishing orbital diameters</i>	143
Glenn Richard Luecke, <i>A class of operators on Hilbert space</i>	153
R. James Milgram, <i>Group representations and the Adams spectral sequence</i>	157
G. S. Monk, <i>On the endomorphism ring of an abelian p-group, and of a large subgroup</i>	183
Yasutoshi Nomura, <i>Homology of a group extension</i>	195
R. Michael Range, <i>Approximation to bounded holomorphic functions on strictly pseudoconvex domains</i>	203
Norman R. Reilly, <i>Inverse semigroups of partial transformations and θ-classes</i>	215
Chris Rorres, <i>Strong concentration of the spectra of self-adjoint operators</i>	237
Saharon Shelah, <i>A combinatorial problem; stability and order for models and theories in infinitary languages</i>	247
George Gustave Weill, <i>Vector space decompositions and the abstract imitation problem</i>	263
Arthur Thomas White, <i>On the genus of the composition of two graphs</i>	275