

Pacific Journal of Mathematics

HOMOMORPHISMS OF BANACH ALGEBRAS WITH MINIMAL IDEALS

GREGORY FRANK BACHELIS

HOMOMORPHISMS OF BANACH ALGEBRAS WITH MINIMAL IDEALS

GREGORY F. BACHELIS

Let A be a semi-simple Banach algebra with socle F , and let ν be a homomorphism of A into a Banach algebra. It is shown that if I is a minimal one-sided ideal of A , then the restriction of ν to I is continuous. This is then used to deduce continuity properties of the restriction of ν to F . In particular, if F has a bounded left or right approximate identity, then ν is continuous on F .

In [1] and [2] we deduced continuity properties of $\nu|_F$ in case A was a semi-simple annihilator Banach algebra. In this paper we obtain essentially the same results, but without the hypothesis that A be an annihilator algebra.

We first show that the restriction of ν to any minimal one-sided ideal is continuous. The proof is almost purely algebraic. We then show that there exists a constant K such that

$$\|\nu(xy)\| \leq K\|x\|\|y\|, \quad x \in F, \quad y \in \bar{F}.$$

As a corollary we obtain that $\nu|_F$ is continuous if F has a bounded left or right approximate identity.

1. Preliminaries. Throughout this section we assume that A is a complex semi-simple Banach algebra. The socle, F , is defined to be the sum of the minimal right ideals. An idempotent e is called *minimal* if eA is a minimal right ideal. We use without reference the basic facts about the socle of a Banach algebra (see e.g. [7, pp. 45-47]).

The following two lemmas, together with the "Main Boundedness Theorem" of Bade and Curtis ([3, Thm. 2.1], [2, Thm. 4.1]) are the basic ingredients in the proofs that follow. The first lemma is due essentially to Barnes.

LEMMA 1.1. *Let $\{x_1, \dots, x_n\} \subset F$. Then there exist idempotents e and f in F such that $\{x_1, \dots, x_n\} \subset eAf$ and eAf is finite-dimensional.*

Proof. By hypothesis, there exist minimal right ideals, I_1, \dots, I_m , whose sum contains $\{x_1, \dots, x_n\}$. By [4, Thm. 2.2], there exists an idempotent $e \in F$ such that $eA = I_1 + \dots + I_m$. Thus $x_k \in eA$, $1 \leq k \leq n$.

Similarly, there exists an idempotent $f \in F$ such that $x_k \in Af$, $1 \leq k \leq n$. Hence $x_k = ex_k f \in eAf$, $1 \leq k \leq n$.

If u and v are minimal idempotents, and $uAv \neq (0)$, then uAv is one-dimensional [9, Lemma 5.1]. Since e and f are each the sum of minimal idempotents, eAf is finite-dimensional.

LEMMA 1.2. *Let e be a minimal idempotent and suppose that eA is infinite-dimensional. Then there exists a sequence of minimal idempotents $\{g_n\}$ such that $g_n g_m = 0$, $n \neq m$, and $eAg_n \neq (0)$ for all n .*

Proof. Let $g_1 = e$. Assume that g_1, \dots, g_n have been chosen with the desired properties. Let $f = g_1 + \dots + g_n$. Then $f = f^2$ and eAf is finite-dimensional. Thus there exists $x \in eA$ such that $x(1-f) \neq 0$. Since eA is a minimal right ideal, there exists $w \in A$ such that $x(1-f)w = e$. Let $g_{n+1} = (1-f)wex(1-f)$. Then $fg_{n+1} = g_{n+1}f = 0$, so $g_k g_{n+1} = g_{n+1}g_k = 0$, $1 \leq k \leq n$. Also

$$\begin{aligned} g_{n+1}^2 &= (1-f)wex(1-f)(1-f)wex(1-f) \\ &= (1-f)wex(1-f)wex(1-f) \\ &= (1-f)wex(1-f) \\ &= g_{n+1}. \end{aligned}$$

Since e is minimal, g_{n+1} is as well. Since

$$exg_{n+1} = ex(1-f) = x(1-f) \neq 0,$$

$eAg_{n+1} \neq (0)$. The conclusion follows by induction.

NOTE. Lemma 1.2 above takes the place of [2, Lemma 2.2] in what follows. Evidently the latter does not hold in this more general situation, since the norm induced in eA as a subset of $(Ae)^*$ (the set of bounded linear functionals on Ae) need not be equivalent to the given norm on eA (see Remark 2.5).

2. The main results. Throughout this section we assume that A is a complex semi-simple Banach algebra with socle F and that ν is a homomorphism of A into a Banach algebra. We first show the following.

THEOREM 2.1. *If I is a minimal one-sided ideal, then $\nu|I$ is continuous.*

Proof. Suppose that I is a minimal right ideal. Then there exists a minimal idempotent e such that $I = eA$.

Let $J = \{x \in A \mid y \mapsto \nu(xy) \text{ is continuous on } A\}$. Then one verifies that J is a two-sided ideal in A [8, p. 153], and that an idempotent g is in J if and only if $\nu \mid gA$ is continuous.

We may assume eA is infinite-dimensional, since otherwise the conclusion trivially follows. Choose $\{g_n\}$ as given by Lemma 1.2. If $g_n \in J$ then there exists $x_n \in g_n A$ such that $\|x_n\| = 1$ and $\|\nu(x_n)\| > n \|g_n\|$. Since $g_n x_n = x_n$ and $g_m x_n = g_m g_n x_n = 0$, $m \neq n$, the Main Boundedness Theorem [2, Thm. 4.1] implies that $g_n \in J$ for some n . Since $eA g_n \neq (0)$ and J is a left ideal, we have that $eA \cap J \neq (0)$. But J is a right ideal and eA is a minimal right ideal. Thus $e \in eA \subset J$, and $\nu \mid eA$ is continuous.

REMARK 2.2. (cf. [3, p. 597]) If $I = eA$ is an infinite-dimensional minimal right ideal, then it is always possible to construct a discontinuous homomorphism ν of eA into a Banach algebra. For let ϕ be a discontinuous linear functional on eA , and define

$$\|x\|_1 = \|x\| + |\phi(x)|, \quad x \in eA.$$

If $x \in eA$, then $x^n = \lambda e$, λ complex. Thus

$$x^n = (xe)^{n-1}x = \lambda^{n-1}x,$$

so

$$\|x^n\|_1^{1/n} = |\lambda|^{(n-1)/n} \|x\|_1^{1/n}.$$

Hence $|\lambda| = \rho(x)$, the spectral radius of x .

If $x, y \in A$, then

$$\begin{aligned} \|xy\|_1 &= \|xy\| + |\phi(xy)| \\ &\leq \|x\| \|y\| + |\phi(xey)| \\ &\leq \|x\| \|y\| + \rho(x) |\phi(y)| \\ &\leq \|x\| (\|y\| + |\phi(y)|) \\ &\leq \|x\|_1 \|y\|_1, \end{aligned}$$

so $\|\cdot\|_1$ is a normed algebra norm on eA .

Now let B be the completion of eA in this norm and define $\nu: eA \rightarrow B$ by $\nu(x) = x$. Then ν is a discontinuous homomorphism of eA . By the above theorem, ν does not extend to a homomorphism of A .

We now have:

THEOREM 2.3. *Let A be a semi-simple Banach algebra with socle F and let ν be a homomorphism of A into a Banach algebra. Then there exists a constant K such that*

$$\| \nu(xy) \| \leq K \| x \| \| y \| \quad x \in F, \quad y \in \bar{F}$$

Proof. Since F is the sum of the minimal right ideals and also the sum of the minimal left ideals, it follows from Theorem 2.1 that, for any $x \in F$, the mappings $y \rightarrow \nu(xy)$ and $y \rightarrow \nu(yx)$ are continuous on A . Thus it suffices to show that

$$\sup_{x, y \in F} \frac{\| \nu(xy) \|}{\| x \| \| y \|} < \infty .$$

In addition, if $e = e^2 \in F$, then $\nu|_{eA}$ and $\nu|_{Ae}$ are continuous. With these observations, the proof is virtually the same as that of [2, Thm. 4.5], with [2, Lemma 2.1] replaced by Lemma 1.1.

COROLLARY 2.4. *If F has a bounded left or right approximate identity, then ν is continuous on F .*

Proof. If F has a bounded left or right approximate identity, then of course so does \bar{F} . The proof now follows as that of [2, Cor. 4.9], with [1, Cor. 4.9] replaced by Lemma 1.1.

REMARK 2.5. Let X and Y be Banach spaces with $Y \subset X$ and such that the inclusion map $i: Y \rightarrow X$ is continuous and $i(Y)^\perp = X$. Let $B(X, Y)$ denote the bounded operators from X to Y and let $B'(X, Y)$ denote the compact operators from X to Y . Then $B(X, Y)$ is a semi-simple Banach algebra with $B'(X, Y)$ as a closed two-sided ideal. If A is a closed two-sided ideal of $B(X, Y)$ containing $B'(X, Y)$, then A is semi-simple with socle F consisting of those bounded operators from X to Y with finite-dimensional range. Each minimal right ideal of A is linearly homeomorphic to X^* and each minimal left ideal is linearly homeomorphic to Y . Now $i^*: Y^* \rightarrow X^*$ is one-to-one and continuous, but not bi-continuous if $Y \neq X$.

If $X = Y$ and satisfies the metric approximation property [5, p. 178], then F has a bounded left approximate identity, so the above corollary applies to A . If in addition X has a continued bisection, then Johnson [6, Thm. 3.5] has shown that any homomorphism of A into a Banach algebra is continuous on $B'(X, X)$ ($= \bar{F}$). He has also shown that any homomorphism of $B(X, X)$ into a Banach algebra is continuous if X has a continued bisection [6, Thm. 3.3].

REFERENCES

1. G. F. Bachelis, *Homomorphisms of annihilator Banach algebras*, Pacific J. Math., **25** (1968), 229-247.
2. ———, *Homomorphisms of annihilator Banach algebras, II*, Pacific J. Math., **30** (1969) 283-291.

3. W. G. Bade and P. C. Curtis, Jr., *Homomorphisms of commutative Banach algebras*, Amer. J. Math., **82** (1960), 589-608.
4. B. A. Barnes, *A generalized Fredholm theory for certain maps in the regular representations of an algebra*, Canad. J. Math., **20** (1968), 495-504.
5. A. Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc., No. **16** (1955).
6. B. E. Johnson, *Continuity of homomorphisms of algebras of operators*, J. London Math. Soc., **42** (1967), 537-541.
7. C. E. Rickart, *General Theory of Banach Algebras*, Van Nostrand, New York, 1960.
8. J. D. Stein, Jr., *Continuity of homomorphisms of von Neumann algebras*, Amer. J. Math., **91** (1969), 153-159.
9. B. Yood, *Faithful *-representations of normed algebras*, Pacific J. Math., **10** (1960), 345-363.

Received May 20, 1971. The author would like to thank Professor Forrest Miller for helpful discussions.

KANSAS STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 108 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Tom M. (Mike) Apostol, <i>Arithmetical properties of generalized Ramanujan sums</i>	281
David Lee Armacost and William Louis Armacost, <i>On p-thetic groups</i>	295
Janet E. Mills, <i>Regular semigroups which are extensions of groups</i>	303
Gregory Frank Bachelis, <i>Homomorphisms of Banach algebras with minimal ideals</i>	307
John Allen Beachy, <i>A generalization of injectivity</i>	313
David Geoffrey Cantor, <i>On arithmetic properties of the Taylor series of rational functions. II</i>	329
Václáv Chvátal and Frank Harary, <i>Generalized Ramsey theory for graphs. III. Small off-diagonal numbers</i>	335
Frank Rimi DeMeyer, <i>Irreducible characters and solvability of finite groups</i>	347
Robert P. Dickinson, <i>On right zero unions of commutative semigroups</i>	355
John Dustin Donald, <i>Non-openness and non-equidimensionality in algebraic quotients</i>	365
John D. Donaldson and Qazi Ibadur Rahman, <i>Inequalities for polynomials with a prescribed zero</i>	375
Robert E. Hall, <i>The translational hull of an N-semigroup</i>	379
John P. Holmes, <i>Differentiable power-associative groupoids</i>	391
Steven Kenyon Ingram, <i>Continuous dependence on parameters and boundary data for nonlinear two-point boundary value problems</i>	395
Robert Clarke James, <i>Super-reflexive spaces with bases</i>	409
Gary Douglas Jones, <i>The embedding of homeomorphisms of the plane in continuous flows</i>	421
Mary Joel Jordan, <i>Period H-semigroups and t-semisimple periodic H-semigroups</i>	437
Ronald Allen Knight, <i>Dynamical systems of characteristic 0</i>	447
Kwangil Koh, <i>On a representation of a strongly harmonic ring by sheaves</i>	459
Hui-Hsiung Kuo, <i>Stochastic integrals in abstract Wiener space</i>	469
Thomas Graham McLaughlin, <i>Supersimple sets and the problem of extending a retracing function</i>	485
William Nathan, <i>Open mappings on 2-manifolds</i>	495
M. J. O'Malley, <i>Isomorphic power series rings</i>	503
Sean B. O'Reilly, <i>Completely adequate neighborhood systems and metrization</i>	513
Qazi Ibadur Rahman, <i>On the zeros of a polynomial and its derivative</i>	525
Russell Daniel Rupp, Jr., <i>The Weierstrass excess function</i>	529
Hugo Teufel, <i>A note on second order differential inequalities and functional differential equations</i>	537
M. J. Wicks, <i>A general solution of binary homogeneous equations over free groups</i>	543