

Pacific Journal of Mathematics

TRANSVERSAL MATROIDS AND HALL'S THEOREM

RICHARD ANTHONY BRUALDI AND JOHN H. MASON

TRANSVERSAL MATROIDS AND HALL'S THEOREM

RICHARD A. BRUALDI AND JOHN H. MASON

Transversal matroids, not necessarily having finite character, are investigated. It is demonstrated that if $\mathfrak{U}(I) = (A_i: i \in I)$ is an arbitrary family of subsets of an arbitrary set E whose transversal matroid has at least one basis and has no coloops, then $\mathfrak{U}(I)$ has a transversal; in fact, each basis is a transversal of $\mathfrak{U}(I)$ but of no proper subfamily of $\mathfrak{U}(I)$. P. Hall's theorem on the existence of a transversal for a finite family, and indeed an extension of it, can be obtained from this result.

Some necessary conditions for a matroid to be a transversal matroid are derived. One of these is that a transversal matroid of rank r can have at most $\binom{r}{k}$ k -flats having no coloops ($1 \leq k \leq r$).

1. Matroids. Let E be a set. A *matroid* [14, 15, 16] on E is a nonempty collection \underline{M} of subsets of E such that

- (i) $A \in \underline{M}, A' \subseteq A$ imply $A' \in \underline{M}$.
- (ii) $A_1, A_2 \in \underline{M}, |A_1| < |A_2| < \infty$ imply there exists $x \in A_2 \setminus A_1$ such that $A_1 \cup x^\dagger \in \underline{M}$.

The members of \underline{M} are called *independent sets*; those subsets of E not in \underline{M} are *dependent sets*. The matroid \underline{M} on E is said to have *finite character* provided

- (iii) $A \in \underline{M}, A' \in \underline{M}$ for all finite sets $A' \subseteq A$ imply $A \in \underline{M}$.

If E is a finite set, a matroid on E is always a finite character matroid, and a matroid on E is the collection of independent sets of a *combinatorial pregeometry* [4] on E . Finite character matroids arise from many mathematical situations including graphs, vector spaces, geometry, and so on. For details the reader is referred to Crapo and Rota [4]. Matroids not necessarily having finite character also arise in important ways, and we shall be concerned with a certain class of such matroids.

Let \underline{M} be a matroid on E . A *basis* of \underline{M} is a maximal, with respect to set-theoretic inclusion, member of \underline{M} . Bases need not exist as is easily seen by taking E to be an uncountable set and \underline{M} to be all finite or countably infinite subsets of E . However, if E is finite,

[†] The set $\{x\}$ is usually denoted by x .

bases surely exist; if \underline{M} has finite character, then Zorn's lemma in conjunction with the finite character property (iii) guarantees the existence of bases and indeed that every independent set is contained in a basis. It is well-known [13, 2] that in a finite character matroid \underline{M} all bases have the same cardinal number called the *rank* of \underline{M} . A *circuit* is a set $C \subseteq E$ which is a minimal dependent set. If the matroid has finite character, it follows from (iii) that circuits are finite sets.

If \underline{M} is a matroid on E and $A \subseteq E$, then we define \underline{M}_A by

$$\underline{M}_A = \{F: F \subseteq A, F \in \underline{M}\}.$$

It is clear that \underline{M}_A is a matroid on A , called the *restriction* of \underline{M} to A . If $(E_i: i \in I)$ is a family of pairwise disjoint sets and \underline{M}_i is a matroid on E_i ($i \in I$), then a matroid \underline{M} on $E = \bigcup_{i \in I} E_i$ can be defined by

$$\underline{M} = \left\{ \bigcup_{i \in I} A_i: A_i \in \underline{M}_i (i \in I) \right\}.$$

The matroid \underline{M} is called the *direct sum* of \underline{M}_i ($i \in I$) and is denoted by $\bigoplus_{i \in I} \underline{M}_i$. If $|I| < \infty$ and \underline{M}_i has finite rank r_i ($i \in I$), then $\bigoplus_{i \in I} \underline{M}_i$ has finite rank $\sum_{i \in I} r_i$.

If \underline{M} is a matroid on E , then \underline{M} is *connected* or *nonseparable* provided it is impossible to partition E into nonempty sets E_1, E_2 in such a way that $\underline{M} = \underline{M}_{E_1} \oplus \underline{M}_{E_2}$. The element x of E is a *loop* of \underline{M} if $\{x\} \in \underline{M}$; thus loops can be part of no independent sets. The element x is called a *coloop* or *isthmus* provided $A \cup x \in \underline{M}$ whenever $A \in \underline{M}$; thus coloops are part of every basis. If x is either a loop or coloop, then $\underline{M} = \underline{M}_{\{x\}} \oplus \underline{M}_{E \setminus x}$ so that \underline{M} cannot be connected. If X is a set of coloops of \underline{M} , then $\underline{M} = \underline{M}_X \oplus \underline{M}_{E \setminus X}$ where \underline{M}_X is the *free matroid* or *Boolean algebra* on X . In what is to follow, matroids having no coloops play an important role; such matroids will be called *coloop-free*. For these matroids it is impossible to 'split off' a Boolean algebra.

Finally, we introduce the notion of a flat. A set $F \subseteq E$ is a *flat* of the matroid \underline{M} on E or is *closed* provided $A \subseteq F, A \in \underline{M}$ and $x \in E \setminus F$ imply that $A \cup x \notin \underline{M}$. If \underline{M}_F has finite rank, then this means that enlarging F in any way increases the rank or equivalently that given $x \in E \setminus F$ there is no circuit C with $x \in C \subseteq F \cup x$. If the rank of \underline{M}_F is $k < \infty$, then F is called a *k-flat*. Observe that each coloop is a 1-flat (but not conversely) and that the set of all loops is the only 0-flat. In case \underline{M} has finite rank, the collection of flats form a geometric lattice [4] with respect to set-theoretic inclusion.

2. **Transversal matroids.** An important class of matroids, dis-

covered by Edmonds and Fulkerson [6], are those known as transversal matroids. These are defined as follows. Let $\mathfrak{A} = \mathfrak{A}(I) = (A_i; i \in I)$ be a family of subsets of E . A set $T \subseteq E$ is a *transversal* of \mathfrak{A} provided there is a bijection $\theta: T \rightarrow I$ such that $x \in A_{\theta(x)}$ ($x \in T$). If θ is only an injection, then T is a *partial transversal*; in this case T is a transversal of the subfamily $\mathfrak{A}(K) = (A_i; i \in K)$ where $K = \theta(T)$. If $\underline{M}(\mathfrak{A})$ denotes the collection of all partial transversals of \mathfrak{A} , then $\underline{M}(\mathfrak{A})$ is a matroid on E [6, 11]. If each element of E is a member of only finitely many sets of the family \mathfrak{A} , then $\underline{M}(\mathfrak{A})$ is a finite-character matroid. A matroid \underline{M} on E is a *transversal matroid* provided there is a family \mathfrak{A} of subsets of E such that $\underline{M} = \underline{M}(\mathfrak{A})$.

The bases of the transversal matroid $\underline{M}(\mathfrak{A})$, if there are any, are the maximal partial transversals of \mathfrak{A} . These need not however be transversals of \mathfrak{A} . However the following theorem is proved in Brualdi and Scrimger [1], although not stated in this form.

THEOREM 2.1. *Let $\mathfrak{A}(I) = (A_i; i \in I)$ be a family of subsets of E . Let B be a basis of $\underline{M}(\mathfrak{A})$ and let $\mathfrak{A}' = \mathfrak{A}(K) = (A_i; i \in K)$ be any subfamily of \mathfrak{A} of which B is a transversal. Then $\underline{M}(\mathfrak{A}) = \underline{M}(\mathfrak{A}')$ and every basis of $\underline{M}(\mathfrak{A})$ is a transversal of \mathfrak{A}' .*

One of the results of this paper is that in Theorem 2.1 K can only be I if $A_i \neq \emptyset$ ($i \in I$) and the matroid $\underline{M}(\mathfrak{A})$ has no coloops. Before getting to this, it is convenient to place our discussion in a graph-theoretic setting, for some of our proofs are graph-theoretic in nature.

A *bipartite graph* may be regarded as a triple (X, Δ, Y) where X and Y are disjoint sets and $\Delta \subseteq X \times Y$. The members of Δ are the *edges* of the graph, which we regard as undirected. Let $A \subseteq X$, $B \subseteq Y$. Then A and B are *linked* in the bipartite graph (or A is linked to B or B is linked to A) provided there is a bijection $\theta: A \rightarrow B$ with $\Delta' = \{(x, \theta(x)): x \in A\} \subseteq \Delta$. The bijection θ is called a *linking* of A to B and the members of Δ' are the *edges of the linking*. If $\mathfrak{A}(I) = (A_i; i \in I)$ is a family of subsets of a set E , then a bipartite graph (E, Δ, I) can be associated where $(e, i) \in \Delta$ if and only if $e \in A_i$ ($e \in E, i \in I$). The partial transversals of \mathfrak{A} are precisely those subsets of E which are linked to some subset of I . It is then clear that a matroid \underline{M} on a set X is a transversal matroid provided there is a bipartite graph (X, Δ, Y) such that \underline{M} consists of those subsets of X which are linked to at least one subset of Y . Such a bipartite graph is said to *induce* \underline{M} on X . The bipartite graph likewise induces a matroid \underline{M}' on Y . For $x \in X$, $A \subseteq X$, we set $\Delta(x) = \{y \in Y: (x, y) \in \Delta\}$ and $\Delta(A) = \bigcup_{x \in A} \Delta(x)$. If $y \in Y$, $B \subseteq Y$, then $\Delta(y)$ and $\Delta(B)$ are defined analogously.

3. Induced matroids. If we phrase the first part of Theorem 2.1 in terms of bipartite graphs, it becomes: *if (X, Δ, Y) is a bipartite graph inducing the matroid \underline{M} on X , if B is a basis of \underline{M} and Z is any subset of Y to which B is linked, then the bipartite graph (X, Δ', Z) where $\Delta' = \Delta \cap \{X \times Z\}$ already induces \underline{M} on X .*

The main result of the section deals with a closer analysis of the above situation. Before stating it we record a lemma.

LEMMA 3.1. *Let (X, Δ, Y) be a bipartite graph inducing the matroid \underline{M} on X . Assume that \underline{M} is coloop-free. Then if B is a basis of \underline{M} and $z \in B$, there exists $x \in X \setminus B$ such that $\{B \setminus z\} \cup x$ is a basis of \underline{M} .*

The result is true for any coloop-free finite-character matroid. The matroid \underline{M} in the lemma above is not necessarily a finite-character one, but is assumed to be a transversal matroid.

Proof. Since \underline{M} is coloop-free, $B \neq X$. Assume $\{B \setminus z\} \cup x$ is not a basis of \underline{M} for all $x \in X \setminus B$. Let $A \in \underline{M}$ with $z \notin A$. We will show that $A \cup z \in \underline{M}$, so that z is a coloop of \underline{M} , a contradiction.

If $A \subseteq B$, then $A \cup z \in \underline{M}$. Thus we may assume $A \setminus B \neq \emptyset$. Let Δ_1 be the edges of a linking of B to a set $Z_1 \subset Y$ and Δ_2 the edges of a linking of A to a set $Z_2 \subset Y$. (We could assume from the result mentioned above that $Z_2 \subseteq Z_1$.) Each $x \in A \setminus B$ determines a path P_x beginning at x whose edges alternate in Δ_2 and Δ_1 . Let Δ_i^x denote the set of edges of Δ_i on this path ($i = 1, 2$). If P_x is either an infinite path or terminates at an element of $Z_2 \setminus Z_1$, then $\{\Delta_1 \setminus \Delta_1^x\} \cup \Delta_2^x$ is the set of edges of a linking of $B \cup x$ to a subset of Y . This contradicts the basis property of B . The only other alternative is that P_x terminates at an element w_x of $B \setminus A$. If $w_x = z$, then $\{\Delta_1 \setminus \Delta_1^x\} \cup \Delta_2^x$ is the set of edges of a linking of $\{B \setminus z\} \cup x$ to a subset of Y . It follows as in [1], that $\{B \setminus z\} \cup x$ is a basis of \underline{M} . Since we are assuming this is not the case $w_x \neq z$. Since this is true for all $x \in A \setminus B$, the path Q_z determined by z whose edges alternate in Δ_1 and Δ_2 must either be infinite or terminate in $Z_1 \setminus Z_2$. For, if Q_z terminates at some $x \in A \setminus B$, the only other alternative, we would have that P_x terminates at z and thus $w_x = z$. Thus following the above convention, $\{\Delta_2 \setminus \Delta_2^z\} \cup \Delta_1^z$ is the set of edges of a linking of $A \cup \{z\}$ to a subset of Y , so that $A \cup \{z\} \in \underline{M}$. Since this is true for all $A \in \underline{M}$, it follows that z is a coloop. This completes the proof of the lemma.

We now state and prove the main result.

THEOREM 3.2. *Let (X, Δ, Y) be a bipartite graph inducing the matroid \underline{M} on X . Assume that \underline{M} is coloop-free. Then if B is any basis of \underline{M} and Z is any subset of Y to which B is linked, then $\Delta(X) = Z$.*

We point out once more that since we are not assuming \underline{M} has finite character, the matroid \underline{M} may not have any bases, in which case the theorem says nothing.

Proof. There is no loss in generality in assuming that $\Delta(X) = Y$, for those elements $y \in Y$ with $\Delta(y) = \phi$ play no role whatsoever. The conclusion of the theorem is then that $Z = Y$.

Let B_1 be a basis of \underline{M} with B_1 linked to $Z_1 \subseteq Y$. If $x \in X \setminus B_1$, then the maximality of B_1 implies that $\Delta(x) \subseteq Z_1 \subseteq \Delta(B_1)$. Suppose $Y \setminus Z_1 \neq \phi$. Then there exists a $w \in B_1$ such that $\Delta(w) \cap \{Y \setminus Z_1\} \neq \phi$. By Lemma 2.1 there exists $x \in X \setminus B_1$ such that $B_2 = \{B_1 \setminus w\} \cup x$ is a basis of \underline{M} . By Theorem 2.1 B_2 is linked to Z_1 . Since $w \notin B_2$, this means that $B_2 \cup w$ is linked to $Z_1 \cup z$ where $z \in \Delta(w) \cap \{Y \setminus Z_1\}$. Hence $B_2 \cup w \in \underline{M}$, and this contradicts the fact that B_2 is a basis of \underline{M} . The $Y = Z_1$ and the theorem is proved.

In case the matroid \underline{M} has finite rank, the above theorem takes on an appealing form. It is proved by Mason in [9, 10]. It can also be proved using a canonical decomposition of bipartite graphs which was derived by Dulmage and Mendelsohn [5] as an extension of a result of Ore [12].

COROLLARY 3.3. *Let (X, Δ, Y) be a bipartite graph inducing the matroid M of finite rank r on X . If M is coloop-free, then $|\Delta(X)| = r$.*

In this case $|B| = |Z| = r$. Effectively what the corollary says is that a coloop-free transversal matroid \underline{M} on a set X with rank r can only be induced by bipartite graphs (X, Δ, Y) where $|Y| = r$. Observe that the corollary applies in case \underline{M} is connected with $|X| > 1$. As an example it applies to any matroid of the form $\underline{M} = \mathcal{P}_r(X) = \{A \subseteq X: |A| \leq r\}$, $1 \leq r \leq |X|$. This matroid is easily seen to be a connected, transversal matroid. If $r = |X| - 1$, it is just a circuit.

THEOREM 3.4. *Let (X, Δ, Y) be a bipartite graph inducing the matroid \underline{M} on X . Let $A \subseteq X$ and suppose \underline{M}_A is coloop-free. If B' is any basis of \underline{M}_A , there is a unique subset of Y , namely $\Delta(A)$, to which B' is linked. In particular, if \underline{M}_A is coloop-free of rank t , then $|\Delta(A)| = t$.*

Proof. It is clear that (A, Δ', Y) induces \underline{M}_A on A where $\Delta' = \Delta \cap \{A \times Y\}$. Let B' be a basis of \underline{M}_A and let Z' be any subset of Y to which B' is linked. Applying Theorem 3.2 to this new bipartite graph, we conclude that $\Delta'(A) = Z'$. Since $\Delta(A) = \Delta'(A)$, this established the theorem.

THEOREM 3.5. *Let the bipartite graph (X, Δ, Y) induce the matroid \underline{M} on X and the matroid \underline{M}' on Y . If \underline{M} has a basis and is coloop-free, then \underline{M}' is the free matroid on Y .*

Proof. Let B be a basis of \underline{M} . Since \underline{M} has no coloops, it follows from Theorem 3.2 that $\Delta(X) \in \underline{M}'$. Thus if $y \in \Delta(X)$, y is a coloop of \underline{M}' , while if $y \in Y \setminus \Delta(X)$, y is a loop of \underline{M}' . The conclusion is now obvious.

A particular case of Theorem 3.5 asserts that if one of \underline{M} and \underline{M}' has a basis (e.g. if one has finite character), not both of \underline{M} and \underline{M}' can be coloop-free and, in particular, not both can be connected.

4. Applications to transversal theory. Let $\mathfrak{A} = \mathfrak{A}(I) = (A_i: i \in I)$ be a family of subsets of a set E . For $K \subseteq I$, let $A(K) = \bigcup_{i \in K} A_i$. If $|I| < \infty$, so that \mathfrak{A} is a finite family, then the well-known theorem of P. Hall [7] asserts that \mathfrak{A} has a transversal if and only if $|A(K)| \geq |K|$ for all $K \subseteq I$. If $|I| = \infty$ but each A_i is a finite set ($i \in I$), the extension due to M. Hall Jr. [8] of this result asserts that \mathfrak{A} has a transversal if and only if $|A(K)| \geq |K|$ for all finite sets $K \subseteq I$. We offer the following theorem.

THEOREM 4.1. *Let $\mathfrak{A}(I) = (A_i: i \in I)$ be a family of nonempty subsets of a set E . If the matroid $\underline{M}(\mathfrak{A})$ has a basis and is coloop-free, then the family $\mathfrak{A}(I)$ has a transversal.*

Proof. Assume, without loss in generality, that $I \cap E = \emptyset$, and consider the bipartite graph (E, Δ, I) where $\Delta = \{(e, i): e \in A_i, i \in I\}$. This bipartite graph induces the matroid $\underline{M}(\mathfrak{A})$ on E . If B is any basis of $\underline{M}(\mathfrak{A})$ and J is any subset of I to which B is linked, then from Theorem 3.2 we conclude that $\Delta(E) = J$. On the other hand since $A_i \neq \emptyset$ ($i \in I$), $\Delta(E) = I$. Hence $J = I$ and B is linked to I in the bipartite graph. But this means that B is a transversal of the family $\mathfrak{A}(I)$.

If in the theorem each element of E is a member of only finitely many A 's, then $\underline{M}(\mathfrak{A})$ is a finite character matroid and hence has bases.

COROLLARY 4.2. *If the matroid of a family $\mathfrak{A}(I)$ of nonempty*

subsets of a set E with $|E| > 1$ has a basis and is connected, then the family has a transversal.

A connected matroid on a set with more than one element cannot have any coloops.

A more detailed analysis produces the following theorem which contains P. Hall's theorem as a special case, but not necessarily M. Hall's theorem. (On the other hand, M. Hall's theorem follows easily from P. Hall's theorem through a simple application of Rado's selection principle or other theorems dependent on the axiom of choice.)

To say that a matroid \underline{M} has only a finite number of coloops is equivalent to saying that an infinite Boolean algebra can not be "split off" from \underline{M} .

THEOREM 4.3. *Let $\mathfrak{A}(I) = (A_i; i \in I)$ be a family of subsets of a set E . Assume the matroid $\underline{M}(\mathfrak{A})$ has a basis and only a finite number of coloops. Then $\mathfrak{A}(I)$ has a transversal if and only if*

$$(1) \quad |A(K)| \geq |K| \quad (K \text{ finite, } K \subseteq I).$$

Proof. Let (E, Δ, I) be the bipartite graph associated as before with the family $\mathfrak{A}(I)$. Let F be the set of coloops of $\underline{M} = \underline{M}(\mathfrak{A})$ and $E' = E \setminus F$. By assumption F is a finite set. The matroid $\underline{M}_{E'}$ has a basis, since \underline{M} has a basis: if B is a basis of \underline{M} , then $B \setminus F$ is a basis of $\underline{M}_{E'}$. Moreover the matroid $\underline{M}_{E'}$ has no coloops. For, if x were a coloop of $\underline{M}_{E'}$ and $A \in \underline{M}$, then $\{A \cap E'\} \cup x \in \underline{M}$ and thus $A \cup x \subseteq \{A \cap E'\} \cup x \cup F \in \underline{M}$. Thus x is a coloop of \underline{M} with $x \notin F$, and this contradicts the choice of F . Let B' be a basis of $\underline{M}_{E'}$. By Theorem 3.4 B' is linked in the bipartite graph to $J = \Delta(E')$ and J is the only subset of I having this property. Since $J = \Delta(E')$, it follows that $\Delta(I \setminus J) \subseteq F$. Thus $\mathfrak{A}(I)$ has a transversal if and only if the subfamily $\mathfrak{A}(I \setminus J)$, which is a family of subsets of the *finite* set F , has a transversal.

Suppose now condition (1) is satisfied for all finite $K \subseteq I$ and thus for all finite $K \subseteq I \setminus J$. Since F is a finite set and $|A(K)| \leq |F|$ for all $K \subseteq I \setminus J$, the set $I \setminus J$ must be finite. Since $B' \cup F \in \underline{M}$ and B' is linked only to J , it follows that F is linked to a subset of $I \setminus J$, so that $|I \setminus J| \geq |F|$. But then

$$|F| \geq |A(I \setminus J)| \geq |I \setminus J| \geq |F|,$$

so that $|I \setminus J| = |F|$. Hence F is linked to $I \setminus J$. This means that $\mathfrak{A}(I)$ has a transversal, namely $B' \cup F$. Since condition (1) is obviously necessary for $\mathfrak{A}(I)$ to have a transversal, the proof of the theorem is complete.

The proof of the theorem indicates how to find a single set $\bar{K} \subseteq I$ such that $\mathfrak{A}(I)$ has a transversal if and only if $|A(\bar{K})| \geq |\bar{K}|$. For taking $\bar{K} = I \setminus J$, it was demonstrated in the proof that if $|A(\bar{K})| \geq |\bar{K}|$, then $\mathfrak{A}(I)$ has a transversal, while if $|A(\bar{K})| < |\bar{K}|$ this would mean that $|F| < |\bar{K}|$ so that $\mathfrak{A}(I)$ could not have a transversal.

As a corollary to Theorem 4.3 we obtain P. Hall's theorem [7].

COROLLARY 4.4. (P. Hall). *Let $\mathfrak{A} = (A_i: 1 \leq i \leq n)$ be a family of subsets of E . Then \mathfrak{A} has a transversal if and only if*

$$|A(K)| \geq |K| \quad (K \subseteq \{1, \dots, n\}).$$

In this case the matroid $\underline{M}(\mathfrak{A})$ has finite rank, so that it has a basis and can only have a finite number of coloops.

We also remark here that Theorem 4.3 applies to any family $\mathfrak{A}(I)$ of subsets of E such that each element of E is a member of only finitely many A 's and the matroid $\underline{M}(\mathfrak{A})$ has only a finite number of coloops.

If in Theorem 4.3 the matroid $\underline{M}(\mathfrak{A})$ has an infinite number of coloops, then condition (1) is no longer sufficient for \mathfrak{A} to have a transversal. This is already seen from M. Hall's much quoted example [8] where $I = E = \{1, 2, \dots\}$ and $A_1 = E$, $A_i = \{i + 1\}$ ($i \geq 2$). In this case $E \in \underline{M}(\mathfrak{A})$ so that each element of E is a coloop. Condition (1) is satisfied but there is no transversal.

5. Transversal matroids. In general it is difficult to decide whether a given matroid is a transversal matroid. A characterization of finite-character transversal matroids in terms of a rank inequality on unions of circuits is given by Mason [9, 10], but it is difficult to check. The following result is contained implicitly in [1].

THEOREM 5.1. *Let \underline{M} be a transversal matroid on a set E . Let B_1 and B_2 be bases of \underline{M} . Then there exists a bijection $\sigma: B_1 \rightarrow B_2$ such that both $\{B_1 \setminus x\} \cup \sigma(x)$ and $\{B_2 \setminus \sigma(x)\} \cup x$ are bases for all $x \in B_1$.*

For finite-character matroids a σ satisfying the exchange property in this theorem can always be defined, as is proved in [2], but σ need not be a bijection or injection. Indeed the example given in [2] for which it is impossible to define a bijective σ amounts to the cycle matroid of the complete graph on 4 nodes, K_4 . (The *cycle matroid* of a graph is the matroid on its edge set such that a set of edges is independent if and only if it does not contain the edges of a polygon; thus the circuits are the edge sets of polygons.) Thus Theorem

5.1 furnishes a necessary, but not sufficient, condition for a matroid to be a transversal matroid. We shall use the results of § 3 to obtain other necessary conditions.

THEOREM 5.2. *Let \underline{M} be a transversal matroid on a set E with finite rank r . Let k be any integer with $1 \leq k \leq r$. Then \underline{M} has at most $\binom{r}{k}$ coloop-free k -flats.*

Proof. Let (E, Δ, Y) be a bipartite graph which induces \underline{M} on E . By Corollary 2.2 we may assume $|Y| = r$. Let $1 \leq k \leq r$ and let $(F_j: j \in J)$ be the family of distinct coloop-free k -flats of \underline{M} , indexed by J . We need to show that $|J| \leq \binom{r}{k}$. By Theorem 3.4, since \underline{M}_{F_j} is coloop-free with rank k , $|\Delta(F_j)| = k$ ($j \in J$). Suppose $|J| > \binom{r}{k}$. It would then follow, since $|Y| = r$, that $\Delta(F_{j_1}) = \Delta(F_{j_2})$ for some $j_1, j_2 \in J$ with $j_1 \neq j_2$. This would mean that $|\Delta(F_{j_1} \cup F_{j_2})| = k$ and thus that $\underline{M}_{F_{j_1} \cup F_{j_2}}$ has rank k . But since F_{j_1}, F_{j_2} are distinct k -flats, $\underline{M}_{F_{j_1} \cup F_{j_2}}$ has rank greater than k . This is a contradiction and the theorem is proved.

As an example consider once again the rank 3 cycle matroid \underline{M} on the set of edges $E = \{1, 2, \dots, 6\}$ of the complete 4-graph K_4 . (Figure 1) The set $\{1, 2, 5\}, \{2, 3, 6\}, \{3, 4, 5\}, \{1, 4, 6\}$ are all coloop-free 2-flats of \underline{M} . Since $4 > \binom{3}{2}$, it follows by Theorem 5.2 that \underline{M} is not a transversal matroid.




FIGURE 1

Theorem 5.2 can also be used to demonstrate that matroids of infinite rank are not transversal. This is so because if \underline{M} is a transversal matroid on E and $A \subseteq E$, then \underline{M}_A is also a transversal matroid. Thus if A is chosen so that \underline{M}_A has finite rank, we can use Theorem 5.2 on \underline{M}_A .

The conditions on the number of k -flats as given in Theorem 5.2 are not sufficient to guarantee that a matroid is a transversal matroid. To obtain an example, consider the rank 4 cycle matroid \underline{M} on the set of edges $E = \{1, 2, \dots, 7\}$ of the graph of Figure 2. Then \underline{M} has two coloop-free 2-flats, namely $\{3, 4, 7\}$ and $\{4, 5, 6\}$ and three coloop-free 3-flats, namely $\{1, 2, 3, 6\}, \{1, 2, 5, 7\}, \{3, 4, 5, 6, 7\}$. Hence the




FIGURE 2

conditions on the numbers of coloop-free k -flats are satisfied ($k = 1$ and $k = r$ are always trivially satisfied). But \underline{M} is not a transversal matroid as can be seen from Theorem 5.1. For $B_1 = \{1, 2, 4, 6\}$ and $B_2 = \{1, 3, 5, 7\}$ are both bases with $\{B_1 \setminus 4\} \cup y$ and $\{B_2 \setminus y\} \cup 4$ both bases only for $y = 7$ of B_2 , and $\{B_1 \setminus 2\} \cup z$ and $\{B_2 \setminus z\} \cup 2$ are both bases only for $z = 7$ of B_2 . Thus the bijection of Theorem 5.1 cannot exist, so that \underline{M} is not a transversal matroid.

Theorem 5.2 is interesting because it gives a bound on the number of coloop-free k -flats of a transversal matroid on E which does not depend on the size of E but on the rank of the matroid. The total number of k -flats cannot be bounded in terms of r . For if E is a set with $|E| \geq r$ and $\underline{M} = \mathcal{S}_r(E) = \{A \subseteq E: |A| \leq r\}$, then \underline{M} is a transversal matroid of rank r and every subset of E of k elements, $1 \leq k \leq r - 1$, is a k -flat. Hence \underline{M} has $\binom{|E|}{k}$ k -flats ($1 \leq k \leq r - 1$), all of which have coloops.

Before getting to another necessary condition for a matroid to be transversal, we require a definition. Let \underline{M} be a matroid on a set E . We say that \underline{M} has the *direct sum property* provided:

Whenever $(E_k: k \in K)$ is a family of pairwise disjoint subsets of E such that \underline{M}_{E_k} is a coloop-free matroid with basis on E_k ($k \in K$), then

$$\underline{M}_{E_k \cup E_l} = \underline{M}_{E_k} \oplus \underline{M}_{E_l} \quad (k, l \in K, k \neq l)$$

imply

$$\underline{M}_{\bigcup (E_j: j \in J)} = \bigoplus (\underline{M}_{E_j}: j \in J).$$

A matroid need not have the direct sum property as the cycle matroid \underline{M} on the set of edges $E = \{1, 2, \dots, 9\}$ of the graph of Figure 3 shows. If we take $E_1 = \{1, 2, 3\}$, $E_2 = \{4, 5, 6\}$, $E_3 = \{7, 8, 9\}$, then \underline{M}_{E_i} is coloop-free ($1 \leq i \leq 3$) and $\underline{M}_{E_i \cup E_j} = \underline{M}_{E_i} \oplus \underline{M}_{E_j}$ ($1 \leq i \neq j \leq 3$). But $\underline{M}_{E_1 \cup E_2 \cup E_3} \neq \underline{M}_{E_1} \oplus \underline{M}_{E_2} \oplus \underline{M}_{E_3}$, for $\{2, 6, 9\}$ is a circuit whose intersections with E_i are independent ($1 \leq i \leq 3$).

We do, however, have the following theorem.




FIGURE 3

THEOREM 5.3. *Let \underline{M} be a transversal matroid on E . Then \underline{M} has the direct sum property.*

Proof. Let (E, Δ, Y) be a bipartite graph which induces \underline{M} on E . Let $(E_k: k \in K)$ be a family of pairwise disjoint subsets of E such that \underline{M}_{E_k} is a coloop-free matroid with basis on E_k ($k \in K$) and $\underline{M}_{E_k \cup E_l} = \underline{M}_{E_k} \oplus \underline{M}_{E_l}$ ($k, l \in K, k \neq l$). Since \underline{M}_{E_k} has a basis B_k , it follows from Theorem 4.2 that B_k is linked to $\Delta(E_k)$ and to no other subset of Y ($k \in K$). Since $\underline{M}_{E_k \cup E_l} = \underline{M}_{E_k} \oplus \underline{M}_{E_l}$ ($k \neq l$), $B_k \cup B_l$ is a basis of $\underline{M}_{E_k \cup E_l}$ where $B_k \cap B_l = \phi$. Since $\underline{M}_{E_k} \oplus \underline{M}_{E_l}$ is coloop-free, $B_k \cup B_l$ is linked to $\Delta(E_k \cup E_l) = \Delta(E_k) \cup \Delta(E_l)$. Since B_k , resp. B_l , is only linked to $\Delta(E_k)$, resp. $\Delta(E_l)$, it follows that $\Delta(E_k) \cap \Delta(E_l) = \phi$. Since this is true for all $k, l \in K$ with $k \neq l$, the result follows.

Since the matroid of the graph of Figure 3 does not have the direct sum property, it follows it is not a transversal matroid.

The direct sum property does not characterize transversal matroids among all matroids. The cycle matroid of the graph of Figure 2 has the direct sum property but is not transversal as we have already seen. In fact the direct sum property holds trivially, for if $F \subseteq E$ with \underline{M}_F coloop-free $|F| \leq 3$. Since $|E| = 7$ in this case, the direct sum property is valid.

To conclude we wish to mention one further consequence of the results of § 3. For this we need another definition which, to keep things simple we make only for finite character matroids. Let \underline{M} be a finite character matroid on E , and let $F \subseteq E$ with B a basis of $\underline{M}_{E \setminus F}$. Let

$$\underline{M}_{\otimes F} = \{A: A \subseteq F, A \cup B \in \underline{M}\}.$$

Then it is well-known [14, 3] that $\underline{M}_{\otimes F}$ is independent of the choice of basis B and that $\underline{M}_{\otimes F}$ is a finite-character matroid on F , called the *contraction of \underline{M} to F* . The contraction of a transversal matroid need not be a transversal matroid. An example which contains 2-element circuits (thus not a combinatorial geometry [4]) is given in [9]. The cycle matroid \underline{M} on the set of edges of the graph of Figure




FIGURE 4

4 is a transversal matroid, as is not difficult to see. If we take $F = E \setminus e$, then $\underline{M}_{\otimes F}$ is isomorphic to the matroid of the graph of Figure 3 and hence is not a transversal matroid. It is therefore of interest to determine when the contraction of a transversal matroid is also a transversal matroid. We offer the following theorem.

THEOREM 5.4. *Let \underline{M} be a finite-character transversal matroid on a set E . Let $F \subseteq E$ and suppose $\underline{M}_{E \setminus F}$ is coloop-free. Then $\underline{M}_{\otimes F}$ is a (finite-character) transversal matroid.*

Proof. Let the bipartite graph (E, \mathcal{A}, Y) induce the matroid \underline{M} on E . Since $\underline{M}_{E \setminus F}$ has no coloops, it follows from Theorem 3.4 that if B is a basis of $\underline{M}_{E \setminus F}$ then B is linked only to the subset $\mathcal{A}(E/F)$ of Y . Let the bipartite graph (F, \mathcal{A}', Z) be defined by $Z = Y \setminus \mathcal{A}(E \setminus F)$ and $\mathcal{A}' = \mathcal{A} \cap \{F \times Z\}$. Let $A \subseteq F$. Then $A \in \underline{M}_{\otimes F}$ if and only if $A \cup B \in \underline{M}$; $A \cup B \in \underline{M}$ if and only if $A \cup B$ is linked in (E, \mathcal{A}, Y) ; $A \cup B$ is linked in (E, \mathcal{A}, Y) if and only if A is linked in (F, \mathcal{A}', Z) . Hence the bipartite graph (F, \mathcal{A}', Z) induces $\underline{M}_{\otimes F}$ on F so that $\underline{M}_{\otimes F}$ is a transversal matroid.

There is no difficulty in obtaining examples where $\underline{M}_{E \setminus F}$ has coloops but $\underline{M}_{\otimes F}$ is a transversal matroid. In fact the matroid of a graph which is a triangle already furnishes an example.

REFERENCES

1. R. A. Brualdi and E. B. Scrimger, *Exchange systems, matchings and transversals*, J. Combinatorial Theory, **5** (1968), 244-257.
2. R. A. Brualdi, *Comments on bases in dependence structures*, Bull. Australian Math. Soc., **1** (1969), 161-167.
3. ———, *Admissible mappings between dependence spaces*, Proc. London Math. Soc., (3) **21** (1970), 296-312.
4. H. Crapo and G.-C. Rota, *Combinatorial Geometries* (M.I.T. Press; Cambridge, 1971).
5. A. L. Dulmage and N. S. Mendelsohn, *Coverings of bipartite graphs*, Canad. J. Math., **10** (1958), 517-534.
6. J. Edmonds and D. R. Fulkerson, *Transversals and matroid partition*, J. Res. Nat. Bur. Standards **69B** (1965), 73-77.
7. P. Hall, *On representatives of subsets*, J. London Math. Soc., **10** (1935), 26-30.

8. M. Hall, Jr., *Distinct representatives of subsets*, Bull. Amer. Math. Soc., **54** (1948), 922-928.
9. J. H. Mason, *Representations of independence spaces*, Ph. D. Dissertation, University of Wisconsin, 1969.
10. ———, *A characterization of transversal independence spaces*, (to be published).
11. L. Mirsky and H. Perfect, *Applications of the notion of independence to problems of combinatorial analysis*, J. Combinatorial Theory, **3** (1967), 327-357.
12. O. Ore, *Graphs and matching theorems*, Duke. J. Math., **22** (1955), 625-639.
13. R. Rado, *Axiomatic treatment of rank in infinite sets*, Canadian J. Math., **1** (1949), 337-343.
14. W. T. Tutte, *Lectures on matroids*, J. Res. Nat. Bur. Standards, **69B** (1965), 1-47.
15. ———, *Introduction to the Theory of Matroids*, (American Elsevier Publishing Co., N.Y. 1971)
16. H. Whitney, *On the abstract properties of linear independence*, Amer. J. Math., **57** (1935), 507-533.

Received March 31, 1970 and in revised form May 25, 1971.

UNIVERSITY OF WISCONSIN

AND

THE OPEN UNIVERSITY, BLETCHLEY, ENGLAND

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics

Vol. 41, No. 3

BadMonth, 1972

| | |
|---|-----|
| George E. Andrews, <i>Two theorems of Gauss and allied identities proved arithmetically</i> | 563 |
| Stefan Bergman, <i>On pseudo-conformal mappings of circular domains</i> | 579 |
| Beverly L. Brechner, <i>On the non-monotony of dimension</i> | 587 |
| Richard Anthony Brualdi and John H. Mason, <i>Transversal matroids and Hall's theorem</i> | 601 |
| Philip Throop Church and James Timourian, <i>Differentiable maps with 0-dimensional critical set. I</i> | 615 |
| John H. E. Cohn, <i>Squares in some recurrent sequences</i> | 631 |
| Robert S. Cunningham, Edgar Andrews Rutter and Darrell R. Turnidge, <i>Rings of quotients of endomorphism rings of projective modules</i> | 647 |
| Eldon Dyer and S. Eilenberg, <i>An adjunction theorem for locally equiconnected spaces</i> | 669 |
| Michael W. Evans, <i>On commutative P. P. rings</i> | 687 |
| Ronald Lewis Graham, Hans Sylvain Witsenhausen and Hans Zassenhaus, <i>On tightest packings in the Minkowski plane</i> | 699 |
| Stanley P. Gudder, <i>Partial algebraic structures associated with orthomodular posets</i> | 717 |
| Karl Edwin Gustafson and Gunter Lumer, <i>Multiplicative perturbation of semigroup generators</i> | 731 |
| Kurt Kreith and Curtis Clyde Travis, Jr., <i>Oscillation criteria for selfadjoint elliptic equations</i> | 743 |
| Lawrence Louis Larmore, <i>Twisted cohomology theories and the single obstruction to lifting</i> | 755 |
| Jorge Martinez, <i>Tensor products of partially ordered groups</i> | 771 |
| Robert Alan Morris, <i>The inflation-restriction theorem for Amitsur cohomology</i> | 791 |
| Leo Sario and Cecilia Wang, <i>The class of (p, q)-biharmonic functions</i> | 799 |
| Manda Butchi Suryanarayana, <i>On multidimensional integral equations of Volterra type</i> | 809 |
| Kok Keong Tan, <i>Fixed point theorems for nonexpansive mappings</i> | 829 |