

Pacific Journal of Mathematics

**THE INFLATION-RESTRICTION THEOREM FOR AMITSUR
COHOMOLOGY**

ROBERT ALAN MORRIS

THE INFLATION—RESTRICTION THEOREM FOR AMITSUR COHOMOLOGY

ROBERT A. MORRIS

In this paper we develop a generalization of the classical exactness of the inflation—restriction sequence in group cohomology. Our main theorems relate the Amitsur cohomology of algebras to that of subalgebras.

1. **Introduction.** Throughout, R is a commutative ring, unadorned \otimes means tensor product over R , all algebras are commutative, and if S is an R -algebra, S^j denotes the tensor product $S \otimes \cdots \otimes S$, j times. $R\text{-Alg}$ and Ab denote the categories of commutative R -algebras and abelian groups, respectively.

For any R -algebra S there are R -algebra maps $\varepsilon_i^n: S^n \rightarrow S^{n+1}$ given by $\varepsilon_i^n(s_0 \otimes \cdots \otimes s_{n-1}) = s_0 \otimes \cdots \otimes s_{i-1} \otimes 1 \otimes s_i \otimes \cdots \otimes s_{n-1}$, $i = 0, 1, \dots, n + 1$. These are called the (n -dimensional) *co-face maps for S/R* . Generally the superscript will be suppressed. The co-face maps are easily seen to satisfy the co-face relations:

$$\varepsilon_i \varepsilon_j = \varepsilon_{j+1} \varepsilon_i \quad \text{for } i \leq j.$$

If $F: R\text{-Alg} \rightarrow Ab$ is any functor, the Amitsur cochain complex, $C(S/R, F)$, is defined by $C^n(S/R, F) = F(S^{n+1})$, $n = 0, 1, 2, \dots$ [1, 2, 6]. The coboundary operator $d^n: F(S^{n+1}) \rightarrow F(S^{n+2})$ is given by $d^n = \sum_{i=0}^{n+1} (-1)^i F(\varepsilon_i)$. It is a consequence of the co-face relations that a complex results, i.e., that $d^{n+1}d^n = 0$. The homology $\text{Ker } d^n / \text{Im } d^{n-1}$ of this complex is the *Amitsur cohomology of S/R with coefficients in F* , denoted $H^n(S/R, F)$. As usual, $H^0(S/R, F) = \text{Ker } d^0$.

Let $F_1: R\text{-Alg} \rightarrow Ab$ be another functor and let $\eta: F \rightarrow F_1$ be a natural transformation. Then $C(1, \eta) = \eta_{S^{n+1}}: F(S^{n+1}) \rightarrow F_1(S^{n+1})$ is a map of complexes and so induces a map $H^n(1, \eta): H^n(S/R, F) \rightarrow H^n(S/R, F_1)$.

We say a sequence $0 \rightarrow F^\omega F_1 \xrightarrow{\chi} F_2 \rightarrow 0$ is exact if $0 \rightarrow F(A) \xrightarrow{\omega_A} F_1(A) \xrightarrow{\chi_A} F_2(A) \rightarrow 0$ is an exact sequence of abelian groups for each R -algebra A . Indeed the usual long sequence results from a short exact sequence of coefficients.

THEOREM 1.1. [6, p. 47]. *Let $0 \rightarrow F \xrightarrow{\omega} F_1 \xrightarrow{\chi} F_2 \rightarrow 0$ be an exact sequence of functors. Then there are maps $\delta^n(S)$ making*

$$\begin{array}{ccccccc} \dots & \longrightarrow & H^{n-1}(S/R, F_2) & \xrightarrow{\delta^{n-1}(S)} & H^n(S/R, F) & \xrightarrow{H^n(1, \omega)} & H^n(S/R, F_1) \\ & & & & \xrightarrow{H^n(1, \chi)} & H^n(S/R, F_2) & \xrightarrow{\delta^n(S)} & H^{n+1}(S/R, F) & \longrightarrow \dots \end{array}$$

exact and this sequence is natural in S .

This is a standard result, a consequence of the fact that $0 \rightarrow F(S^{n+1}) \xrightarrow{\omega_{S^{n+1}}} F_1(S^{n+1}) \xrightarrow{\chi_{S^{n+1}}} F_2(S^{n+1}) \rightarrow 0$ is a short exact sequence of complexes.

REMARK. The entire discussion thus far in no way depends upon F being defined on all of $R\text{-Alg}$. If A is a full subcategory of $R\text{-Alg}$ containing S^n and T^n and F, F_1 are abelian group valued functors on A , then all the preceding material is still valid.

2. **Inflation-restriction.** By an (R -based) *Grothendieck Topology* T (cf. [7]) we mean a category, $\text{Cat } T$, of commutative R -algebras and a collection, $\text{Cov } T$, of families called *covers* $\{U \rightarrow U_i\}$ of morphisms satisfying axioms dual to those of [3, pp. 1-2]. (In particular, fiber products are replaced by tensor products.) With this convention a *presheaf*, F (of abelian groups) is simply a functor $\text{Cat } T \rightarrow \text{Ab}$ and a presheaf F is a *sheaf* if for every cover $\{U \rightarrow U_i\}$, the induced diagram

$$F(U) \longrightarrow \prod_i F(U_i) \rightrightarrows \prod_{i,j} F(U_i \otimes_U U_j)$$

is an equalizer diagram (equivalently, the natural map $F(U) \rightarrow H_T^n(\{U \rightarrow U_i\}, F)$ [3, I, Sec. 3] is an isomorphism).

REMARK 3.1. If $A \rightarrow B$ is a map in $\text{Cat } T$ and F a presheaf, the complex yielding $H_T^n(\{A \rightarrow B\}, F)$ coincides with the Amitsur complex. The definition of sheaf may be phrased as follows: the natural maps $F(S) \rightarrow H^0(T/S, F)$ and $F(S^2) \rightarrow H^0(T^2/S^2, F)$ are isomorphisms and $F(S^3) \rightarrow H^0(T^3/S^3, F)$ is a monomorphism. Among others which we examine in the next section, the functor which assigns to each R -algebra its multiplicative group of units satisfies this hypothesis if S/R and T/S are faithfully flat. (This follows, for example, from [6, Lemma 3.8].)

Another cohomology theory is defined as follows: the category \mathcal{S} of sheaves on T is abelian with enough injectives [3, Ch. II, Thm. 1.6 (i) and 1.8 (i)]. For any object U in $\text{Cat } T$, the evaluation functor $E_U: \mathcal{S} \rightarrow \text{Ab}$ given by $E_U(F) = F(U)$ is left exact [3, Ch. II, Thm. 1.8 (iii)]. The n th right derived functor of E_U is denoted $H_T^n(U, -)$ and the group $H_T^n(U, F)$ is called the n th Grothendieck cohomology group of U with coefficients in F .

Let $R \xrightarrow{i} S \xrightarrow{j} T$ and \underline{A} be a full subcategory of $R\text{-Alg}$ which is closed under tensor products. Regard S and T as R -algebras and T as S -algebra via i, ij , and j respectively.

The map $H^n(j, 1): H^n(S/R, F) \rightarrow H^n(T/R, F)$ induced by j is called

inflation and denoted *inf*.

Now *i* induces an *R*-algebra map $T^n \rightarrow T \otimes_S \cdots \otimes_S T$ given by $t_1 \otimes \cdots \otimes t_n \rightarrow t_1 \otimes_S \cdots \otimes_S t_n$. This is easily seen to commute with the face maps and so induces a map of complexes and in turn a map of cohomology $H^n(T/R, F) \rightarrow H^n(T/S, F)$, called restriction and denoted *res*.

Note that if $A \rightarrow B$ is a map in \underline{A} , then so is the multiplication map $B \otimes_A B \rightarrow B$ given by $x \otimes y \rightarrow xy$, being simply the composition $B \otimes_A B \rightarrow B \otimes_B B \xrightarrow{\simeq} B$.

Our main theorem is

THEOREM 3.2. (Exactness of the inflation—restriction sequence)
Let X be a Grothendieck topology whose category \underline{A} is such that $\{i\}$ and $\{j\}$ are covers. If F is a sheaf on X, then the inflation—restriction sequence

$$0 \longrightarrow H^n(S/R, F) \xrightarrow{\text{inf}} H^n(T/R, F) \xrightarrow{\text{res}} H^n(T/S, F)$$

*is exact if $n = 1$. Suppose $n \geq 1$ and let Σ be the set of algebras S^i, T^i , or $T \otimes_S \cdots \otimes_S T$ (*i* times), $i \leq n + 1$. If $H_x^j(A, F) = 0$ for all $j < n$ and for all *A* in Σ , then the inflation—restriction sequence is exact for *n*.*

Proof. We induce on *n*.

The case $n = 1$ can be deduced from the spectral sequence of Čech cohomology [3, Ch. II, (3.1)] but a tedious direct argument can be given mimicing the corresponding proof for group cohomology [4, Ch. IV, Sec. 5, Prop. 5]. We illustrate the proof that *inf*. is a monomorphism:

Consider the diagram whose rows are exact since *F* is a sheaf:

$$(1) \quad \begin{array}{ccccc} 0 & \longrightarrow & F(S \otimes_K S) & \xrightarrow{F(j \otimes j)} & F(T \otimes_R T) \\ & & \uparrow F(M_S) & & \uparrow F(M_T) \\ & & \downarrow d^0 & & \downarrow d^0 \\ 0 & \longrightarrow & F(S) & \xrightarrow{F(j)} & F(T) \end{array} \begin{array}{c} \searrow F(\rho) \\ \xrightarrow{F(\varepsilon_0) - F(\varepsilon_1)} \\ \downarrow \\ F(T \otimes_S T) \end{array}$$

with $d^0 = F\varepsilon'_0 - F\varepsilon'_1$, $d^0 = F\varepsilon_0 - F\varepsilon_1$, with ε'_i and ε_i the face maps for S/R and T/R respectively and where $\rho(x \otimes_R y) = x \otimes_S y$, M_S and M_T are the multiplication maps from $S \otimes S$ to S and $T \otimes T$ to T respectively.

The solid arrows of the diagram clearly commute, the commutativity of the square being an example of an *R*-algebra map inducing a map of complexes.

If x in $F(S \otimes S)$ is a one cocycle whose cohomology class gets mapped to zero by inf , then $F(j \otimes j)(x) = (F(\varepsilon_0) - F(\varepsilon_1))(y)$ for some y in $F(T)$. We must show that the cohomology class of x was already zero, i.e., that there is an element z in $F(S)$ such that $(F(\varepsilon'_0) - F(\varepsilon'_1))(z) = x$. By commutativity of solid arrows and exactness of the rows in (1), it clearly suffices to show that $(F(\bar{\varepsilon}_0) - F(\bar{\varepsilon}_1))(y) = 0$. But by the definition of y and the commutativity of (1), this is the same as establishing

$$(2) \quad F(\rho)F(j \otimes j)(x) = 0 .$$

Now in (1) the square with the dotted arrows clearly commutes as does

$$(3) \quad \begin{array}{ccc} T & \xrightarrow{\varepsilon_0} & T \otimes T \\ \varepsilon_1 \downarrow & & \downarrow M_T \\ T \otimes_R T & \xrightarrow{M_T} & T \end{array}$$

so that

$$F(j)F(M_S)(x) = F(M_T)F(j \otimes j)(x) = F(M_T)(F(\varepsilon_0) - F(\varepsilon_1))(y) \quad (\text{by hypothesis on } x)$$

and this is zero by the commutativity of (3). But since F is a sheaf, the map $F(j)$ is monic so we have

$$(4) \quad F(M_S)(x) = 0 .$$

Finally if $\lambda: S \rightarrow T \otimes_s T$ is given by $\lambda(s) = s \otimes 1 = 1 \otimes s$ we clearly have $\rho(j \otimes j) = \lambda M_S$ as maps from $S \otimes_R S \rightarrow T \otimes_s T$ and multiplying (4) by $F(\lambda)$ shows

$$0 = F(\lambda)F(M_S)(x) = F(\rho)F(j \otimes j)(x) .$$

Thus (2) is established, completing the proof that inf is monic.

The remainder of the case $n = 1$ is proved by similar arguments.

For the induction we will make a “dimension shifting” argument. Let $n > 1$. Choose an injective sheaf F^* and a sheaf F' so that

$$0 \longrightarrow F' \longrightarrow F^* \longrightarrow F'' \longrightarrow 0$$

is exact in \mathcal{S} . Now in general this is not exact at each A in \underline{A} so we can not immediately derive a long exact sequence of Amitsur cohomology.

However, since $H^1_X(A, \)$ is the derived functor of “evaluation at A ” we have an exact sequence

$$0 \longrightarrow F(A) \longrightarrow F^*(A) \longrightarrow F'(A) \longrightarrow H^1_x(A, F) .$$

By hypothesis the last term is 0 for all A in Σ .

Consequently we get an exact (up to dimension n) sequence of Amitsur cochain groups

$$0 \longrightarrow C^i(S/R, F) \longrightarrow C^i(S/R, F^*) \longrightarrow C^i(S/R, F') \longrightarrow 0 \quad i \leq n$$

and similar sequences upon replacing S/R by T/R and T/S respectively.

In the usual fashion (cf. Thm. 1.1), these induce exact columns in the following diagram

$$\begin{array}{ccccc}
 & & H^n(S/R, F^*) & \xrightarrow{\text{inf}} & H^n(T/R, F^*) & \xrightarrow{\text{res}} & H^n(T/S, F^*) \\
 & & \uparrow & & \uparrow & & \uparrow \\
 0 & \longrightarrow & H^n(S/R, F) & \xrightarrow{\text{inf}} & H^n(T/R, F) & \xrightarrow{\text{res}} & H^n(T/S, F) \\
 & & \uparrow \delta & & \uparrow \delta & & \uparrow \delta \\
 0 & \longrightarrow & H^{n-1}(S/R, F') & \xrightarrow{\text{inf}} & H^{n-1}(T/R, F') & \xrightarrow{\text{res}} & H^{n-1}(T/S, F') \\
 & & \uparrow & & \uparrow & & \uparrow \\
 & & H^{n-1}(S/R, F^*) & \xrightarrow{\text{inf}} & H^{n-1}(T/R, F^*) & \xrightarrow{\text{res}} & H^{n-1}(T/S, F^*) .
 \end{array}$$

That this diagram commutes is an immediate consequence of definitions of inf and res.

Now F^* is injective and so Čech cohomology of any cover with coefficients in F^* vanishes [3, Prop. 4.3 (iv), p. 40]. But again using Remark 3.1, we conclude that in the above diagram all the Amitsur cohomology with coefficients in F^* also vanishes.

Hence the maps δ are isomorphisms and the desired exactness will follow by induction if we can show that $H^j_x(A, F') = 0$ for all A in Σ and for all $1 \leq j \leq n - 1$. But this is immediate: F^* being injective, the short exact sequence $0 \rightarrow F \rightarrow F^* \rightarrow F' \rightarrow 0$ of sheaves yields $H^j_x(A, F') \cong H^{j+1}_x(A, F) = 0$ for all $1 \leq j < n - 1$ and for all A in Σ .

This completes the proof of the theorem.

The full strength of the definition of sheaf is in fact not needed in case $n = 1$ in the above theorem. All that is required is that the sheaf property hold on $\{S^n \rightarrow T^n\}$, $n = 1, 2, 3$, however in practice the functors of interest which are not sheaves do not even satisfy this.

4. Etale sheaves and group cohomology. In this section we briefly sketch how the classical inflation—restriction theorem for

group cohomology can be recovered from our results by use of the étale topology.

Let G be a finite group, H a normal subgroup and choose fields $k \subseteq L$ with $G = \text{Gal}(L/k)$. Let $N = L^H$ be the fixed field of H and let A be any G module. By a straight-forward modification of the results of I, Sec. 4 and 5 of [7] (cf. "Supplements" in [7]) one can show that there is a topology $T = T_{L/k}$, analogous to the usual étale topology, which has the following properties: (1) Every sheaf on T is additive [9, p. 9]. (2) The category \mathcal{S} of sheaves on T is naturally equivalent to the category of G -modules. This equivalence associates to any sheaf F the module $F(L)$ with g in G acting as $F(g)$. If A is a module and M a subfield of L normal over k (all such subfields are among the objects of $\text{Cat } T$) then the sheaf F_A associated to A has $F_A(M) = A^{M'}$ where M' is the subgroup of G which fixes M .

With these observations one can prove the classical group cohomology theorem:

THEOREM 4.1. [4, Ch. IV, Sec. 5, Prop. 5] *Let G be a finite group, H a normal subgroup and A a G -module. Then*

$$0 \longrightarrow H^n(G/H, A^H) \xrightarrow{\text{inf}} H^n(G, A) \xrightarrow{\text{res}} H^n(H, A)$$

is exact for $n = 1$. If $H^i(H, A) = 0$ for $1 \leq i < n$, then the sequence is exact for n .

Proof. Letting F be the sheaf associated to A one deduces from [5, Thm. 5.4] a commutative diagram

$$\begin{array}{ccccc} 0 \longrightarrow & H^n(N/k, F) & \xrightarrow{\text{inf}} & H^n(L/k, F) & \xrightarrow{\text{res}} & H^n(L/N, F) \\ & \downarrow & & \downarrow & & \downarrow \\ 0 \longrightarrow & H^n(G/H, F(L)^H) & \xrightarrow{\text{inf}} & H^n(G, F(L)) & \xrightarrow{\text{res}} & H^n(H, F(L)) \end{array}$$

with each vertical map an isomorphism. It thus suffices to show the exactness of the upper sequence.

Since N/k , L/k and L/N are Galois, the set Σ of Theorem 3.2 consists of algebras which are the products of copies of N or copies of L .

The arguments of the Supplements, of [7] show that $H_T^n(X, F) \cong H^n(\text{Gal}(L/X), F(L))$ for $X = N$ or L . Since sheaves on T are additive, dimension shifting shows $H_T^n(A \times B, F) \cong H_T^n(A, F) \oplus H_T^n(B, F)$ for any algebras A and B in $\text{Cat } T$. It then follows that the hypotheses of Theorem 3.2 reduce to requiring $H^j(H, A) = 0$ and $H^j(\text{Gal}(L/L), A) = 0$ for $j < n$. The latter is trivial and the former is assumed, completing the proof.

REFERENCES

1. S. A. Amitsur, *Simple Algebras and cohomology groups of arbitrary fields*, Trans. Amer. Math. Soc., **90** (1959), 73-112.
2. ———, *Homology groups and double complexes for arbitrary fields*, J. Math. Soc. Jap., **14** (1962), 1-25.
3. M. Artin, *Grothendieck Topologies*, Seminar Notes, Harvard University, Cambridge, Mass., 1962.
4. J. W. S. Cassels and A. Frohlich, eds., *Algebraic Number Theory* (proceedings of the 1965 Brighton conference), Thompson Book Co., Washington D. C., 1967.
5. S. U. Chase, D. K. Harrison and A. Rosenberg, *Galois theory and galois cohomology of commutative rings*, Amer. Math. Soc. Memoir, no. **52** (1965).
6. S. U. Chase and A. Rosenberg, *Amitsur cohomology and the Brauer group*, Amer. Math. Soc. Memoir, no. **52** (1965).
7. D. Dobbs, *Cohomological Dimension of Fields, Ch. 1 in "Cech cohomology and dimension theory for rings"*, Lecture Notes in Mathematics No. 147, Springer, Berlin, 1970.
8. A. Grothendieck, *Eléments de Geometrie Algebrique*, Ch. IV, Publ. Math. de l'Institut de Hautes Etudes Scientifiques, nos. **20** (1964), **24** (1965), **28** (1966), **32** (1967).
9. S. Lang, *Algebra*, Addison Wesley, 1965.
10. A. Rosenberg and D. Zelinsky, *On Amitsur's complex*, Trans. Math. Soc., **97** (1960), 327-356.
11. J. P. Serre, *Cohomologie Galoisienne*, Springer, Berlin, 1965.

Received August 6, 1970 and in revised form September 10, 1971. The results in this paper are contained in this author's Ph. D. thesis written under the supervision of Prof. Alex Rosenberg at Cornell University. The research was supported in part by the National Science Foundation, NSF GP9345 and GU3171.

STATE UNIVERSITY OF NEW YORK AT ALBANY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics

Vol. 41, No. 3

BadMonth, 1972

George E. Andrews, <i>Two theorems of Gauss and allied identities proved arithmetically</i>	563
Stefan Bergman, <i>On pseudo-conformal mappings of circular domains</i>	579
Beverly L. Brechner, <i>On the non-monotony of dimension</i>	587
Richard Anthony Brualdi and John H. Mason, <i>Transversal matroids and Hall's theorem</i>	601
Philip Throop Church and James Timourian, <i>Differentiable maps with 0-dimensional critical set. I</i>	615
John H. E. Cohn, <i>Squares in some recurrent sequences</i>	631
Robert S. Cunningham, Edgar Andrews Rutter and Darrell R. Turnidge, <i>Rings of quotients of endomorphism rings of projective modules</i>	647
Eldon Dyer and S. Eilenberg, <i>An adjunction theorem for locally equiconnected spaces</i>	669
Michael W. Evans, <i>On commutative P. P. rings</i>	687
Ronald Lewis Graham, Hans Sylvain Witsenhausen and Hans Zassenhaus, <i>On tightest packings in the Minkowski plane</i>	699
Stanley P. Gudder, <i>Partial algebraic structures associated with orthomodular posets</i>	717
Karl Edwin Gustafson and Gunter Lumer, <i>Multiplicative perturbation of semigroup generators</i>	731
Kurt Kreith and Curtis Clyde Travis, Jr., <i>Oscillation criteria for selfadjoint elliptic equations</i>	743
Lawrence Louis Larmore, <i>Twisted cohomology theories and the single obstruction to lifting</i>	755
Jorge Martinez, <i>Tensor products of partially ordered groups</i>	771
Robert Alan Morris, <i>The inflation-restriction theorem for Amitsur cohomology</i>	791
Leo Sario and Cecilia Wang, <i>The class of (p, q)-biharmonic functions</i>	799
Manda Butchi Suryanarayana, <i>On multidimensional integral equations of Volterra type</i>	809
Kok Keong Tan, <i>Fixed point theorems for nonexpansive mappings</i>	829