

Pacific Journal of Mathematics

**RELICH DENSITIES AND AN APPLICATION TO
UNCONDITIONALLY NONOSCILLATORY ELLIPTIC
EQUATIONS**

JOHN PIEPENBRINK

RELLICH DENSITIES AND AN APPLICATION TO UNCONDITIONALLY NONOSCILLATORY ELLIPTIC EQUATIONS

JOHN PIEPENBRINK

Sufficient conditions for embeddings between weighted Sobolev spaces to be compact are derived. These theorems are generalizations of the well known selection principle of Rellich. These results are then applied to the study of the oscillational properties of self-adjoint second order elliptic equations. In addition to reproving some results of Headley and Swanson, new nonoscillation criteria are furnished for these equations.

1. Introduction. Let Ω be a domain, bounded or unbounded, in Euclidean n -space E^n , $p(x)$ a positive measurable function, $x = (x_1, \dots, x_n)$, and $a(x)$ a symmetric matrix with measurable entries such that the smallest eigenvalue of $a(x)$ for each x in Ω positive. Define the weighted strong Sobolev spaces $H_\Omega(p)$ and $H_\Omega(p, a)$ as the closure of the sets of functions u, C^1 on Ω for which the integrals

$$(1.1) \quad \int_{\Omega} p(x) [u(x)]^2 dx$$

$$(1.2) \quad \int_{\Omega} \{p(x) [u(x)]^2 + \sum a_{ij}(x) u_i(x) u_j(x)\} dx$$

are finite. The closures are taken with respect to norms given by (1.1) and (1.2). The weighted weak Sobolev spaces $W_\Omega(p)$ and $W_\Omega(p, a)$ consist of functions u with (1.1) or (1.2) respectively being finite. Here $u_i(x)$ is the distributional derivative $\partial u / \partial x_i$.

We will say that the pair (p, a) has the *strong Rellich compactness property* if the inclusion map $H_\Omega(p, a) \rightarrow H_\Omega(p)$ is compact. This means that each sequence in $H_\Omega(p, a)$ which is uniformly bounded in its norm has a subsequence which is convergent in the norm for $H_\Omega(p)$. The classical Rellich selection principle states that if Ω is bounded and smooth, then $(1, I)$ has the strong Rellich compactness property where I is the identity matrix. The *weak Rellich compactness property* is defined analogously with $W_\Omega(p)$, $W_\Omega(p, a)$ taking the place of $H_\Omega(p)$, $H_\Omega(p, a)$.

This paper investigates the case where $\Omega = E^n$, $n \geq 2$. The arguments however apply equally well to quasi-conical domains, i.e. domains which contain a cone $\{x \mid x \cdot v \geq \alpha |x|\}$, where v is some unit vector, and α is a positive constant. Theorem 3.1 and 3.2 of § 3 provide sufficient conditions for (p, a) to have either the strong or

weak Rellich compactness property. Theorem 3.2 is based on a Sobolev type lemma communicated to the author by N. Meyers. A proof of this lemma is included as an appendix. Theorem 3.3 gives a simple condition on $p(x)$ in case $a(x)$ is uniformly definite on E^n which ensures that the inclusion map $H_{E^n}(p, a) \cap L^2(E^n) \rightarrow H_{E^n}(p)$ is compact. This weaker result is still sufficient for the application in § 5. In § 4 we give a necessary condition on p when a is uniformly definite. An example shows that the sufficient conditions of Theorems 3.2 and 3.3 are the best of their kind in this case.

In § 5 we apply the preceding results to the determination of the oscillatory properties of the elliptic equation

$$(1.3) \quad Lu = \sum_{i,j=1}^n \frac{\partial}{\partial x_i} (a_{ij}(x)u_j) + p(x)u = 0.$$

We say that (1.3) is *nonoscillatory* if there is a positive constant R such that for each bounded domain N with smooth boundary exterior to the sphere $\{x \mid |x| \leq R\}$, the Dirichlet problem

$$\begin{aligned} Lu &= 0 & \text{in } N \\ u &= 0 & \text{on } \partial N \end{aligned}$$

has nontrivial solution. (1.3) is said to be *unconditionally nonoscillatory* if for each positive λ , (1.3) with $p(x)$ replaced by $\lambda p(x)$ is nonoscillatory.

Theorem 5.1, which asserts that if (p, a) has the Rellich compactness property then (1.3) is unconditionally nonoscillatory, combined with the results of § 3 yields results which differ from the recent nonoscillation theorems of Swanson and Headley, [1] and [2], in two respects. Our conditions apply directly to the coefficients $a_{ij}(x)$ and $p(x)$, rather than to pointwise majorants. Furthermore our results are not based on the oscillation theory for ordinary differential equations. There is, however, some overlap between the results of [1] and [2] and ours, which will be pointed out later.

2. The case of a bounded domain. Our results for the unbounded domain E^n will follow from a process of Cantor diagonalization over compact subdomains. Therefore we will need conditions guaranteeing that (p, a) has a Rellich compactness property for a bounded domain Ω with smooth boundary $\partial\Omega$. The first result in this direction follows from an imbedding theorem for weighted Sobolev spaces due to Stampachia and Murthy [7], which in turn followed easily from the corresponding result for classical Sobolev spaces, see Sobolev [5]. A special case of the Stampachia-Murthy Theorem is stated as a lemma:

LEMMA 2.1. *Let Ω be a bounded domain in E^n with smooth boundary. Suppose $\lambda(x)$ is the smallest eigenvalue of $a(x)$ and $\lambda^{-1}(x)$ and $p^{-1}(x)$ are in $L^t(\Omega)$ for some $t \geq n$. If $1 + 1/t < 2 < n(1 + 1/t)$, then the embedding $W_\Omega(p, a) \rightarrow L^{q-\varepsilon}(\Omega)$ is compact, where q is defined by*

$$(2.1) \quad \frac{1}{q} = \frac{1}{2} \left(1 + \frac{1}{t} \right) - \frac{1}{n} .$$

From Lemma 2.1 follows our first result.

THEOREM 2.1. *Suppose the conditions of Lemma 2.1 hold. If in addition $p(x)$ is in $L^s(\Omega)$ where $s = (q - \varepsilon)(q - \varepsilon - 2)^{-1}$, then (p, a) has the weak Rellich compactness property on Ω .*

By Hölder's inequality

$$\left[\int_\Omega p(x) |u(x)|^2 dx \right]^{1/2} \leq \left[\left(\int_\Omega |u(x)|^{q-\varepsilon} dx \right)^{2/(q-\varepsilon)} \left(\int_\Omega |p(x)|^s dx \right)^{1/s} \right]^{1/2} ,$$

or

$$\left[\int_\Omega p(x) |u(x)|^2 dx \right]^{1/2} \leq c \|u\|_{q-\varepsilon} ,$$

where $\|u\|_{q-\varepsilon}$ is the norm of u in $L^{q-\varepsilon}(\Omega)$. Thus the imbedding $L^{q-\varepsilon}(\Omega) \rightarrow W_\Omega(p)$ is continuous. Since by Lemma 2.1 the imbedding $W_\Omega(p, a) \rightarrow L^{q-\varepsilon}(\Omega)$ is compact, and since the composition of a compact map and a continuous map is compact we have the desired conclusion.

A second compactness theorem can be obtained independently of the Sobolev imbedding theorems. Yet despite its apparent simplicity it permits $p(x)$ which are inadmissible in Lemma 2.1. This generality is obtained at the expense of restricting ourselves to the strong spaces.

THEOREM 2.2. *Let Ω be bounded and convex and $p(x)$ be non-negative and measurable. If $p(x)$ is bounded above and has a positive lower bound on some open subset of Ω , then (p, I) has the strong Rellich compactness property.*

Proof. Let us assume that B is a bounded subset of $H_\Omega(p, I)$. It suffices to assume that $B \subset C^1(\Omega)$. Pick x_0 in Ω such that $p(x) \geq p_0$ for $|x - x_0| \leq \varepsilon_1$. Thus if u is in B , ∇u is in $L^2(\Omega)$ and u is in $L^2(S(x_0, \varepsilon_1))$, where $S(x_0, \varepsilon_1) = \{x \mid |x - x_0| \leq \varepsilon_1\}$. Now fix ε_0 , $0 < \varepsilon_0 < \varepsilon_1$. For any ε such that $\varepsilon_0 \leq \varepsilon \leq \varepsilon_1$ set $u_\varepsilon(x) = u(x_0 + \varepsilon(x - x_0)/(|x - x_0|))$.

Introduce spherical coordinates (r, θ) , $r = |x - x_0|$ and

$$\theta = \frac{x - x_0}{|x - x_0|}.$$

Then

$$\begin{aligned} \int_{\varepsilon_0 < |x - x_0| < \varepsilon_1} |u(x)|^2 dx &= \int_{\varepsilon_0}^{\varepsilon_1} r^{n-1} \int_{\theta} |u(x_0 + r\theta)|^2 d\theta dr \\ &\geq \frac{\varepsilon_1^n - \varepsilon_0^n}{n} \min_{\varepsilon_0 < r < \varepsilon_1} \int_{\theta} |u(x_0 + r\theta)|^2 d\theta. \end{aligned}$$

So for u in B there is an ε , $\varepsilon_0 \leq \varepsilon \leq \varepsilon_1$ with

$$(2.2) \quad \int_{\theta} |u(x_0 + \varepsilon\theta)|^2 d\theta \leq C_1,$$

where C_1 is a constant independent of the particular u in B . Now if R is the diameter of Ω , also by (2.2)

$$\begin{aligned} \int_{|x - x_0| \geq \varepsilon} |u_\varepsilon(x)|^2 dx &\leq \int_{\varepsilon}^R r^{n-1} \int_{\theta} |u(x_0 + \varepsilon\theta)|^2 d\theta dr \\ &\leq C_1 \frac{R^n - \varepsilon^n}{n} < C_1 \frac{R^n}{n} \equiv C_2. \end{aligned}$$

So for each u in B there is an $\varepsilon = \varepsilon(u)$, $\varepsilon_0 \leq \varepsilon \leq \varepsilon_1$, such that

$$(2.3) \quad \int_{|x - x_0| \geq \varepsilon_1} |u_\varepsilon(x)|^2 dx \leq C_2,$$

and C_2 is independent of u and ε .

Also if $|x - x_0| \geq \varepsilon$

$$u(x) - u\left(x_0 + \varepsilon \frac{x - x_0}{|x - x_0|}\right) = \int_{\varepsilon}^{|x - x_0|} \nabla u\left(x_0 + s \frac{x - x_0}{|x - x_0|}\right) \cdot \frac{x - x_0}{|x - x_0|} ds.$$

Letting $t = s/(|x - x_0|)$ we see that

$$|u(x) - u_\varepsilon(x)| \leq |x - x_0| \int_{\varepsilon/|x - x_0|}^1 |\nabla u(x_0 + t(x - x_0))| dt.$$

Now square both sides and use the Schwartz inequality, integrating over $|x - x_0| \geq \varepsilon_1$ to derive

$$\int_{|x - x_0| \geq \varepsilon_1} |u(x) - u_\varepsilon(x)|^2 dx \leq R^2 \int_{|x - x_0| \geq \varepsilon_1} \int_{\varepsilon/R}^1 |\nabla u(x_0 + t(x - x_0))|^2 dt dx.$$

We reverse the integrals on the right and set $y = x_0 + t(x - x_0)$, obtaining

$$\int_{|x - x_0| \geq \varepsilon_1} |u(x) - u_\varepsilon(x)|^2 dx \leq R^2 \int_{\varepsilon_0/R}^1 \frac{1}{t^n} \int_{\Omega} |\nabla u(y)|^2 dy dt,$$

or

$$(2.4) \quad \int_{|x-x_0| \geq \varepsilon_1} |u(x) - u_\varepsilon(x)|^2 dx \leq C_3 \int_\Omega |\nabla u(y)|^2 dy,$$

where C_3 only depends on R and ε_0 . From (2.3) and (2.4) we finally see that

$$\begin{aligned} \int_\Omega |u(x)|^2 dx &= \int_{|x-x_0| \leq \varepsilon_1} |u(x)|^2 dx + \int_{|x-x_0| \geq \varepsilon_1} |u(x)|^2 dx \\ &\leq C_4 + 2 \int_{|x-x_0| \geq \varepsilon_1} |u(x) - u_\varepsilon(x)|^2 dx + 2 \int_{|x-x_0| \geq \varepsilon} |u_\varepsilon(x)|^2 dx \\ &\leq C_4 + 2C_3 \int_\Omega |\nabla u|^2 dy + 2C_2. \end{aligned}$$

Thus there is a C independent of u such that

$$(2.5) \quad \int_\Omega |u(x)|^2 dx \leq C$$

for all u in B .

Now

$$|u(x+h) - u(x)| \leq \int_0^{|h|} \left| \nabla u \left(x + t \frac{h}{|h|} \right) \right| dt$$

or

$$|u(x+h) - u(x)|^2 \leq |h| \int_0^{|h|} \left| \nabla u \left(x + t \frac{h}{|h|} \right) \right|^2 dt.$$

Integrate both sides with respect to x and interchange the order of integration on the right to obtain

$$(2.6) \quad \int_\Omega |u(x+h) - u(x)|^2 dx \leq |h|^2 \int_\Omega |\nabla u(x)|^2 dx \leq C' |h|^2.$$

By a well known theorem of Bochner, see [6] p. 38, (2.5) and (2.6) imply that B is compact in $L^2(\Omega)$ and since $p(x)$ is bounded above, B is compact in $W_\Omega(p)$.

We see that the proof of Theorem 2.2 has yielded the following embedding theorem.

THEOREM 2.3. *Let $p(x)$ and Ω be as in Theorem 2.2. Then $H_\Omega(p, I)$ is embedded in $L^2(\Omega)$ by a compact mapping.*

It should be observed that in the applications to differential equations in § 5 only smooth functions are used. Also in many applications $a(x)$ will be uniformly definite and $p(x)$ bounded from below

and above by positive constants on bounded subdomains Ω of E^n . In this case, where Theorem 2.1 applies anyway, the weak Rellich compactness of (p, a) over Ω follows from the classical Rellich selection principle referred to in § 1.

3. $\Omega = E^n$. Through this and the remaining sections of this paper the hypothesis of Theorem 2.1 or Theorem 2.2 are assumed to hold on each bounded subdomain of E^n .

Our first compactness result gives conditions on $p(x)$ sufficient for the mapping $H_{E^n}(p, I) \rightarrow H_{E^n}(p)$ to be compact. So it may be assumed that the hypothesis either of Theorem 2.1 or 2.2 hold on bounded subdomains. Spherical coordinates (r, θ) , $r = |x|$, $\theta = x/|x|$ are introduced.

THEOREM 3.1. *If $p(x)$ satisfies the local conditions of Theorem 2.1 or 2.2, is continuous and in addition*

$$(3.1) \quad \lim_{a \rightarrow \infty} \left[\sup_{\theta} \int_a^{\infty} r^n p(r\theta) dr \right] = 0, \quad \text{and} \quad \inf_{\theta} \int_1^a r^{n-1} p(r\theta) dr > 0$$

then (p, I) has the strong Rellich compactness property.

Proof. The theorem is proved if we can establish the inequality

$$(3.2) \quad \int_{E^n} p(x) |u(x)|^2 dx \leq \gamma \left[\int_{|x| < a} p(x) |u(x)|^2 dx + \varepsilon(a) \int_{E^n} |\nabla u(x)|^2 dx \right],$$

where $\varepsilon(a) \rightarrow 0$ as $a \rightarrow \infty$, $\varepsilon(a)$ and γ are independent of u in $H_{E^n}(p, I)$. For if $\{u_n\}$ is a bounded sequence in $H_{E^n}(p, I)$, by the results of the preceding section and Cantor diagonalization we can select a subsequence $\{u_{n_k}\}$ which is Cauchy in $H_{\Omega}(p)$ for each bounded subdomain Ω of E^n . But then (3.2) would show that $\{u_{n_k}\}$ was Cauchy in $H_{E^n}(\Omega)$, and hence convergent. This would establish the desired conclusion.

Thus we need to prove (3.2) for u in $H_{E^n}(p, I)$. Without loss of generality assume that u is in $C^1(E^n)$. Fix θ and let $1 \leq s < t$. Then

$$u(t\theta) = u(s\theta) + \int_s^t \nabla u(s\theta + (\tau - s)\theta) \cdot \theta d\tau$$

$$u^2(t\theta) \leq 2u^2(s\theta) + 2(t - s) \int_1^{\infty} |\nabla u(\tau\theta)|^2 \tau^{n-1} d\tau.$$

Multiply both sides of the inequality by $s^{n-1}p(s\theta)$ and integrate with respect to s from $s = 1$ to $s = a$, then divide by $\int_1^a s^{n-1}p(s\theta)ds$ to derive

$$u^2(t\theta) \leq \frac{2}{\int_1^a s^{n-1} p(s\theta) ds} \left[\int_1^a s^{n-1} p(s\theta) |u(s\theta)|^2 ds + \int_1^a (t-s) s^{n-1} p(s\theta) ds \right. \\ \left. \times \int_1^\infty |\nabla u(\tau\theta)|^2 \tau^{n-1} d\tau \right].$$

Now multiply both sides by $t^{n-1} p(t\theta)$ and integrate with respect to t from $t = a$ to $t = \infty$, to find that

$$(3.3) \quad \int_a^\infty t^{n-1} p(t\theta) |u(t\theta)|^2 dt \leq \frac{2 \int_0^a s^{n-1} p(s\theta) |u(s\theta)|^2 ds}{\int_1^a s^{n-1} p(s\theta) ds} \int_a^\infty t^{n-1} p(t\theta) dt \\ + \frac{2 \int_a^\infty \int_0^a t^{n-1} (t-s) s^{n-1} p(t\theta) p(s\theta) ds dt}{\int_1^a s^{n-1} p(s\theta) ds} \\ \times \int_0^\infty |\nabla u(\tau\theta)|^2 \tau^{n-1} d\tau.$$

By reversing the order of integrations

$$\int_a^\infty \int_0^a t^{n-1} (t-s) s^{n-1} p(t\theta) p(s\theta) ds dt = \left[\int_0^a s^{n-1} p(s\theta) ds \right] \left[\int_a^\infty t^n p(t\theta) dt \right] \\ - \left[\int_a^\infty t^{n-1} p(t\theta) dt \right] \left[\int_0^a s^n p(s\theta) ds \right].$$

Thus condition (3.1) shows that if $\varepsilon(a)$ is defined by

$$\varepsilon(a) = \sup_\theta \int_a^\infty \int_0^a t^{n-1} (t-s) s^{n-1} p(t\theta) p(s\theta) ds dt,$$

then

$$\varepsilon(a) \longrightarrow 0 \quad \text{as } a \longrightarrow \infty.$$

Define

$$\gamma \geq 2 \left(\inf_\theta \int_1^a s^{n-1} p(s\theta) d\theta \right)^{-1} \quad \text{for all } a \geq A.$$

Then (3.3) implies

$$\int_a^\infty t^{n-1} p(t\theta) |u(t\theta)|^2 dt \leq \gamma \left[\int_0^a s^{n-1} p(s\theta) |u(s\theta)|^2 ds \right. \\ \left. + \varepsilon(a) \int_0^\infty |\nabla u(\tau\theta)|^2 \tau^{n-1} d\tau \right].$$

Now if this is integrated over all θ , (3.2) results.

It is clear from the proof that the same conclusion follows if the condition

$$\inf_{\theta} \int_1^a r^{n-1} p(r\theta) dr > 0$$

is replaced by

$$\inf_{\theta} \int_{\delta}^a r^{n-1} p(r\theta) dr > 0,$$

where δ is a positive constant.

Our next theorem depends on a lemma related to a Sobolev type embedding theorem, see [3]. It differs from such embedding results in that instead of asserting that a function belongs to a certain Lebesgue class, it states that the function minus a suitable constant is in the Lebesgue space. This lemma was communicated to the author by N. Meyers. The proof is given in an appendix.

LEMMA 3.1. *Let r be a number with $1 < r < n$ and define r^* by the expression $(r^*)^{-1} = r^{-1} - n^{-1}$. Then if $u(x)$ is a function on E^n whose gradient is in $L^r(E^n)$, there is a function v in $L^{r^*}(E^n)$ and a constant k such that*

$$u(x) = v(x) + k.$$

Furthermore the inequality

$$\|v\|_{r^*} \leq c \|\nabla u\|_r$$

holds, where c is a constant depending only on n and r .

Now define a function $q(x)$ such that

$$(3.4) \quad \sum_{i,j=1}^n a_{ij}(x) y_i y_j \geq q(x) |y|^2$$

for all x and y in E^n . Theorem 3.2 gives conditions on p and q guaranteeing that (p, a) has the Rellich compactness property.

THEOREM 3.2. *Let q be such that (3.4) holds and α a constant such that $(2n)(n+2)^{-1} < \alpha \leq 2$. Then if*

- (i) p is in $L^{\alpha/(\alpha-2)}(E^n)$, and
- (ii) $1/q$ is in $L^{\alpha(2-\alpha)^{-1}}(E^n)$

the pair (p, a) has the weak Rellich compactness property.

Proof. By (3.4) we may assume without loss of generality that $a(x) = q(x) I$. Let B be a bounded set in $W(p, a)$, so that there is a constant K with

$$\int p(x) u^2 dx + \int q(x) |\nabla u|^2 dx \leq K$$

for each u in B . We must find a sequence in B which is convergent in the norm for $W(p)$.

By Hölder's inequality if $\alpha < 2$

$$\begin{aligned} \int |\nabla u|^\alpha dx &= \int [q(x)^{-1}]^{\alpha/2} |q^{1/2}(x)\nabla u|^\alpha dx \\ &\leq \left[\int [q(x)^{-1}]^{\alpha/(2-\alpha)} dx \right]^{(2-\alpha)/2} \left[\int q(x) |\nabla u|^2 dx \right]^{\alpha/2}. \end{aligned}$$

Thus for each u in B

$$(3.5) \quad \int |\nabla u|^\alpha dx \leq C_1 K$$

where C_1 depends only on $q(x)$. Clearly (3.5) also holds when $\alpha = 2$. By Lemma 3.1 and (3.5) for each u in B there is a v_u in L^{α^*} and a constant k_u such that

$$(3.6) \quad u = v_u + k_u,$$

and

$$(3.7) \quad \|v_u\|_{\alpha^*} \leq C_2,$$

where C_2 does not depend on u .

Now we claim there is a sequence $\{u_n\}$ from B such that the v_{u_n} associated with u_n by (3.6) converge in $W(p)$. For by Hölder's inequality

$$\int_{|x|>R} p(x)v_u^2 dx \leq \left(\int_{|x|>R} |v_u(x)|^{\alpha^*} dx \right)^{2/\alpha^*} \left(\int_{|x|>R} [p(x)]^{\alpha^*/(\alpha^*-2)} dx \right)^{(\alpha^*-2)/\alpha^*}.$$

So by hypothesis (i) and (3.7) the estimate

$$(3.8) \quad \int_{|x|>R} p(x)v_u^2 dx \leq C_2^2 \varepsilon(R)$$

holds, where $\varepsilon(R) \rightarrow 0$ as $R \rightarrow \infty$. By the results of § 2 and Cantor diagonalization there is a sequence $\{v_{u_n}\}$ which converges in the $W(p)$ norm on each compact subset of E^n . But then the estimate (3.8) show that this sequence converges globally in $W(p)$.

If $q(x)$ is not in $L^1(E^n)$, the fact that $k = 0$ in (3.6) follows. Then we are done since $u_n = v_{u_n}$. But if $p(x)$ is in $L^1(E^n)$, then clearly from (3.6) and (3.8) with $R = 0$ we have the inequality

$$|k| = \text{const.} \quad \|k\|_{W(p)} \leq \text{const.} \quad (\|u\|_{W(p)} + \|v_u\|_{W(p)}) \leq C,$$

where C is independent of u in B . Thus $\{k_n\}$ is a bounded sequence in R^1 and must have a convergent subsequence in R^1 . But this subsequence also converges in $W(p)$. Hence $u_n = v_{u_n} + k_{u_n}$ has a con-

vergent subsequence in $W(p)$. This completes the proof.

If $a(x)$ is uniformly definite then we may choose $q(x)$ to be a positive constant. Then we may take $\alpha = 2$ in Theorem 3.2. As a special case of Theorem 3.2 we have the following corollary.

COROLLARY 3.2. *If $a(x)$ is uniformly definite and $p(x)$ is in $L^{n/2}(E^n)$ for $n \geq 3$, then (p, a) has the weak Rellich compactness property.*

In the case where $a(x)$ is uniformly definite a simple bound on the growth of $p(x)$ as $x \rightarrow \infty$ suffices to ensure a weaker compactness property of the pair (p, a) .

THEOREM 3.3. *If $a(x)$ is uniformly definite and $p(x) = o(|x|^{-2})$ as $x \rightarrow \infty$, then the inclusion map $H(p, a) \cap L^2(E^n) \rightarrow H(p)$ is compact.*

Proof. Without loss of generality assume that $a(x) = I$ and let B be a bounded set in $H(p, I) \cap L^2(E^n)$. For any function in $L^2(E^n)$ with its gradient in $L^2(E^n)$ the inequality

$$(3.9) \quad \int |x|^2 u^{-2} dx \leq (n-2)^{-2} \int |\nabla u|^2 dx$$

holds, see [4] where (3.9) is proved in greater generality. But then because of the growth condition on $p(x)$ we have the estimate

$$(3.10) \quad \int_{|x|>R} p(x)u^2 dx \leq \varepsilon(R) \int |\nabla u|^2 dx,$$

where $\varepsilon(R) \rightarrow 0$ as $R \rightarrow \infty$. As before we pick a sequence $\{u_n\}$ from B which converges in $H(p)$ on each compact subset of E^n . But (3.10) then shows that $\{u_n\}$ converges globally in $H(p)$, and the proof is complete.

The following section contains an example to show that in the case $a(x) = I$ the conditions $p(x) \in L^{n/2}(E^n)$ and $p(x) = o(|x|^{-2})$ of Corollary 3.2 and Theorem 3.2 are the best of their kind. By this is meant that $p(x) \in L^{n/2+\varepsilon}(E^n)$ and $p(x) = O(|x|^{-2})$ will not be sufficient.

4. A necessary condition. In this section we limit ourselves to the case $a(x) = I$. In this case Theorem 4.1 provides a necessary condition for (p, I) to have the Rellich compactness property.

THEOREM 4.1. *Suppose (p, I) has the Rellich compactness property.*

If $n = 2$, then $p(x)$ must be in $L^1(E^2)$. If $n > 2$ then necessarily

$$(4.1) \quad \lim_{t \rightarrow \infty} t^{2-n} \int_{|x| < t} p(x) dx = 0 .$$

Proof. For positive numbers a and k consider the piecewise C^1 function $u(x; a, k)$ defined to be identically zero for $|x| \leq a$ and $|x| \geq a + k + 1$, to be one for $|x| = a + 1$, and to be linear in $|x|$ for $a \leq |x| \leq a + 1$ and $a + 1 \leq |x| \leq a + k + 1$. If $Q[u]$ represents the quotient $\left(\int |\nabla u|^2 dx \right) \left(\int p u^2 dx \right)^{-1}$, then

$$Q[u(x; a, k)] = \frac{\text{const.} [(a + 1)^n - a^n + k^{-2}((a + k + 1)^n - (a + 1)^n)]}{\int p u^2 dx} .$$

By obvious estimation of the denominator and multiplication of numerator and denominator by k^{2-n} we derive the estimate

$$(4.2) \quad Q[u(x; a, k)] \leq \frac{\text{const.} [k^{2-n}((a + 1)^n - a^n) + (1 + k^{-1}(a + 1))^n - (k^{-1}(a + 1))^n]}{k^{2-n} \int_{a+1 < |x| < a+1+1/2k} p(x) dx} .$$

If $n = 2$ and $\int p(x) dx = \infty$, (4.2) shows that for any a there is a k such that $Q[u(x; a, k)] \leq 1$. If $n > 2$ and (4.1) does not hold there is a $\delta > 0$ and a sequence $\{t_m\}$ with $t_m \rightarrow \infty$ such that

$$(4.3) \quad t_m^{2-n} \int_{|x| < t_m} p(x) dx \geq 2\delta$$

for all m . Now if $k = k_m \equiv 2(t_m - a - 1)$, the denominator of the right side of (4.2) becomes essentially

$$\begin{aligned} & (t_m - (a + 1))^{2-n} \left[\int_{0 < |x| < t_m} p(x) dx - \int_{0 < |x| < a+1} p(x) dx \right] \\ &= \left(1 - \frac{a + 1}{t_m} \right)^{2-n} \left[t_m^{2-n} \int_{0 < |x| < t_m} p(x) dx - t_m^{2-n} \int_{0 < |x| < a+1} p(x) dx \right] \\ &\geq \delta \end{aligned}$$

for all large m . Thus in either case there is a constant K such that for any given a , there is a k such that $Q[u(x; a, k)] \leq K$.

Thus if either condition of our theorem is violated we can pick a sequence $\{u_n\}$ of piecewise C^1 functions with disjoint supports such that $Q[u_n] \leq K$. But if we set $v_n = \left[\int p u_n^2 dx \right]^{-1/2} u_n$, then

$$\int p v_n^2 dx + \int |\nabla v_n|^2 dx = 1 + Q[u_n] \leq 1 + K$$

for each n . But $\{v_n\}$ clearly cannot have any convergent subsequences since

$$\int p(v_n - v_m)^2 dx = 2$$

if $n \neq m$. Thus if either condition in our theorem is violated, the pair (p, I) will not have the Rellich compactness property.

As an example consider $p(x) = |x|^{-2}$. By Theorem 4.1 (p, I) does not have the Rellich compactness property. Yet $p(x)$ is in $L^{n/2+\varepsilon}(E^n)$ for each $\varepsilon > 0$ and $p(x) = 0(|x|^{-2})$ as $x \rightarrow \infty$.

5. Unconditionally nonoscillatory equations. Let $p(x)$ and $a(x)$ be as in § 1 and in addition let them be C^∞ . We shall show that if (p, a) has the Rellich compactness property then for each $\lambda > 0$ the equation

$$(5.1) \quad \sum_{i,j=1}^n \frac{\partial}{\partial x_i} (a_{ij} u_j) + \lambda p u = 0$$

is nonoscillatory. We say that in this case the equation

$$(5.2) \quad \sum_{i,j=1}^n \frac{\partial}{\partial x_i} (a_{ij} u_j) + p u = 0$$

is unconditionally nonoscillatory.

THEOREM 5.1. *Suppose that (p, a) has either the weak or strong Rellich compactness property. Then (5.2) is unconditionally nonoscillatory.*

Proof. Define $C_0^\infty(|x| > R)$ to be the space of C^∞ functions with compact supports contained in the set of x with $|x| > R$. Define the function $f(R)$ by

$$(5.3) \quad f(R) = \inf \{Q[\varphi] \mid \varphi \text{ is in } C_0^\infty(|x| > R)\},$$

where

$$Q[\varphi] = \left[\int \sum_{i,j=1}^n a_{ij}(x) \varphi_i \varphi_j dx \right] \left[\int p(x) \varphi^2 dx \right]^{-1}.$$

Now $f(R)$ is a continuous function of R and furthermore $\lim_{R \rightarrow \infty} f(R) = \infty$. For if $\lim_{R \rightarrow \infty} f(R) = L < \infty$, then we could select a sequence of functions in C_0^∞ , say $\{\varphi_m\}$, with disjoint supports such that

$$\lim_{m \rightarrow \infty} Q[\varphi_m] = L.$$

But this contradicts the fact that (p, a) has the Rellich compactness property since if

$$\psi_m = \left(\int p \varphi_m dx \right)^{-1/2} \varphi_m,$$

then

$$\int p \psi_m^2 dx + \int \sum_{i,j} \psi_{m,i} \psi_{m,j} dx = 1 + Q[\varphi_m] \leq K$$

for all m , and

$$\int p(\psi_m - \psi_l)^2 dx = 2$$

if $m \neq l$. So on the one hand $\{\psi_m\}$ forms a bounded set in $W(p, a)$ or $H(p, a)$ and on the other contains no subsequences convergent in $W(p)$ or $H(p)$.

Now if $\lambda > 0$ pick R so that $f(R) > \lambda$. If N is any bounded smooth subdomain of $\{x \mid |x| < R\}$, the first eigenvalue σ_1 of the problem

$$\sum_{i,j=1}^n \frac{\partial a}{\partial x_0} (a_{ij}(x)u_j) + \sigma p(x)u = 0, \quad \text{in } N$$

$$u = 0 \quad \text{on } \partial N$$

is greater than $f(R)$ and so $\sigma_1 > \lambda$. Thus N cannot be a nodal domain for the equation

$$(5.4) \quad \sum_{i,j=1}^n \frac{\partial}{\partial x_i} (a_{ij}(x)u_j) + \lambda p(x)u = 0.$$

So (5.4) is nonoscillatory, and (5.2) is unconditionally nonoscillatory.

COROLLARY 5.1. *If p and q satisfy the conditions of Theorem 3.2, then (5.2) is unconditionally nonoscillatory.*

COROLLARY 5.2. *If $a(x)$ is uniformly definite on E^n , $n \geq 3$, either of the following conditions is sufficient for (5.2) to be unconditionally nonoscillatory:*

- (i) $p(x)$ is in $L^{n/2}(E^n)$,
- (ii) $p(x) = o(|x|^{-2})$ as $|x| \rightarrow \infty$.

Condition (ii) of Corollary (5.2) is also obtained as a special case of a theorem of Headley and Swanson, see [2] Theorem 5. However, Corollary 5.1 and condition (i) of Corollary 5.2 seem to be new.

Theorem 3.1 coupled with Theorem 5.1 leads to the next corollary.

COROLLARY 5.3. *Suppose $a(x)$ is uniformly definite on E^n and*

$p(x)$ is positive and continuous for each x . If

$$\lim_{a \rightarrow \infty} \left[\sup_{\theta} \int_a^{\infty} t^n p(r\theta) dr \right] = 0,$$

then (5.2) is unconditionally nonoscillatory.

Corollary 5.3 can be used to prove a theorem of Headley, see [1], Theorem 2. If the smallest eigenvalue of $a(x)$ is greater than or equal to a positive constant K and $g_0(r) = \max_{|x|=r} b(x)$, the equation

$$(5.5) \quad K \sum_{i=1}^n \frac{\partial^2 v}{\partial x_i^2} + g_0(r)v = 0$$

is a Sturmian majorant of the equation

$$(5.6) \quad \sum_{i,j=1}^n \frac{\partial}{\partial x_i} (a_{ij}(x)u_j) + b(x)u = 0.$$

If we write (5.5) as an ordinary differential equation in r and transform to remove the first derivative term, as Headley does, we arrive at the equation

$$Ky'' + [g_0(r) - K(n-1)(n-3)/4r^2]y = 0,$$

which in turn is majorized by

$$(5.7) \quad Ky'' + g_1^+(r)y = 0,$$

where $g_1^+(r) = \max(0, g_0(r) - K(n-1)(n-3)/4r^2)$. So if (5.7) is nonoscillatory so is (5.6). But Corollary 5.3 in the case $n=1$ shows that the condition

$$\int_a^{\infty} r g_1^+(r) dr < \infty$$

suffices. This is Headley's condition.

Appendix: Proof of Lemma 3.1. Suppose that f_1, \dots, f_n are functions in $L^r(E^n)$ with the property that $(\partial f_i)/(\partial x_j) = (\partial f_j)/(\partial x_i)$ as distributions. We shall construct a function v such that

$$(A.1) \quad \frac{\partial v}{\partial x_i} = f_i, \quad i = 1, \dots, n.$$

and

$$(A.2) \quad \|v\|_{r^*} \leq c \|f\|_r,$$

where $(r^*)^{-1} = r^{-1} - n^{-1}$, $f = \{f_1, \dots, f_n\}$, and c is a constant that

depends only on n and r . The proof is based on the validity of the inequality

$$(A.3) \quad \|\varphi\|_{r^*} \leq c \|\nabla\varphi\|_r$$

for functions with compact support, see [3].

Let $f_i^{(k)}$ be a modification of f_i so that $f_i^{(k)} \rightarrow f_i$ in L^r as $k \rightarrow \infty$. Set $\Gamma(x) = \text{const. } |x|^{2-n}$ if $n \geq 3$ and $\Gamma(x) = \text{const. } \log|x|$ if $n = 2$, so that $\Delta\Gamma = \varepsilon$, where ε is the Dirac distribution with support at zero, and $\Delta = \sum_{i=1}^n \partial^2/(\partial x_i^2)$. Let $\xi_h(x)$ be a smooth function with $0 \leq \xi_h \leq 1$, $\xi_h(x) = 1$ for $|x| \leq h$, $\xi_h(x) = 0$ for $|x| \geq 2h$, $|\nabla \xi_h| \leq 2h^{-1}$. Now consider the function

$$v_h^{(k)} = \sum_{i=1}^n \left(\xi_h \frac{\partial \Gamma}{\partial x_i} \right) * (f_i^{(k)} \xi_h).$$

Differentiate and use the fact that $(\partial f_i)/(\partial x_j) = (\partial f_j)/(\partial x_i)$, so that

$$\begin{aligned} \frac{\partial v_h^{(k)}}{\partial x_j} &= \sum_{i=1}^n \left(\xi_h \frac{\partial \Gamma}{\partial x_i} \right) * \left(\frac{\partial f_j^{(k)}}{\partial x_i} \xi_h \right) + \sum_{i=1}^n \left(\xi_h \frac{\partial \Gamma}{\partial x_i} \right) * \left(\frac{\partial \xi_h}{\partial x_j} f_i^{(k)} \right) \\ &= \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\xi_h \frac{\partial \Gamma}{\partial x_i} \right) * (f_j^{(k)} \xi_h) + \sum_{i=1}^n \xi_h \frac{\partial \Gamma}{\partial x_i} * \left(\frac{\partial \xi_h}{\partial x_j} f_i^{(k)} - f_j^{(k)} \frac{\partial \xi_h}{\partial x_j} \right). \end{aligned}$$

Finally

$$(A.4) \quad \begin{aligned} &\frac{\partial v_h^{(k)}}{\partial x_j} \\ &= f_j^{(k)} \xi_h + \sum_{i=1}^n \left(\frac{\partial \xi_h}{\partial x_i} \frac{\partial \Gamma}{\partial x_i} \right) * (f_j^{(k)} \xi_h) + \sum_{i=1}^n \xi_h \frac{\partial \Gamma}{\partial x_i} * \left(\frac{\partial \xi_h}{\partial x_j} f_i^{(k)} - f_j^{(k)} \frac{\partial \xi_h}{\partial x_j} \right). \end{aligned}$$

For each i

$$\left\| \frac{\partial \xi_h}{\partial x_i} \frac{\partial \Gamma}{\partial x_i} * f_j^{(k)} \xi_h \right\|_r \leq \|f_j^{(k)}\|_r \left\| \frac{\partial \xi_h}{\partial x_i} \frac{\partial \Gamma}{\partial x_i} \right\|_1 \leq \text{const. } \|f\|_r,$$

so that the first summation term on the right in (A.4) is bounded in L^r uniformly in h . Furthermore we find that for each i the same term is tending to zero in L^r_{loc} as $h \rightarrow 0$. For if B_a is the set of x with $|x| \leq a$, Hölder's inequality implies that

$$\begin{aligned} \int_{B_a} \left| \frac{\partial \xi_h}{\partial x_i} \frac{\partial \Gamma}{\partial x_i} * f_j^{(k)} \xi_h \right|^r dx &\leq \text{const.} \int |f_j^{(k)}(x)|^r \int_{B_a} \left| \frac{\partial \xi_h}{\partial x_i} \frac{\partial \Gamma}{\partial x_i}(y-x) \right| dy dx \\ &\leq \text{const.} \int_{|x| \geq h-a} |f_j(x)|^r dx \longrightarrow 0 \quad \text{as } h \longrightarrow \infty. \end{aligned}$$

Furthermore Hölder's inequality also shows that all the other terms in the right side of (A.4) tend to zero in L^r as $h \rightarrow \infty$.

So $(\partial v_h^{(k)})/(\partial x_j) - f_j^{(k)}$ is bounded in L^r as $h \rightarrow \infty$ and converges to

zero in L^r_{10c} . For some sequence of h 's then tending to infinity,

$$\frac{\partial v_h^{(k)}}{\partial x_j} \longrightarrow f_j^{(k)}$$

weakly in L^r . Also (A.3) holds for each $v_h^{(k)}$ so there is a further subsequence of h 's with $v_h^{(k)} \rightarrow v^{(k)}$ weakly in L^{r^*} . Hence $(\partial v^{(k)})/(\partial x_j) = f_j^{(k)}$ as distributions and (A.2) holds for $v = v^{(k)}$ and $f = f^{(k)}$. Finally for a subsequence of h 's tending to ∞ , $v^{(k)} \rightarrow v$ weakly in L^{r^*} and $(\partial v^{(k)})/(\partial x_j) \rightarrow f_j$ in L^r . So $(\partial v)/(\partial x_j) = f_j$ and (A.2) holds. This completes the proof.

REFERENCES

1. V. B., Headley, *Some oscillation properties of selfadjoint elliptic equations*, Proc. Amer. Math. Soc., **25** (1970), 824-829.
2. V. B. Headley and C. A. Swanson, *Oscillation criteria for elliptic equations*, Pacific J. Math., **27** (1968), 501-506.
3. L. Nirenberg, *On elliptic partial differential equations*, Ann. della Scuola Nor. Sup. Pisa, **13** (1959), 115-162.
4. John Piepenbrink, *Integral inequalities and theorems of Liouville type*, J. Math. Anal. Appl., **26** (1969), 630-639.
5. S. L. Sobolev, *On a theorem of functional analysis*, Amer. Math. Soc. Translations, (2) **34** (1963), 39-68.
6. ———, *Einige Anwendungen der Funktionalanalysis auf Gleichungen der Mathematischen Physik*, Akademie-Verlag, Berlin 1964.
7. G. Stampachia and M. K. V. Murthy, *Boundary value problems for some degenerate-elliptic operators*, Ann. di Mat. pura ed. Appl. LXXX, (1968) 1-122.

Received June 24, 1971. This work was in part supported by NSF grant GP-7041X, and Air Force Grant AFOSR 883-67.

UNIVERSITY OF MINNESOTA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 42, No. 1

January, 1972

Tage Bai Andersen, <i>On Banach space valued extensions from split faces</i>	1
David Marion Arnold, <i>A duality for quotient divisible abelian groups of finite rank</i>	11
Donald Pollard Ballou, <i>Shock sets for first order nonlinear hyperbolic equations</i>	17
Leon Brown and Lowell J. Hansen, <i>On the range sets of H^p functions</i>	27
Alexander Munro Davie and Arne Stray, <i>Interpolation sets for analytic functions</i>	33
M. G. Deshpande, <i>Structure of right subdirectly irreducible rings. II</i>	39
Barry J. Gardner, <i>Some closure properties for torsion classes of abelian groups</i>	45
Paul Daniel Hill, <i>Primary groups whose subgroups of smaller cardinality are direct sums of cyclic groups</i>	63
Richard Allan Holzinger, <i>When certain natural maps are equivalences</i>	69
Donald William Kahn, <i>A note on H-equivalences</i>	77
Joong Ho Kim, <i>R-automorphisms of $R[t][[X]]$</i>	81
Shin'ichi Kinoshita, <i>On elementary ideals of polyhedra in the 3-sphere</i>	89
Andrew T. Kitchen, <i>Watts cohomology and separability</i>	99
Vadim Komkov, <i>A technique for the detection of oscillation of second order ordinary differential equations</i>	105
Charles Philip Lanski and Susan Montgomery, <i>Lie structure of prime rings of characteristic 2</i>	117
Andrew Lenard, <i>Some remarks on large Toeplitz determinants</i>	137
Kathleen B. Levitz, <i>A characterization of general Z.P.I.-rings. II</i>	147
Donald A. Lutz, <i>On the reduction of rank of linear differential systems</i>	153
David G. Mead, <i>Determinantal ideals, identities, and the Wronskian</i>	165
Arunava Mukherjea, <i>A remark on Tonelli's theorem on integration in product spaces</i>	177
Hyo Chul Myung, <i>A generalization of the prime radical in nonassociative rings</i>	187
John Piepenbrink, <i>Rellich densities and an application to unconditionally nonoscillatory elliptic equations</i>	195
Michael J. Powers, <i>Lefschetz fixed point theorems for a new class of multi-valued maps</i>	211
Aribindi Satyanarayan Rao, <i>On the absolute matrix summability of a Fourier series</i>	221
T. S. Ravisankar, <i>On Malcev algebras</i>	227
William Henry Ruckle, <i>Topologies on sequences spaces</i>	235
Robert C. Shock, <i>Polynomial rings over finite dimensional rings</i>	251
Richard Tangeman, <i>Strong heredity in radical classes</i>	259
B. R. Wenner, <i>Finite-dimensional properties of infinite-dimensional spaces</i>	267