

Pacific Journal of Mathematics

INTERPOLATION BY ANALYTIC FUNCTIONS

ARNE STRAY

INTERPOLATION BY ANALYTIC FUNCTIONS

ARNE STRAY

It is shown that interpolation problems for $R(X)$, $A(X)$ and $H^\infty(X^\circ)$ are local problems whenever X is a compact plane set.

Introduction and notation. Let X be compact plane set, X° its interior and $\partial X = X \setminus X^\circ$ its boundary.

$H^\infty(X^\circ)$ denotes all bounded complex-valued analytic functions on X° . $A(X)$ is all continuous functions on X which are analytic in X° . $R(X)$ denotes the uniform closure on X of the rational functions with poles outside X .

A subset E of X is an interpolation set for $A(X)$ if $A(X) \setminus E$ (the restrictions to E of the functions in $A(X)$) equals the space $C(E)$ of all continuous complex-valued functions on E .

E is called a peak set for $A(X)$ if there exists $f \in A(X)$ such that $f = 1$ on E and $|f(x)| < 1$ if $x \in X \setminus E$.

A peak interpolation set for $A(X)$ is a set E which has both these properties. Peak and interpolation sets for $R(X)$ are defined in the same way.

A sequence $S = \{z_n\}$ of distinct points is called an interpolating sequence for $H^\infty(X^\circ)$ if for any bounded sequence $\{w_n\}$ of complex numbers there exists $f \in H^\infty(X^\circ)$ such that $f(z_n) = w_n$ for each n . (For more about interpolating sequences see Ch. 10 in [3].)

If F is a subset of the complex plane we give it (as a topological space) the topology induced from \mathbb{C} . $C_b(F)$ is the Banach space of all bounded continuous complex-valued functions on F . We also consider $H^\infty(X^\circ)$, $R(X)$ and $A(X)$ as Banachspaces with the usual sup norm.

Let us mention two other Banach-spaces of analytic functions which has not been much studied yet, but which may be useful in characterizing interpolation sets for $R(X)$ and $A(X)$ among other things.

$HR(X)$ denotes all functions on X° which are pointwise limits on X° of bounded sequences in $R(X)$. For each $f \in HR(X)$ we define

$$\|f\|_{HR} = \inf \{ \sup_n \|f_n\| : \{f_n\} \subset R(X), f_n \longrightarrow f \text{ pointwise on } X^\circ \}.$$

With this norm $HR(X)$ clearly is a Banach space. In the same way we define $HA(X)$ corresponding to $A(X)$ and it is also a Banach space with the norm $\| \cdot \|_{HA}$. Very recently A. M. Davie has shown that the norm $\| \cdot \|_{HA}$ is the same as sup norm on X° and the same is proved for $\| \cdot \|_{HR}$ if almost every point of ∂X (w.r.t. area) is a peak point for $R(X)$. We shall not need these interesting results

here. (See [1] for his results.) Some results about $HR(X)$ can be found in [2].

If f is a complex-valued function defined on a set F and $S \subset F$ is a subset we define $\|f\|_S$ as $\sup \{|f(z)| : z \in S\}$.

A typical problem we shall study in this paper is the following:

Let S be a sequence in X^0 . What local conditions on S are sufficient to conclude that S is an interpolating sequence for $H^\infty(X^0)$?

An obvious necessary condition is that $S \cap \Delta_z$ is an interpolating sequence for $H^\infty(X^0)$ whenever Δ_z is an open disc centered at z for which $\Delta_z \cap S \neq \emptyset$.

Suppose that the following weaker condition is satisfied:

(*): For every $z \in \bar{S}$ (the closure of S) there exists $\delta_z > 0$ such that $S \cap \Delta_z$ is an interpolating sequence for $H^\infty(\Delta_z \cap X^0)$ where $\Delta_z = \{w : |w - z| < \delta_z\}$.

We shall then by definition say that S admits local H^∞ -interpolation w.r.t. X^0 .

Our main result is the following:

THEOREM 1. *Let X be a compact set with nonempty interior X^0 . A sequence S in X^0 is an interpolating sequence for $H^\infty(X^0)$ if and only if S admits local H^∞ -interpolation w.r.t. X^0 .*

Some time after Theorem 1 was proved we learnt about a result of J. Rainwater which has some connection with Theorem 1. If in the definition of local H^∞ -interpolation the condition (*) had been replaced by the other necessary condition for interpolation mentioned above Theorem 1 would be a somewhat weaker result.

We want to point out this weaker result is easy to deduce from J. Rainwaters paper. (See [4].) We also want to point out that a theorem of E. L. Stout on interpolating sequences in multiply connected domains is an easy consequence of Theorem 1. (See [5].)

Interpolating sequences can clearly also be defined for $HR(X)$ and $HA(X)$. It should also be clear what is meant by saying that a sequence $S \subset X^0$ admits local HR -interpolation (or HA -interpolation) w.r.t. X .

It will follow from our proof that Theorem 1 also holds for $HR(X)$ and $HA(X)$. We shall give some reasons for this at the end of the proof.

LEMMA 1. *Let X be as in Theorem 1 and $z_0 \in \partial X$. Let $0 < r_1 < r_2$ and define $O_1 = \{w : |w - z_0| < r_1\}$ and $O_2 = \{w : |w - z_0| > r_2\}$. Suppose there exists $z_1 \in C \setminus X$ such that $r_2 > |z_1 - z_0| > r_1$*

Let S_i be an interpolating sequence for $H^\infty(X^0 \cap O_i)$ for $i = 1, 2$.

Suppose $\bar{S}_i \subset 0_i$ for $i = 1, 2$.

Then $S = S_1 \cup S_2$ is an interpolating sequence for $H^\infty(X^0)$.

Proof. Put $\Gamma_i = \partial 0_i$ for $i = 1, 2$.

Then $\text{dist} \cdot (S, \Gamma_i) > 0$.

Assume $h \in H^\infty(X^0 \cap 0_1)$. Extend it to C by defining $h(z) = 0$ if $z \in X^0 \cap 0_1$.

Let $\delta > 0$ be given. Then cover C by open discs $\Delta_n = \Delta(z_n, \delta)$ (of radius δ and centered at z_n) and choose continuously differentiable functions g_n supported on Δ_n as in the scheme for approximation described on page 210 in [2].

Let T_{g_n} be the integral operator on $L^\infty(dx dy)$ defined by

$$\begin{aligned} T_{g_n}(f)(w) &= \frac{1}{\pi} \iint \frac{f(w) - f(z)}{w - z} \frac{\partial g_n}{\partial \bar{z}} dx dy \\ &= f(w) \cdot g_n(w) + \frac{1}{\pi} \iint \frac{f(z)}{z - w} \frac{\partial g_n}{\partial \bar{z}} dx dy. \end{aligned}$$

We mention that $T_{g_n}(f)$ is analytic outside the support of g_n and wherever f is and that $T_{g_n}(f)$ is continuous wherever f is.

Also $f - T_{g_n}(f)$ is analytic in the interior of the set where g_n attains the value 1. (See on p. 28-29 in [2] for more details.)

Put $h_n = T_{g_n}(h)$. We are only interested in those n for which $\bar{\Delta}_n \cap \Gamma_1 \cap X \neq \emptyset$. Assume this happens if and only if $1 \leq n \leq N$.

Then $h - \sum_{i=1}^N h_n = h - T_{(\sum_{i=1}^N g_n)}(h)$ is analytic near $\Gamma_1 \cap X$ since $\sum_{i=1}^N g_i$ equals 1 near $\Gamma_1 \cap X$.

Now there exist functions $\{H_n\}_{n=1}^N$ analytic outside a compact subset of $D_n = \{w: |w - z_n| < 2\delta\} \setminus 0_1$ such that $h_n - H_n$ has a triple zero in the Taylor expansion at infinity and in our situation we can obtain $\|H_n\| \leq c_1 \|h\|$ where c_1 is an absolute constant. (See Theorem 7.4 on p. 213 in [2] and the proof of it.)

Now one has to observe two important facts.

(a) If B is a subset of C and $\text{dist}(B, \Gamma_1 \cap X) > 0$ and $\varepsilon > 0$ one can choose δ depending only on ε and $\text{dist}(B, \Gamma_1 \cap X)$ so small that the sum $f_\delta = h - \sum_{i=1}^N (h_n - H_n)$ satisfies

$$(1) \quad \|h - f_\delta\|_B \leq \varepsilon \|h\|.$$

(b) The functions H_n can be chosen such that its singularities lies on a fixed compact subset of D_n independent of h .

In fact one can find two functions $F_{n,1}$ and $F_{n,2}$ analytic outside a compact subset of D_n such that $\|F_{n,1}\| + \|F_{n,2}\| \leq 20$ and $H_n = \lambda_{n,1}(h)F_{n,1} + \lambda_{n,2}(h)F_{n,2}$. (See lemma 6.3 on page 209 in [2].)

Here $\lambda_{n,k}(h)$ is a complex number and we have

$$(2) \quad |\lambda_{n,k}(h)| \leq c_2 \|h\| \quad \text{for } k = 1, 2,$$

where c_2 in our situation is an absolute constant. If $F_{n,k}$ is constructed as in the mentioned lemma in [2]. We also mention that the maps $h \rightarrow \lambda_{n,k}(h)$ are linear.

(Some details indicating how this can be done, can be found in the proof of Lemma 3.1 in [4].)

Given $\varepsilon > 0$ we first choose δ so small that

$$(3) \quad \|h - f_\delta\|_S < \frac{\varepsilon \|h\|}{4}$$

whenever h is as above. Then choose rational functions $r_{n,k}$ with poles only at z_1 such that

$$(4) \quad \sum_{n=1}^N (\|F_{n,1} - r_{n,1}\|_{0_1 \cup 0_2} + \|F_{n,2} - r_{n,2}\|_{0_1 \cup 0_2}) < \frac{\varepsilon}{4c_2}.$$

Now define $A_1: H^\infty(X^0 \cap 0_1) \rightarrow H^\infty(X^0)$ by

$$A_1(h) = [h - \sum_1^N (h_n - (\lambda_{n,1}(h)r_{n,1} + \lambda_{n,2}(h)r_{n,2}))] \Big| X^0.$$

From (1), (2), (3) and (4) we deduce that

- (i) $\|A_1(h)\| \leq c_4 \|h\|$ where c_4 depends only on the rational functions $r_{n,k}$.
- (ii) $\|A_1(h) - h\|_S \leq \varepsilon \|h\|/4 + \varepsilon \|h\|/4 = \varepsilon \|h\|/2$.

In addition we also mention that A_1 is linear but this fact will not be needed.

In exactly the same way we define a map $A_2: H^\infty(X^0 \cap 0_2) \rightarrow H^\infty(X^0)$.

Suppose now $f \in C_b(S)$. By the open mapping theorem applied to the restriction $H^\infty(0_i \cap X^0) \rightarrow C_b(S_i)$ for $i = 1, 2$, there exists a constant M independent of f and functions $h_i \in H^\infty(0_i \cap X^0)$ such that

$$\|h_i\| \leq M \|f\| \text{ and } h_i = f \text{ on } S_i \text{ for } i = 1, 2.$$

Put $h_i = 0$ outside $0_i \cap X^0$ and define $g = A_1(h_1) + A_2(h_2)$.

Then $g \in H^\infty(X^0)$, $\|g\| \leq 2c_4 M \|f\|$ and $\|f - g\|_S = \|A_1(h_1) - h_1 + A_2(h_2) - h_2\|_S \leq \varepsilon M \|f\| \leq 1/2 \|f\|$ if we choose $\varepsilon \leq 1/2M$.

Put $g_1 = g$ and assume g_1, \dots, g_n constructed such that

$$\|g_k\| \leq 2^{-k+2} c_4 M \|f\| \text{ for } 1 \leq k \leq n$$

and

$$\left\| f - \sum_1^n g_j \right\|_S \leq \frac{\|f\|}{2^n}.$$

By the approximation technique above one easily find $g_{n+1} \in H^\infty(X^0)$ such that $\|g_{n+1}\| \leq 2^{-n-1} c_4 M \|f\|$ and

$$\left\| f - \sum_1^{n+1} g_j \right\|_S \leq \frac{\|f\|}{2^{n+1}}.$$

By induction the series $\sum_1^\infty g_n \in H^\infty(X^0)$ interpolates f on S .

LEMMA 2. *Let S be a sequence in X^0 with no clusterpoints in X^0 . Assume there exist n points z_1, \dots, z_n and numbers $r_k > s_k > t_k$ for $1 \leq k \leq n$ such that the open discs $\{\Delta(z_k, t_k)\}_{k=1}^n$ cover \bar{S} .*

Assume also that $(C \setminus X) \cap \{w: r_k > |w - z_k| > s_k\}$ and $(C \setminus X) \cap \{w: |s_k > |w - z_k| > t_k\}$ are nonempty for each k . If for each k , $\Delta(z_k, r_k) \cap S$ is an interpolating sequence for $H^\infty(X^0)$ then also S is.

Proof. We can assume $n \geq 2$ and by induction the lemma to be true if n is replaced by $n - 1$.

Put $S_1 = S \cap \Delta(z_n, t_n)$.

By hypothesis $S_2 = S \cap (\bigcup_1^{n-1} \Delta(z_k, s_k))$ is an interpolating sequence for $H^\infty(X^0)$ and given $f \in C(S)$ we can find $h_1 \in H^\infty(X^0)$ such that $h_1 = f$ on S_2 .

The choose $h_2 \in H(X^0)$ equal to $f - h_1$ on $\Delta(z_n, r_n)$.

By Lemma 1 we can find h_3 in $H^\infty(X^0)$ such that $h_3 = 1$ on S_1 and $h_3 = 0$ on $S_2 \setminus \Delta(z_n, s_n)$.

Then $h_1 + h_2 h_3 = f$ on S .

Proof of Theorem 1. We have to show that the local condition implies that S is an interpolating sequence.

S has no clusterpoints in X^0 and for each $z \in (\partial X) \cap \bar{S}$ we can find $r_z > 0$ and such that $\Delta(z, r_z) \cap S$ is an interpolating sequence for $H^\infty(X_z^0)$ where $X_z^0 = \{w: |w - z| < 2r_z\} \cap X^0$. By Lemma 1 $S \cap \Delta(z, r_z)$ is an interpolating sequence for $H^\infty(X^0)$.

Since $z \in \partial X$ we can choose $s_z > t_z > 0$ such that $(C \setminus X) \cap \{w: r_z > |w - z| > s_z\}$ and $(C \setminus X) \cap \{w: s_z > |w - z| > t_z\}$ are nonempty.

Since $\bar{S} \cap (\partial X)$ is compact we can obtain the hypothesis of Lemma 2 for a set $S' \subset S$ such that $S \setminus S'$ is finite.

But if S' is an interpolating sequence for $H^\infty(X^0)$ then clearly also S is.

REMARK. To prove Theorem 1 in case $H = HR(X)$ one must modify the arguments slightly in the proof of Lemma 1. We use the notation from that lemma.

Given $f \in C(S)$ one finds $h_i \in HR(\bar{0}_i \cap X)$ equal to f on S_i such that $\|h_i\|_{HR} \leq M \|f\|$ where M is a constant independent of S found by using the open mapping theorem.

Then we find a sequence $\{g_n^i\}_{n=1}^\infty \subset C(S^2)$ analytic in a neighbourhood

of $X \cap \bar{0}_i$ (depending on n) such that $\sup_n \|g_n^i\| \leq 2M\|f\|$ and such that $g_n^i \rightarrow h_i$ pointwise on the interior of $X \cap \bar{0}_i$. (S^2 denotes the extend complex plane with the usual topology.) We can also assume g_n^i converges in the w^* -topology of $L^\infty(dx dy)$ to a function \tilde{h}_i equal to h_i on $0_i \cap X^0$ such that $\|\tilde{h}_i\|_\infty \leq 2M\|f\|$.

We can assume $\tilde{h}_i = 0$ outside $\bar{0}_i$.

Then it is easy to see that $\sum_{i=1}^2 A_i(\tilde{h}_i)$ will approximate f well on S and that $A_i(g_n^i)|X$ belongs to $R(X)$ for all n and that $A_i(g_n^i) \rightarrow A_i(\tilde{h}_i)$ pointwise on X^0 . Also $\|A_i(\tilde{h}_i)\|_{HR} \leq k \cdot M\|f\|$. (k is independent of f .)

With these remarks Lemma 1 also applies for $HR(X)$. It is clear that similar modifications give Lemma 1 also for $HA(X)$.

But then the rest of the proof of Theorem 1 including the proof of Lemma 2 applies almost directly.

COROLLARY 1. *Let X be a compact plane set and E_a closed subset.*

Then E is an interpolation set for $R(X)$ if and only if for each $z \in E$ there exists a closed disc $N_z = \{w: |w - z| \leq r_z\}$ such that $E \cap N_z$ is an interpolation set for $R(X \cap N_z)$.

Proof. Clearly $E_z = E \cap \{w: |w - z| \leq z/2\}$ is an interpolation set for $R(X \cap N_z)$.

The approximation technique used in the proof of Lemma 1 shows that E_z then is an interpolation for $R(X)$.

But then the corollary follows from Rainwaters result.

REMARK. A similar corollary also clearly holds for $A(X)$.

Finally we state a theorem for $R(X)$ which is not difficult to prove. Perhaps it makes the space $HR(X)$ a little more attractive.

THEOREM 3. *Let S be a closed subset of a compact plane set X . Suppose that*

- (i) $S \cap \partial X$ is a peak interpolation set for $R(X)$
- (ii) $S \cap X^0$ is an interpolating sequence for $HR(X)$.

Then $R(X)|S = C(S)$.

One proves Theorem 3 by showing that for every $f \in C(S)$ there exists $g \in R(X)$ such that

$$\|f - g\|_S \leq \frac{1}{2} \|f\| \quad \text{and} \quad \|g\| \leq M \|f\|$$

where M is independent of f . This is sufficient by the approximation

argument at the end of the proof of Lemma 1.

First choose $f_1 \in R(X)$ such that $f_1 = f$ on $S \cap \partial K$ and $\|f_1\| \leq \|f\|$.

Interpolate then $f - f_1$ on $S \cap X^0$ by $f_2 \in HR(X)$ such that $\|f_2\|_{HR} \leq M_1 \|f\|$ where M_1 is independent of f .

If $\varepsilon > 0$ choose an open set $V_\varepsilon \supset S \cap \partial X$ such that $|f_2| < \varepsilon$ on $S \cap X^0 \cap V_\varepsilon$.

Choose also $f_3 \in R(X)$ such that $\|f_3\| \leq 2$, $f_3 = 0$ on $S \cap \partial X$ and $|1 - f_3| < \varepsilon$ on $X \setminus V_\varepsilon$ and $f_4 \in R(X)$ such that $|f_4(z) - f_2(z)| \leq \varepsilon$ for all $z \in S$ where $|f_3(z)| \geq \varepsilon$ and such that $\|f_4\| \leq 2\|f_2\|_{HR} \leq 2M_1 \|f\|$.

Then put $g = f_1 + f_3 f_4$. We have $\|g\| \leq (1 + 4M_1) \|f\|$ and $\|f - g\|_\varepsilon \leq \varepsilon \|f\| (2 + 3M_1)$.

So with $\varepsilon = 1/(4 + 6M_1)$ we have what we want.

Finally I want to thank Dr. A. M. Davie for some very helpful correspondence which gave our results considerably greater generality.

REFERENCES

1. A. M. Davie, *Bounded limits of analytic functions*, (to appear).
2. T. W. Gamelin, *Uniform Algebras*, Prentice Hall, Englewood Cliffs. N. J. 1969.
3. K. Hoffmann, *Banach Spaces of Analytic Functions*, Prentice Hall, Englewood Cliffs. N. J. 1962.
4. J. Rainwater, *A remark on regular Banach algebras*, Proc. Amer. Math. Soc., **18** (1967), 255-256.
5. E. L. Stout, *Two theorems concerning functions holomorphic in multiply connected domains*, B. A. M. S., **69** (1963), 527-530.
6. A. Stray, *Approximation and Interpolation*. (to appear in Pacific J. Math.).

Received April 23, 1971.

UNIVERSITY OF OSLO

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Stephen Richard Bernfeld, <i>The extendability of solutions of perturbed scalar differential equations</i>	277
James Edwin Brink, <i>Inequalities involving f_p and $f^{(n)}_q$ for f with n zeros</i>	289
Orrin Frink and Robert S. Smith, <i>On the distributivity of the lattice of filters of a groupoid</i>	313
Donald Goldsmith, <i>On the density of certain cohesive basic sequences</i>	323
Charles Lemuel Hagopian, <i>Planar images of decomposable continua</i>	329
W. N. Hudson, <i>A decomposition theorem for biadditive processes</i>	333
W. N. Hudson, <i>Continuity of sample functions of biadditive processes</i>	343
Masako Izumi and Shin-ichi Izumi, <i>Integrability of trigonometric series. II</i>	359
H. M. Ko, <i>Fixed point theorems for point-to-set mappings and the set of fixed points</i>	369
Gregers Louis Krabbe, <i>An algebra of generalized functions on an open interval: two-sided operational calculus</i>	381
Thomas Latimer Kriete, III, <i>Complete non-selfadjointness of almost selfadjoint operators</i>	413
Shiva Narain Lal and Siya Ram, <i>On the absolute Hausdorff summability of a Fourier series</i>	439
Ronald Leslie Lipsman, <i>Representation theory of almost connected groups</i>	453
James R. McLaughlin, <i>Integrated orthonormal series</i>	469
H. Minc, <i>On permanents of circulants</i>	477
Akihiro Okuyama, <i>On a generalization of Σ-spaces</i>	485
Norberto Salinas, <i>Invariant subspaces and operators of class (S)</i>	497
James D. Stafney, <i>The spectrum of certain lower triangular matrices as operators on the l_p spaces</i>	515
Arne Stray, <i>Interpolation by analytic functions</i>	527
Li Pi Su, <i>Rings of analytic functions on any subset of the complex plane</i>	535
R. J. Tondra, <i>A property of manifolds compactly equivalent to compact manifolds</i>	539