

Pacific Journal of Mathematics

ON ABSOLUTE DE LA VALLÉE POUSSIN SUMMABILITY

B. T. Y. KWEE

ON ABSOLUTE DE LA VALLÉE POUSSIN SUMMABILITY

B. KWEE

Gronwall proved that $(C, r) \subseteq (V - P)$ for $r \geq 0$, where (C, r) and $(V - P)$ denote Cesàro and de la Vallée Poussin summability. It is proved in this paper that $|C, r| \subseteq |V - P|$ for $r \geq 0$.

1. Introduction. Let

$$V_n = \sum_{k=1}^n \frac{(n!)^2}{(n-k)!(n+k)!} a_k \quad (n \geq 0).$$

If $\lim_{n \rightarrow \infty} V_n = s$, we say that the series is summable $(V - P)$ to s .
 If

$$\sum_{n=1}^{\infty} |V_n - V_{n-1}| < \infty.$$

The series $\sum_{n=0}^{\infty} a_n$ is said to be summable $|V - P|$.

Hyslop [2] proved that the $(V - P)$ method is equivalent to the $(A, 2)$ method defined by

$$\lim_{x \rightarrow 0+} \sum_{n=0}^{\infty} a_n e^{-n^2 x} = s$$

for all series $\sum_{n=0}^{\infty} a_n$ which satisfy the condition $a_n = O(n^c)$, where c is any constant, and that the inclusion $(A, 2) \subseteq (V - P)$ is false without restriction.

Kuttner [3] has shown that $(V - P) \subseteq (A, 2)$ without restriction.

Gronwall [1] proved that $(C, r) \subseteq (V - P)$ for $r \geq 0$, where (C, r) denotes the Cesàro summability of order r .

In this paper, we shall prove

THEOREM A. $|C, r| \subseteq |V - P|$ for $r \geq 0$.

2. Proof of Theorem A. Since it is well-known that $|C, r|$ implies $|C, r'|$ for $-1 < r \leq r'$, it is enough to consider the case r an integer. Now, writing

$$V_n = v_0 + v_1 + \cdots + v_n,$$

we find that

$$(1) \quad \begin{cases} v_0 = a_0, \\ v_n = \sum_{k=1}^n \frac{((n-1)!)^2}{(n-k)!(n+k)!} k^2 a_k \quad (n \geq 1). \end{cases}$$

Now write $\tau_k = \tau_k^r$ for the (C, r) mean of the sequence $\{ka_k\}$; thus the assumption that $\sum_{n=0}^{\infty} a_n$ is summable $|C, r|$ is equivalent to

$$(2) \quad \sum_{n=0}^{\infty} \frac{|\tau_n|}{n} < \infty .$$

If we take $((n-1)!)/(n-k)(n+k)!$ as meaning 0 whenever $k > n$, we deduce from (1) by n partial summations that, for $n \geq 1$,

$$v_n = \sum_{k=1}^n \Delta_k^r \left\{ \frac{((n-1)!)^2 k}{(n-k)!(n+k)!} \right\} \binom{k+r}{k} \tau_k .$$

Now it is well-known that in order that the series-to-series transformation

$$b_n = \sum_{k=0}^{\infty} \alpha_{nk} a_k$$

should be that $\sum_{n=0}^{\infty} |b_n|$ converges whenever $\sum_{n=0}^{\infty} |a_n|$ does so, it is necessary and sufficient that

$$\sum_{n=0}^{\infty} |\alpha_{nk}|$$

should be bounded. Thus it is enough to show that, for $k \geq 1$,

$$(3) \quad \sum_{n=k}^{\infty} \left| \Delta_k^r \left\{ \frac{((n-k)!)^2 k}{(n-k)!(n+k)!} \right\} \right| = O(k^{-r-1}) .$$

It is easily seen by induction on r that

$$\Delta_k^r \left\{ \frac{((n-1)!)^2 k}{(n-k)!(n+k)!} \right\} = \frac{A^r(n, k)((n-1)!)^2}{(n-k)!(n+k+r)!} ,$$

where $A^r(n, k)$ is defined inductively by

$$(4) \quad \begin{cases} A^0(n, k) = k , \\ A^{r+1}(n, k) = (n+k+r+1)A^r(n, k) - (n-k)A^r(n, k+1) . \end{cases}$$

Write $P_j(k)$ for a polynomial in k of degree not exceeding j , possibly different at each occurrence (thus $P_0(k)$ denotes a constant). We deduce from (4) by induction that

$$\begin{aligned} A^{2s}(n, k) &= \sum_{j=0}^s P_{2j+1}(k) n^{s-j} , \\ A^{2s+1}(n, k) &= \sum_{j=0}^{s+1} P_{2j}(k) n^{s+1-j} . \end{aligned}$$

Hence, uniformly in the ranges stated

$$A^r(n, k) = \begin{cases} O(n^{(r+1)/2}) & (1 \leq k \leq n^{1/2}), \\ O(K^{r+1}) & (n^{1/2} < k \leq n). \end{cases}$$

Next, for large n uniformly in $k \leq n^{2/3}$ we have, by Stirling's formula

$$\frac{(n!)^2}{(n - k)!(n + k)!} = O(H(n, k)),$$

where

$$H(n, k) = \left(1 - \frac{k}{n}\right)^{-n+k-1/2} \left(1 + \frac{k}{n}\right)^{-n-k-1/2}.$$

We have

$$\log H(n, k) = -\frac{k^2}{n} + O\left(\frac{k^3}{n^2}\right).$$

Now since we supposing that $k \leq n^{2/3}$ we have

$$(5) \quad \exp\left\{O\left(\frac{k^3}{n^2}\right)\right\} = O(1)$$

so that

$$\frac{(n!)^2}{(n - k)!(n + k)!} = O\left\{\exp\left(-\frac{k^2}{n}\right)\right\}.$$

This will not apply if $k > n^{2/3}$. Since we cannot then assert (5). However, for fixed n , $(n!)^2/(n - k)!(n + k)!$ is a decreasing function of k so that, for $k > n^{2/3}$,

$$\frac{(n!)^2}{(n - k)!(n + k)!} = O\{\exp(-n^{1/3})\}.$$

Also, it is trivial that

$$\frac{((n - 1)!)^2}{(n - k)!(n + k + 2)!} = \frac{(n!)^2}{(n - k)!(n + k)!} O(n^{-r-2}).$$

Combining these results, we find that

$$A_k^r\left\{\frac{((n - 1)!)^2 k}{(n - k)!(n + k)!}\right\} = \begin{cases} O(n^{-(r+3)/2}) & (1 \leq k \leq n^{1/2}), \\ O\left(\frac{k^{r+1}}{n^{r+2}} \exp\left(-\frac{k^2}{n}\right)\right) & (n^{1/2} < k \leq n^{2/3}), \\ O(n^{-1} \exp(-n^{-(1/3)})) & (n^{2/3} < k \leq n). \end{cases}$$

Thus the sum (3) is

$$O\left\{\sum_{k \leq n < k^{3/2}} \frac{1}{n} \exp(-n^{-(1/3)})\right\} + O\left\{\sum_{k^{3/2} \leq n < k^2} \frac{k^{r+1}}{n^{r+2}} \exp\left(-\frac{k^2}{n}\right)\right\} + O\left\{\sum_{n \geq k^2} \frac{1}{n^{(r+3)/2}}\right\}$$

$$= O(I_1) + O(I_2) + O(I_3),$$

say. It is clear that

$$I_1 = O(k^{-r-1}),$$

$$I_3 = O(k^{-r-1})$$

so we need consider only I_2 . Now for fixed k

$$\frac{k^{r+1}}{y^{r+2}} \exp\left(-\frac{k^2}{y}\right)$$

is increasing for $y < y_0$ and decreasing for $y > y_0$, where $y_0 = y_0(k) = k^2/(r+2)$. Hence

$$(6) \quad I_2 \leq k^{r+1} \int_{k^{3/2-1}}^{k^{2+1}} \frac{1}{y^{r+2}} \exp\left(-\frac{k^2}{y}\right) dy + \frac{k^{r+1}}{y_0^{r+2}} \exp\left(-\frac{k^2}{y_0}\right).$$

The second term on the right of (6) is a constant multiple of k^{-r-3} . The first does not exceed

$$k^{r+1} \int_0^\infty \frac{1}{y^{r+2}} \exp\left(-\frac{k^2}{y}\right) dy.$$

Putting $y = k^2/w$, this becomes

$$k^{-r-1} \int_0^\infty w^r e^{-w} dw = \Gamma(r+1) k^{-r-1},$$

hence the result.

REFERENCES

1. T. H. Gronwall, *Über eine summationsmethode und ihre anwendung auf die Fouriersche reihe*, J. Für Math., **147** (1916), 16-25.
2. J. M. Hyslop, *Some relations between the de la Vallée Poussin and Abel methods of summability*, Proc. London Math. Soc., (2), **40** (1936), 449-467.
3. B. Kuttner, *The relation between de la Vallée Poussin and Abel summability*, Proc. London Math. Soc., **44** (1938), 92-99.

Received May 28, 1971.

UNIVERSITY OF MALAYA
KUALA LUMPUR
MALAYSIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 108 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics

Vol. 42, No. 3

March, 1972

Catherine Bandle, <i>Extensions of an inequality by Pólya and Schiffer for vibrating membranes</i>	543
S. J. Bernau, <i>Topologies on structure spaces of lattice groups</i>	557
Woodrow Wilson Bledsoe and Charles Edward Wilks, <i>On Borel product measures</i>	569
Eggert Briem and Murali Rao, <i>Normpreserving extensions in subspaces of $C(X)$</i>	581
Alan Seymour Cover, <i>Generalized continuation</i>	589
Larry Jean Cummings, <i>Transformations of symmetric tensors</i>	603
Peter Michael Curran, <i>Cohomology of finitely presented groups</i>	615
James B. Derr and N. P. Mukherjee, <i>Generalized quasicenter and hyperquasicenter of a finite group</i>	621
Erik Maurice Ellentuck, <i>Universal cosimple isols</i>	629
Benny Dan Evans, <i>Boundary respecting maps of 3-mainfolds</i>	639
David F. Fraser, <i>A probabilistic method for the rate of convergence to the Dirichlet problem</i>	657
Raymond Taylor Hoobler, <i>Cohomology in the finite topology and Brauer groups</i>	667
Louis Roberts Hunt, <i>Locally holomorphic sets and the Levi form</i>	681
B. T. Y. Kwee, <i>On absolute de la Vallée Poussin summability</i>	689
Gérard Lallement, <i>On nilpotency and residual finiteness in semigroups</i>	693
George Edward Lang, <i>Evaluation subgroups of factor spaces</i>	701
Andy R. Magid, <i>A separably closed ring with nonzero torsion pic</i>	711
Billy E. Rhoades, <i>Commutants of some Hausdorff matrices</i>	715
Maxwell Alexander Rosenlicht, <i>Canonical forms for local derivations</i>	721
Cedric Felix Schubert, <i>On a conjecture of L. B. Page</i>	733
Reinhard Schultz, <i>Composition constructions on diffeomorphisms of $S^p \times S^q$</i>	739
J. P. Singhal and H. M. (Hari Mohan) Srivastava, <i>A class of bilateral generating functions for certain classical polynomials</i>	755
Richard Alan Slocum, <i>Using brick partitionings to establish conditions which insure that a Peano continuum is a 2-cell, a 2-sphere or an annulus</i>	763
James F. Smith, <i>The p-classes of an H^*-algebra</i>	777
Jack Williamson, <i>Meromorphic functions with negative zeros and positive poles and a theorem of Teichmüller</i>	795
William Robin Zame, <i>Algebras of analytic functions in the plane</i>	811