Pacific Journal of Mathematics

THE p-CLASSES OF AN H*-ALGEBRA

JAMES F. SMITH

Vol. 42, No. 3

March 1972

THE p-CLASSES OF AN H^* -ALGEBRA

JAMES F. SMITH

This paper considers a family of *-subalgebras of a semisimple H*-algebra A. For 0 a nonnegative extended $real value <math>|a|_p$ is associated with each a in A; then the p-class A_p is defined to be $\{a \in A: |a|_p < \infty\}$. If $1 \leq p \leq \infty$, A_p is then a two-sided *-ideal of A (proper only if p < 2), and $(A_p, |\cdot|_p)$ is a normed *-algebra. $(A_2, |\cdot|_2)$ is $(A, ||\cdot||)$; and for $1 \leq p < 2$, $(A_p, |\cdot|_p)$ is a Banach *-algebra, for which structure theorems are given.

1. Introduction. Let A be a semisimple H^* -algebra with inner product and norm denoted by (,) and $||\cdot||$, respectively. The trace class of A, that is, the set $\tau(A) = \{xy: x, y \in A\}$, has been studied by Saworotnow and Friedell [8], who show, first of all, that for any nonzero $a \in A$ there exists a positive element $[a] \in A$ such that $[a]^2 =$ a^*a , and $a \in \tau(A)$ if and only if $[a] \in \tau(A)$. An algebra norm τ is then introduced on $\tau(A)$ by defining $\tau(a) = tr[a]$ for each $a \in \tau(A)$, where in turn the trace functional tr is unambiguously defined on $\tau(A)$ by letting $tr xy = (x, y^*) = \Sigma(xyp_{\omega}, p_{\omega}), \{p_{\omega}: \omega \in \Omega\}$ being any maximal family of mutually orthogonal nonzero self-adjoint idempotents. With this norm, $\tau(A)$ is actually a Banach algebra [9, Corollary to Theorem 1]. This presentation parallels that of Schatten [10] for τc , the trace class of σc , the Schmidt class of operators on a Hilbert space.

In a somewhat similar sense our central development in §3 brings over into the present context some of the work of McCarthy [6] on the operator algebras c_p . We preface this with a basic spectral theorem established in §2; in §4 we study the structure of the Banach *-algebras A_p , where $1 \leq p < 2$. Finally, in §5 we relate A_p to the class c_p of operators on a Hilbert space [6; 2, ch. XI. 9] and also to \mathcal{C}_p spaces [3, pp. 70 ff.; 5].

2. Preliminary spectral theory. Throughout the remainder of this paper A will continue to denote a semisimple H^* -algebra. By a projection p in A we shall mean a nonzero self-adjoint idempotent. A projection p is primitive if p cannot be expressed as $p = p_1 + p_2$, where p_1 and p_2 are orthogonal projections. By a projection base in A we mean a maximal family of mutually orthogonal projections (not necessarily primitive); note that if $a \in A$ and $\{p_{\omega}: \omega \in \Omega\}$ is a projection base, then $a = \sum a p_{\omega} = \sum p_{\omega} a$ [1, Theorem 4.1, where primitivity of the projections is not needed to establish this point]. Finally, we shall say that an element a in A is positive if $(ax, x) \ge 0$ for every $x \in A$; a is then necessarily self-adjoint.

LEMMA 2.1. Let b be a nonzero normal element of A. There is a well-defined family $\{p_{\omega}: \omega \in \Omega\}$ of mutually orthogonal projections in A, and a well-defined set $\{\alpha_{\omega}: \omega \in \Omega\}$ of complex numbers, such that

 $(1) \quad b = \Sigma \alpha_{\omega} p_{\omega}$

(2) $bp_{\omega} = p_{\omega}b = \alpha_{\omega}p_{\omega}$ for each $\omega \in \Omega$.

The nonzero α_{ω} are precisely the nonzero elements of the spectrum of b.

Proof. Let A_0 be the intersection of all maximal commutative *-subalgebras of A containing b. A_0 is a proper H*-algebra in the inner product and involution of A. Let $\{p_{\omega}: \omega \in \Omega\}$ be the collection of projections of A_0 which are primitive in A_0 ; then each $p_{\omega}A_0$ is a minimal ideal of A_0 , and if $\omega_1 \neq \omega_2$ we have $p_{\omega_1} p_{\omega_2} = 0$ and $(p_{\omega_1}, p_{\omega_2}) = 0$. Also, $A_0 = \Sigma p_{\omega} A_0$, the orthogonal direct sum of the minimal ideals $p_{\omega}A_0$, each of which is one-dimensional and consists of scalar multiples of p_{ω} [1, Corollary 4.1]. Therefore $b = \sum \alpha_{\omega} p_{\omega}$, where $\{\alpha_{\omega} : \omega \in \Omega\}$ is a set of complex numbers. Property (2) is immediate from the orthogonality of the p_{ω} . We shall show that the nonzero α_{ω} are the nonzero elements of $sp(b|A_0)$, the spectrum of b relative to A_0 . Let ϕ be any multiplicative linear functional on A_0 . We have $\phi(p_{\omega}) = \phi(p_{\omega}^2) = [\phi(p_{\omega})]^2$, and hence the value of ϕ at each projection p_{ω} must be either 0 or 1. ϕ cannot have the value 0 at every p_{ω} or else ϕ would vanish on A_0 ; nor can we have $\phi(p_{\omega_1}) = 1 = \phi(p_{\omega_2})$ if $\omega_1 \neq \omega_2$, for then 1 = $\phi(p_{\omega_1})\phi(p_{\omega_2}) = \phi(p_{\omega_1}p_{\omega_3}) = \phi(0) = 0.$ Therefore, each multiplicative linear functional on A_{\circ} is of the form $\phi_{\nu}(p_{\omega}) = \delta_{\nu\omega}$, where $\nu \in \Omega$. We have, for each $\nu \in \Omega$, $\phi_{\nu}(b) = \Sigma_{\omega} \alpha_{\omega} \phi_{\nu}(p_{\omega}) = \alpha_{\nu} = b(\phi_{\nu})$, where b denotes the Gelfand transform of $b \in A_0$. Since the nonzero α_{ω} are therefore the nonzero elements of the range of b, they are by the Gelfand theory precisely the nonzero elements of $sp(b|A_0)$. However, $sp(b|A) = sp(b|A_0)$, since if $c \in A_0$ has a quasi-inverse c^0 in A, then, as is well-known, c^0 belongs to every maximal commutative *-subalgebra of A containing c, or equivalently, $c^0 \in A_0$. Finally, it is clear that the element b uniquely determines the algebra A_0 , along with its set of primitive projections $\{p_{\omega}: \omega \in \Omega\}$ and the corresponding numbers α_{ω} , since $\alpha_{\omega}p_{\omega}$ is the orthogonal projection of b on the closed ideal $p_{\omega}A_0$ of A_0 .

LEMMA 2.2. Let b be a nonzero normal element of A, and let $b = \Sigma \mu_n q_n$, where $\{q_n\}$ is a countable (possibly finite) family of mutually orthogonal projections, and the μ_n are nonzero complex numbers such that $\mu_m \neq \mu_n$ if $m \neq n$. Let h be any self-adjoint element of A which commutes with b. Then for each n, $hq_n = q_nh$.

Proof. Extend $\{q_n\}$ to a projection base $\{q_i: \gamma \in \Gamma\}$. For each γ , if $q_{\gamma} = q_n$ for some *n*, let $\mu_{\gamma} = \mu_n$; otherwise, let $\mu_{\gamma} = 0$. (Note that $bq_{\gamma} = q_{\gamma}b = \mu_{\gamma}q_{\gamma}$ for each $\gamma \in \Gamma$.) Then for any q_n we have $q_nh =$

 $\Sigma_{\tau}q_nhq_{\tau}$. Also, since b and h commute, $\mu_nq_nhq_{\tau} = q_nbhq_{\tau} = q_nhbq_{\tau} = \mu_{\tau}q_nhq_{\tau}$. If $q_{\tau} \neq q_n$ then $\mu_{\tau} \neq \mu_n$ and consequently $q_nhq_{\tau} = 0$. Thus $q_nh = q_nhq_n$. Taking adjoints we have $hq_n = q_nhq_n$; therefore $hq_n = q_nh$.

COROLLARY 2.3. Let b, $\{\mu_n\}$, and $\{q_n\}$ be as in the lemma, and let A_0 be, as before, the intersection of all maximal commutative *subalgebras of A containing b. Then for each $n, q_n \in A_0$.

Proof. Let A_1 be any maximal commutative *-subalgebra of A containing b. Since A_1 is a *-algebra, each $x \in A_1$ is of the form x = h + ik, where $h, k \in A_1$, and h and k are self-adjoint. Therefore, each q_n commutes with every element of A_1 , and by maximality of $A_1, q_n \in A_1$. Therefore, finally, $q_n \in A_0$.

LEMMA 2.4. Let b, $\{\mu_n\}$, and $\{q_n\}$ be as in Lemma 2.2. Then each q_n is a finite sum of the projections p_{ω} of Lemma 2.1.

Proof. Each q_n belongs to A_0 , and therefore, as in the proof of Lemma 2.1, $q_n = \Sigma \beta_w p_w$ for suitable numbers β_w . Also, $q_n = q_n^2 = \Sigma \beta_w^2 p_w$, and therefore each β_w is either 0 or 1. Only finitely many can be 1, since $||q_n||^2 = \Sigma \beta_w^2 ||p_w||^2 \ge \Sigma \beta_w^2$.

Now let $q_n = p_{n_1} + \cdots + p_{n_{k(n)}}$. The orthogonal projection of bon the closed left ideal Aq_n is $bq_n = \mu_n q_n = \mu_n (p_{n_1} + \cdots + p_{n_{k(n)}})$. From Lemma 2.1, since $b = \Sigma \alpha_{\omega} p_{\omega}$, this projection of b is also $\alpha_{n_1} p_{n_1} + \cdots + \alpha_{n_{k(n)}} p_{n_{k(n)}}$. Therefore $\alpha_{n_i} = \mu_n$, $i = 1, \cdots + k(n)$, and in the representation $b = \Sigma \alpha_{\omega} p_{\omega}$ we may replace the sum $\alpha_{n_1} p_{n_1} + \cdots + \alpha_{n_{k(n)}} p_{n_{k(n)}}$ by $\mu_n q_n$. If this is done for each n indexing the countable set $\{q_n\}$, the procedure evidently replaces the representation $b = \Sigma \alpha_{\omega} p_{\omega}$ by $b = \Sigma \mu_n q_n$, and therefore makes use of every term $\alpha_{\omega} p_{\omega}$ except those for which $\alpha_{\omega} = 0$. We thus have the following spectral theorem.

THEOREM 2.5. Let b be a nonzero normal element of A. Then b may be represented uniquely (apart from the order of the terms) as a sum

$$(*) \qquad \qquad b=\varSigma\lambda_n e_n$$
 ,

in which

(1) $\{\lambda_n\}$ is a countable family of distinct nonzero complex numbers consisting of the nonzero elements of the spectrum of b, and

(2) $\{e_n\}$ is a countable family of mutually orthogonal projections. We have $be_n = e_n b = \lambda_n e_n$ for each n; b is self-adjoint if and only if each λ_n is real, and b is positive if and only if each $\lambda_n > 0$.

DEFINITION 2.6. Let b be a nonzero normal element of A. A representation (*) of b having properties (1) and (2) of Theorem 2.5

JAMES F. SMITH

will be called a spectral representation of b. If b is a positive element of A, we shall refer to the spectral representation of b, meaning the one in which $\lambda_m < \lambda_n$ if m > n. For any nonzero normal element b, the set E_b of mutually orthogonal projections in a spectral representation of b will be called the spectral family of b.

DEFINITION 2.7. Let b be a nonzero normal element of A, and let E_b be its spectral family. A projection base $\{e_{\omega}: \omega \in \Omega\}$ containing every e_n in E_b will be called a *projection base associated with b*. (Note that by a simple maximality argument, E_b can always be extended to a projection base associated with b.)

The classes A_{p} and their basic properties. We begin this 3. section by recalling some basic results from [8]. Corresponding to each a in A there is a unique positive element [a] of A such that $[a]^2 = a^*a$. Moreover, there is, for each nonzero a in A, a well-defined partial isometry W on A, having initial set $\overline{[a]A}$ and final set \overline{aA} , such that a = W[a], $[a] = W^*a$, and ||W|| = 1. We shall call W the partial isometry associated with a. We define a left centralizer on A to be an operator S in B(A) such that S(xy) = (Sx)y for all $x, y \in A$. (This terminology, though widely used, is not universal; the type of operator just defined is called a right centralizer in [8] and [9], and elsewhere.) Evidently, each left multiplication operator L_a , $a \in A$, is a left centralizer on A; also, for any nonzero a in A, the partial isometry W associated with a is a left centralizer (see [8, p. 97]). We note, finally, for fairly frequent use, that for any $x \in A$, ||ax|| =||[a]x||, since $||ax||^2 = (ax, ax) = (a^*ax, x) = ([a]^2x, x) = ([a]x, [a]x) =$ $||[a]x||^2$.

DEFINITION 3.1. Let a be a nonzero element of A, and let $[a] = \Sigma \lambda_n e_n$ be the spectral representation of [a]. We define

$$\|a\|_p = (\Sigma\lambda_n^p \||e_n||^2)^{1/p} ext{ for } 0 , $\|a\|_\infty = \lambda_1.$$$

For a = 0, we define $|a|_p = 0$, 0 .

DEFINITION 3.2. For $0 , <math>A_p = \{a \in A \colon |a|_p < \infty\}$.

Remark 3.3. For 0 ,

(1) $a \in A_p$ if and only if $[a] \in A_p$, since [a] = [[a]] implies $|a|_p = |[a]|_p$;

(2) if e is a projection, $e \in A_p$ and $|e|_p = ||e||^{2/p}$.

REMARK 3.4. Let $\{e_{\omega}: \omega \in \Omega\}$ be a projection base associated with [a]. We shall write $[a] = \Sigma \lambda_{\omega} e_{\omega}$, always assuming that $\lambda_{\omega} = \lambda_n$ if $e_{\omega} \notin E_{[a]}$. Then $|a|_p = (\Sigma \lambda_{\omega}^p ||e_{\omega}||^2)^{1/2}$ for $0 ; and we continue to write <math>|a|_{\infty} = \lambda_1$, understanding λ_1 to be $\sup \{\lambda_{\omega}: \omega \in \Omega\}$.

REMARK 3.5. Let $\{e_{\omega}: \omega \in \Omega\}$ be a projection base associated with $[a] \in A$.

(1) $|a|_{2}^{2} = |[a]|_{2}^{2} = \Sigma \lambda_{\omega}^{2} ||e_{\omega}||^{2} = \Sigma ||\lambda_{\omega}e_{\omega}||^{2} = \Sigma ||[a]e_{\omega}||^{2} = \Sigma ||ae_{\omega}||^{2} = ||a||^{2}$. Hence $|a|_{2} = ||a||$ and $A_{2} = A$.

(2) $|a|_1 = |[a]|_1 = \Sigma \lambda_{\omega} ||e_{\omega}||^2 = \Sigma (\lambda_{\omega} e_{\omega}, e_{\omega}) = \Sigma ([a] e_{\omega}, e_{\omega}) = tr [a] = \tau(a)$ [8, Lemma 3]. Hence $|a|_1 = \tau(a)$ and $A_1 = \tau(A)$, the trace class of A.

DEFINITION 3.6. Let b be a nonzero positive element of A, with spectral representation $b = \Sigma \lambda_n e_n$. For $0 , <math>b^p = \Sigma \lambda_n^p e_n$, provided that this sum exists in A.

REMARK 3.7. From [8, Lemma 3] we have that $a \in A_p$ if and only if $[a]^p \in A_1 = \tau(A)$. This occurs if and only if $[a]^{p/2}$ exists in A; we then have $|a|_p^p = \Sigma \lambda_n^p ||e_n||^2 = \tau([a]^p) = |[a]^p|_1 = ||[a]^{p/2}||^2 = \Sigma([a]^p p_{\omega}, p_{\omega})$ for any projection base $\{p_{\omega} : \omega \in \Omega\}$.

REMARK 3.8. For $0 , clearly <math>|\alpha|_p \geq 0$, and $|\alpha|_p = 0$ if and only if a = 0. Also, since $[\alpha a] = |\alpha| [a]$ for any complex number α , we have $|\alpha a|_p = |\alpha| |a|_p$.

LEMMA 3.9. For any $a \in A$ and $0 , <math>|a|_{\infty} \leq |a|_{p}$.

Proof. For a = 0 the result is obvious. Otherwise, using the spectral representation of [a], we have $|a|_{\infty}^{p} = \lambda_{1}^{p} \leq \Sigma \lambda_{n}^{p} ||e_{n}||^{2} = |a|_{p}^{p}$.

LEMMA 3.10. For any $a \in A$, $||ax|| \leq |a|_{\infty} ||x||$.

Proof. For $a \neq 0$, let $\{e_{\omega} \colon \omega \in \Omega\}$ be a projection base associated with [a]. Then $[a]x = \Sigma \lambda_{\omega} e_{\omega} x$ and $||[a]x||^2 = \Sigma \lambda_{\omega}^2 ||e_{\omega} x||^2 \leq \lambda_1^2 \Sigma ||e_{\omega} x||^2 = \lambda_1^2 ||x||^2$. Hence $||ax|| = ||[a]x|| \leq |a|_{\infty} ||x||$.

COROLLARY 3.11. For any $a \in A$, $|a|_{\infty} = ||L_a||$.

Proof. For $a, x \neq 0$, $||ax||/||x|| \leq |a|_{\infty}$, by the lemma. But $||ae_1||/||e_1|| = ||[a]e_1||/||e_1|| = \lambda_1 = |a|_{\infty}$.

PROPOSITION 3.12. For $a \in A$ and $0 , <math>|a|_q \leq |a|_p$.

Hence $A_p \subset A_q$, and if $2 \leq p \leq \infty$ then $A_p = A$.

Proof. Using the spectral representation of [a], we have $|a|_q^q = \Sigma \lambda_n^q ||e_n||^2 = \Sigma \lambda_n^{q-p} \lambda_n^p ||e_n||^2 \leq \lambda_1^{q-p} \Sigma \lambda_n^p ||e_n||^2 = |a|_{\infty}^{q-p} |a|_p^p \leq |a|_p^q$, by Lemma 3.9.

REMARK 3.13. By 3.7, $a \in A_{2p}$ $(0 if and only if <math>[a]^p$ exists in A. For $1 \leq p < \infty$, $A_{2p} = A$ and hence $[a]^p$ is defined.

PROPOSITION 3.14. If A is infinite-dimensional, then for $0 , <math>A_q$ is properly larger than A_p .

Proof. From the structure theory of H^* -algebras [1], we see that if A is infinite-dimensional then A contains a countably infinite set $\{e_n: n \in N\}$ of mutually orthogonal projections. Choose r such that p < r < q; then the series $\sum_{n=1}^{\infty} n^{-1/r} ||e_n||^{-2/q} e_n$ converges to a positive element of A (since the squares of the norms of its terms have a finite sum). Denoting this element by a, we observe that the given series (or one obtained from it by grouping and rearranging terms) is the spectral representation of a. Thus $a \in A_q$, since $|a|_q^q = \sum_{n=1}^{\infty} n^{-q/r} < \infty$; however $a \notin A_p$, since $|a|_p^p = \sum_{n=1}^{\infty} n^{-p/r} ||e_n||^{2-(2p/q)} \ge \sum_{n=1}^{\infty} n^{-p/r} = \infty$.

Some elements of the following lemma appear in [8, p. 96]. For most of it, however, the author is indebted to M. Kervin.

LEMMA 3.15. Let a be any nonzero element of A, and let $[a] = \Sigma \lambda_n e_n$ be the spectral representation of [a]. For each n, let $f_n = \lambda_n^{-2} a e_n a^*$. Then $[a^*] = \Sigma \lambda_n f_n$ is the spectral representation of $[a^*]$, and $||f_n|| = ||e_n||$ for each n.

Proof. Clearly, the λ_n are distinct positive numbers and the f_n are self-adjoint. We recall, first of all, that $[a]^2 = \Sigma \lambda_n^2 e_n = a^* a$, and therefore $a^* a e_n = e_n a^* a = \lambda_n^2 e_n$. Thus $f_m f_n = (\lambda_m^{-2} a e_m a^*)(\lambda_n^{-2} a e_n a^*) = \lambda_n^{-2} \lambda_n^{-2} a e_m (a^* a e_n) a^* = \lambda_m^{-2} a e_m e_n a^* = \delta_{mn} f_n$. Therefore, the f_n are mutually orthogonal idempotents. Also, $\lambda_n^2 ||f_n||^2 = \lambda_n^{-2} (a e_n a^*, a e_n a^*) = (e_n a^*, e_n a^*) = \lambda_n^2 ||e_n||^2$, and therefore $||f_n|| = ||e_n||$ and the f_n are nonzero. Now we wish to show that $[a^*] = \Sigma \lambda_n f_n$. We shall show first that $a = \Sigma a e_n$. Extend the family $E_{[a]}$ to a projection base $\{e_\omega : \omega \in \Omega\}$. Then $a = \Sigma a e_\omega$ and $a^* a = \Sigma a^* a e_\omega$. But if $e_\alpha \notin E_{[a]}$ then $a^* a e_\alpha = 0$, since $a^* a = \Sigma \lambda_n^2 e_n = \Sigma a^* a e_n$. Therefore, for $e_\alpha \notin E_{[a]}$ we have $e_\alpha a^* a e_\alpha = 0 = (a e_\alpha)^* (a e_\alpha)$, and thus $a e_\alpha = 0$ [1, Lemma 2.2]. We conclude that $a = \Sigma a e_n$. Finally, $(\Sigma \lambda_n f_n)^2 = \Sigma \lambda_n^2 f_n = \Sigma a e_n a^* = a a^*$, and therefore $\Sigma \lambda_n f_n$ is the (unique) positive square root of $a a^*$; that is, $\Sigma \lambda_n f_n = [a^*]$.

782

COROLLARY 3.16. For any $a \in A$ and $0 , <math>|a|_p = |a^*|_p$. Hence $a \in A_p$ if and only if $a^* \in A_p$.

In order to arrive at the results announced in our opening synopsis, we shall need to establish several crucial inequalities. Lemmas 3.17, 3.18, and 3.22 are adapted from [6, Lemmas 2.1, 2.2].

LEMMA 3.17. For $0 , let b be a positive element of <math>A_{2p}$ (so that b^p exists in A). Then for any nonzero $x \in A$,

 $(1) \quad (b^p x, x) \ge (bx, x)^p ||x||^{2(1-p)} \ if \ 1 \le p < \infty$,

 $(2) \quad (b^{p}x, x) \leq (bx, x)^{p} ||x||^{2(1-p)} \text{ if } 0$

Proof. (1) Suppose $1 \leq p < \infty$. Let $\{e_{\omega} : \omega \in \Omega\}$ be a projection base associated with b, where, as usual, we take $\lambda_{\omega} = \lambda_n$ if $e_{\omega} = e_n \in E_b$, and $\lambda_{\omega} = 0$ if $e_{\omega} \notin E_b$. We have, by Hölder's inequality,

$$egin{aligned} (bx,\,x) &= \Sigma\lambda_\omega(e_\omega x,\,x) \ &\leq [\Sigma\lambda_\omega^p(e_\omega x,\,x)]^{1/p}[\Sigma(e_\omega x,\,x)]^{1-(1/p)} \ &= [(\Sigma\lambda_\omega^pe_\omega x,\,x)]^{1/p}[\Sigma\,||\,e_\omega x\,||^2]^{(p-1)/p} \ &= (b^p x,\,x)^{1/p}\,||\,x\,||^{2(p-1)/p} \ . \end{aligned}$$

Hence $(b^{p}x, x) \ge (bx, x)^{p} ||x||^{2(1-p)}$.

(2) Suppose $0 . Replace the element b in (1) by <math>b^p$ and the exponent p by 1/p to obtain the desired inequality.

LEMMA 3.18. Let $a \in A$, and let $\{q_{\omega} : \omega \in \Omega\}$ be a projection base for A. Then

 $(1) |a|_p^p \leq \Sigma ||aq_w||^p ||q_w||^{2-p} ext{ if } 1 \leq p \leq 2$,

 $(2) |a|_p^p \geq \Sigma ||aq_\omega||^p ||q_\omega||^{2-p} \text{ if } 2 \leq p < \infty.$

In each case, equality holds if $\{q_{\omega} : \omega \in \Omega\}$ is a projection base associated with [a].

Proof. We note first that $[a]^p$ exists, since $p \ge 1$.

(1) Suppose $1 \leq p \leq 2$. By (2) of Lemma 3.17 we have for each q_{ω} ,

$$egin{aligned} ([a]^p q_{\omega}, q_{\omega}) &= (([a]^2)^{p/2} q_{\omega}, q_{\omega}) \ &\leq ([a]^2 q_{\omega}, q_{\omega})^{p/2} ||q_{\omega}||^{2-p} \ &= ||aq_{\omega}||^p ||q_{\omega}||^{2-p} \ . \end{aligned}$$

Summing over Ω gives, by 3.7,

$$a|_p^p = \varSigma([a]^p q_w, q_w) \leq \varSigma ||aq_w||^p ||q_w||^{2-p}$$
 .

If $\{q_{\omega}\}$ is a projection base associated with [a], then by 3.4 we have

$$\begin{split} \Sigma ||aq_{\omega}||^{p}||q_{\omega}||^{2-p} &= \Sigma ||[a]q_{\omega}||^{p}||q_{\omega}||^{2-p} \\ &= \Sigma \lambda_{\omega}^{p}||q_{\omega}||^{p}||q_{\omega}||^{2-p} \\ &= \Sigma \lambda_{\omega}^{p}||q_{\omega}||^{2} \\ &= |a|_{p}^{p} . \end{split}$$

(2) is proved similarly, using (1) of Lemma 3.17.

PROPOSITION 3.19. For $1 \leq p \leq \infty$, let $a \in A_p$, and let S be a left centralizer on A. Then $Sa \in A_p$, and $|Sa|_p \leq ||S|| |a|_p$.

Proof. The result is standard for $p = \infty$. Suppose $1 \le p \le 2$; let $\{e_{\omega}: \omega \in \Omega\}$ be a projection base associated with [a]. By Lemma 3.18 (1), $|Sa|_p^p \le \Sigma||(Sa)e_{\omega}||^p||e_{\omega}||^{2-p} = \Sigma||S(ae_{\omega})||^p||e_{\omega}||^{2-p} \le ||S||^p\Sigma||ae_{\omega}||^p||e_{\omega}||^{2-p} =$ $||S||^p|a|_p^p$. Now suppose $2 \le p < \infty$, and this time let $\{e_{\omega}: \omega \in \Omega\}$ be a projection base associated with [Sa]. We have, using (2) of Lemma 3.18, $|Sa|_p^p = \Sigma||(Sa)e_{\omega}||^p||e_{\omega}||^{2-p} = \Sigma||S(ae_{\omega})||^p||e_{\omega}||^{2-p} \le ||S||^p\Sigma||ae_{\omega}||^p||e_{\omega}||^{2-p} \le$ $||S||^p|a|_p^p$.

COROLLARY 3.20. For $1 \leq p \leq \infty$, let $a \in A_p$, $x \in A$. Then xa and ax belong to A_p , and $|xa|_p \leq |x|_{\infty} |a|_p$, $|ax|_p \leq |a|_p |x|_{\infty}$.

Proof. By Corollary 3.11 the statements about xa are immediate, since L_x is a left centralizer. We also have, by Corollary 3.16, $|ax|_p = |(ax)^*|_p = |x^*a^*|_p \le |x^*|_{\infty} |a^*|_p = |a|_p |x|_{\infty}$.

COROLLARY 3.21. For $1 \leq p \leq \infty$, let $a, b \in A_p$. Then $|ab|_p \leq |a|_p |b|_p$.

In our next lemma we shall make use of a special operator decomposition given by McCarthy [6, p. 250]. Suppose $T \in B(A)$; then $T = (TT^*)^{1/4}U(T^*T)^{1/4}$, where U is a partial isometry with ||U|| = 1.

LEMMA 3.22. Suppose $1 \leq p < \infty$. Let $a \in A$, and let $\{q_{\omega} : \omega \in \Omega\}$ be any projection base for A. Then $\Sigma |(aq_{\omega}, q_{\omega})|^{p} ||q_{\omega}||^{2(1-p)} \leq |a|_{p}^{p}$.

Proof. We use the operator decomposition just mentioned: $L_a = (L_a L_a^*)^{1/4} U(L_a^* L_a)^{1/4} = L_{[a^*]}^{1/2} U L_{[a]}^{1/2}$. We have, by two applications of the Schwarz inequality,

$$\begin{split} & \Sigma \left| \left(a q_{\omega}, q_{\omega} \right) |^{p} \left| \left| q_{\omega} \right| \right|^{2(1-p)} = \Sigma \left| \left(U L_{[a]}^{1/2} q_{\omega}, L_{[a*]}^{1/2} q_{\omega} \right) |^{p} \right| \left| q_{\omega} \right| \right|^{2(1-p)} \\ & \leq \Sigma \left| \left| L_{[a]}^{1/2} q_{\omega} \right| |^{p} \left| \left| L_{[a*]}^{1/2} q_{\omega} \right| |^{p} \right| \left| q_{\omega} \right| \right|^{2(1-p)} \\ & = \Sigma \left(\left| \left| L_{[a]}^{1/2} q_{\omega} \right| \right|^{p} \left| \left| q_{\omega} \right| \right|^{1-p} \right) \left(\left| \left| L_{[a*]}^{1/2} q_{\omega} \right| \right|^{p} \right| \left| q_{\omega} \right| \right|^{1-p} \right) \\ & \leq \left[\Sigma \left| \left| L_{[a]}^{1/2} q_{\omega} \right| \right|^{2p} \left| \left| q_{\omega} \right| \right|^{2(1-p)} \right]^{1/2} \left[\Sigma \left| \left| L_{[a*]}^{1/2} q_{\omega} \right| \right|^{2p} \left| \left| q_{\omega} \right| \right|^{2(1-p)} \right]^{1/2} \end{split}$$

$$= \left[\Sigma(L_{[a]}^{1/2}q_{\omega}, L_{[a]}^{1/2}q_{\omega})^{p} || q_{\omega} ||^{2(1-p)} \right]^{1/2} \left[\Sigma(L_{[a^{*}]}^{1/2}q_{\omega}, L_{[a^{*}]}^{1/2}q_{\omega})^{p} || q_{\omega} ||^{2(1-p)} \right]^{1/2} \\= \left[\Sigma([a]q_{\omega}, q_{\omega})^{p} || q_{\omega} ||^{2(1-p)} \right]^{1/2} \left[\Sigma([a^{*}]q_{\omega}, q_{\omega})^{p} || q_{\omega} ||^{2(1-p)} \right]^{1/2} \\\leq \left[\Sigma([a]^{p}q_{\omega}, q_{\omega}) \right]^{1/2} \left[\Sigma([a^{*}]^{p}q_{\omega}, q_{\omega}) \right]^{1/2} \text{ by Lemma 3.17 (1)} \\= |a|_{p}^{p/2} |a^{*}|_{p}^{p/2} \text{ by 3.7} \\= |a|_{p}^{p}.$$

PROPOSITION 3.23. For $1 \leq p \leq \infty$, let $a, b \in A_p$. Then $|a + b|_p \leq |a|_p + |b|_p$.

Proof. The result is well-known for $p = \infty$. For $1 \leq p < \infty$, let $\{e_{\omega}: \omega \in \Omega\}$ be a projection base associated with [a + b], and let W be the partial isometry associated with a + b. Then

$$\begin{split} |a + b|_{p} &= [\Sigma([a + b]e_{\omega}, e_{\omega})^{p} ||e_{\omega}||^{2(1-p)}]^{1/p} \\ &= [\Sigma|([a + b]e_{\omega}, e_{\omega})|^{p} ||e_{\omega}||^{2(1-p)}]^{1/p} \\ &= [\Sigma|((W^{*}(a + b))e_{\omega}, e_{\omega})|^{p} ||e_{\omega}||^{2(1-p)}]^{1/p} \\ &= [\Sigma|((W^{*}a)e_{\omega}, e_{\omega})||e_{\omega}||^{2(1-p)/p} + ((W^{*}b)e_{\omega}, e_{\omega})||e_{\omega}||^{2(1-p)/p}|^{p}]^{1/p} \\ &\leq [\Sigma|((W^{*}a)e_{\omega}, e_{\omega})|^{p} ||e_{\omega}||^{2(1-p)}]^{1/p} + [\Sigma|((W^{*}b)e_{\omega}, e_{\omega})|^{p} ||e_{\omega}||^{2(1-p)}]^{1/p} \end{split}$$

by Minkowski's inequality

 $\leq |W^*a|_p + |W^*b|_p$ by Lemma 3.22 $\leq ||W^*|| |a|_p + ||W^*|| |b|_p$ by Proposition 3.19 $= |a|_p + |b|_p.$

COROLLARY 3.24. For $1 \leq p \leq \infty$, A_p is a normed linear space. Hence A_p is a two-sided *-ideal of A and $(A_p, |\cdot|_p)$ is a normed algebra.

Now for $1 \leq p \leq \infty$ we wish to investigate the relationship between A_p and the dual space of A_q , where (1/p) + (1/q) = 1. In what follows we shall omit proofs for the cases $p = 1, q = \infty$ and $p = \infty, q = 1$; these are given in [9].

LEMMA 3.25. Let (1/p) + (1/q) = 1, where $1 \leq p, q \leq \infty$. Let $a \in A_p$, $b \in A_q$. Then $|tr \ ab| = |tr \ ba| \leq |a|_p |b|_q$.

Proof. We shall assume with no loss of generality that 1 $and hence <math>2 \leq q < \infty$. Let $\{e_{\omega}: \omega \in \Omega\}$ be a projection base associated with [a]. Then $|tr ab| = |tr ba| = |\Sigma(bae_{\omega}, e_{\omega})| \leq \Sigma |(ae_{\omega}, b^*e_{\omega})| \leq \Sigma ||ae_{\omega}|| ||b^*e_{\omega}|| = \Sigma ||ae_{\omega}|| ||e_{\omega}||^{(2-p)/p}||b^*e_{\omega}|| ||e_{\omega}||^{(2-q)/q}$, since ((2-p)/p) + ((2-q)/q) = 0. By Hölder's inequality, the last sum does not exceed $|\Sigma||ae_{\omega}||^p ||e_{\omega}||^{2-p}]^{1/p} [\Sigma||b^*e_{\omega}||^q ||e_{\omega}||^{2-q}]^{1/q}$. But the first sum in brackets is $|a|_p^p$, and the second is less than or equal to $|b^*|_q^q$, by Lemma 3.18 (2). Hence $|tr ab| \leq |a|_p |b|_q$.

For each $a \in A_p$ we now define $\phi_a(x) = tr \ xa$ for all $x \in A_q$. From the linearity of tr on the trace class $\tau(A)$, it is evident that ϕ_a is a linear functional on A_q ; moreover, ϕ_a is bounded and $||\phi_a|| \leq |a|_p$, by Lemma 3.25. We shall show that the opposite inequality holds as well.

PROPOSITION 3.26. For $1 \leq p \leq \infty$, the mapping $a \to \phi_a$ is a linear isometry of A_p into A'_q , the dual space of A_q .

Proof. Again using the linearity of tr on $\tau(A)$ one easily verifies that the mapping is linear. In view of our above remarks, therefore, we need only prove that $|a|_p \leq ||\phi_a||$. Let $[a] = \Sigma \lambda_n e_n$ be the spectral representation of [a], and let $w_k = \Sigma_{n=1}^k \lambda_n^{p-2} e_n a \in A_q$. We shall compute $|w_k|_q$. First of all, $w_k^* w_k = (\sum_{m=1}^k \lambda_n^{p-2} a e_m)(\sum_{n=1}^k \lambda_n^{p-2} e_n a^*) = \sum_{m,n=1}^k \lambda_n^{p-2} \lambda_n^{p-2} a e_m a^* = \sum_{n=1}^k \lambda_n^{2(p-1)} a e_n a^* = \sum_{n=1}^k \lambda_n^{2(p-1)} \lambda_n^{-2} a e_n a^* = \sum_{n=1}^k \lambda_n^{2(p-1)} f_n$, where $f_n = \lambda_n^{-2} a e_n a^*$. Since, by Lemma 3.15, the f_n are mutually orthogonal projections with $||f_n|| = ||e_n||$, we have $[w_k] = \sum_{n=1}^k \lambda_n^{p-1} f_n$, and $|w_k|_q = [\sum_{n=1}^k \lambda_n^{q(p-1)} ||f_n||^2]^{1/q} = [\sum_{n=1}^k \lambda_n^{p} ||e_n||^2]^{1/q}$. We also have $\sum_{n=1}^k \lambda_n^{p-1} f_n$, and $|w_k|_q = ||\phi_a|| [\sum_{n=1}^k \lambda_n^{p-1} \lambda_n^{p-1} e_n a^*] = |tr(\sum_{n=1}^k \lambda_n^{p-1} e_n a^* a)| = |tr w_k a| = |\phi_a(w_k)| \leq ||\phi_a|| ||w_k|_q = ||\phi_a|| [\sum_{n=1}^k \lambda_n^{p-1} ||e_n||^2]^{1/q}$. Thus $[\sum_{n=1}^k \lambda_n^{p-1} ||e_n||^2]^{1/p} \leq ||\phi_a||$, and since $\sum_{n=1}^k \lambda_n^{p-1} ||e_n||^2 \leq ||\phi_a||^p$ for every k, we have $|a|_p^p \leq ||\phi_a||^p$.

THEOREM 3.27. For $1 \leq p \leq 2$, the mapping $a \rightarrow \phi_a$ is a linear isometry of A_p onto A'_q .

Proof. Let ϕ be any bounded linear functional on A_q . Then for all $x \in A_q(=A)$, $|\phi(x)| \leq ||\phi|| ||x|_q \leq ||\phi|| ||x||$, by Proposition 3.12. Therefore ϕ is a bounded linear functional on A, and by the Riesz representation theorem there exists $a \in A$ such that $\phi(x) = (x, a^*) =$ $tr \, xa$ for all $x \in A$. We need only show that $a \in A_p$. But if we again consider the spectral representation $[a] = \Sigma \lambda_n e_n$ and define w_k as in the preceding proof, the same computations show that $\sum_{n=1}^k \lambda_n^n ||e_n||^2 \leq$ $||\phi||^p$ for every k, and hence $\Sigma \lambda_n^n ||e_n||^2 < \infty$ and $a \in A_p$.

COROLLARY 3.28. For $1 \leq p \leq 2$, $(A_p, |\cdot|_p)$ is a Banach *-algebra.

We conclude this section with an example to show that if $2 and <math>A(=A_p)$ is infinite-dimensional, then $(A^p, |\cdot|_p)$ is incomplete. First of all, if $(A_p, |\cdot|_p)$ is complete, then from the inverse mapping theorem and the fact that $|\cdot|_p$ is dominated by $||\cdot||$, we can conclude that these two norms are equivalent on A. But this is not so if A is infinite-dimensional, for if $\{e_n: n \in N\}$ is a countably infinite set of mutually orthogonal projections in A and we let $s_k = \sum_{n=1}^k n^{-1/2} ||e_n||^{-2/p} e_n$, then $\{s_k\}$ is a Cauchy sequence in the $|\cdot|_p$ -topology but not in the

786

||•||-topology.

4. The structure of the Banach *-algebras A_p . In this section we shall confine our attention mainly to the algebras A_p , where $1 \leq p \leq 2$, although some of our results hold for p > 2 as well. Unless otherwise indicated, therefore, we shall assume throughout that $1 \leq p < 2$. We begin by observing that for these vealues of p, A_p is a quite special instance of an *IP*-algebra, as introduced and studied by Yood in [12]; hence the entire theory of that paper is at our disposal. Furthermore, it is readily verified that $(A_p, ||\cdot||)$ is a (normed) Hilbert algebra; we shall immediately note some properties of this Hilbert algebra. Our first lemma is a simple consequence of the $||\cdot||$ -continuity of multiplication.

LEMMA 4.1. If R is any right ideal of A_p , then \overline{R} , the closure of R in A, is a closed right ideal of A.

LEMMA 4.2. If R is a right ideal of A_p and P is the orthogonal projection operator of A onto \overline{R} , the closure of R in A, then for any $a \in A_p$, $Pa \in A_p$. In particular, if R is relatively $|| \cdot ||$ -closed in A_p then $Pa \in R$.

Proof. This is immediate from Proposition 3.19, inasmuch as P is a left centralizer on A.

PROPOSITION 4.3. If R is a relatively $||\cdot||$ -closed right ideal of A_p , then $A_p = R \bigoplus R^{\perp}$, where R^{\perp} is the orthogonal complement of R in A_p .

Proof. Considering the closures in A of these right ideals, we have, for any $a \in A_p$, $a = a_1 + a_2$, where $a_1 \in \overline{R}$, $a_2 \in \overline{R^{\perp}}$. But by Lemma 4.2, $a_1 \in R$ and $a_2 \in R^{\perp}$.

REMARK 4.4. For a closed right ideal R in any Hilbert algebra, we have $\mathscr{L}(R) = R^{\perp *}$, where $\mathscr{L}(R)$ is the left annihilator of R. This is readily established by the argument used for an H^* -algebra [5, Theorem 12]. Combining this fact with Proposition 4.3 we obtain the following.

COROLLARY 4.5. $(A_p, || \cdot ||)$ is a dual Hilbert algebra.

Our next proposition, along with the known structure theory of H^* -algebras [1, Theorem 4.2], enables us to obtain a structure theorem for the Hilbert algebras A_p .

PROPOSITION 4.6. Let I be a closed two-sided ideal of A (and therefore an H^* -algebra). Then $I \cap A_p = I_p$, the p-class of I.

Proof. If $a \in I_p$ then [a], as an element of the H^* -algebra I, has a spectral decomposition $[a] = \Sigma \lambda_n e_n$, where $e_n \in I$ for each n, and $\Sigma \lambda_n^p ||e_n||^2 < \infty$. This is therefore the (unique) spectral decomposition of [a] in A, and therefore $a \in I \cap A_p$. Conversely, suppose $a \in I \cap A_p$. Since $a \in I$, [a] has a spectral decomposition $[a] = \Sigma \lambda_n e_n$ in I, and again this is its unique spectral decomposition in A. Since $a \in A_p$ we have $\Sigma \lambda_n^p ||e_n||^2 < \infty$, and therefore $a \in I_p$.

REMARK 4.7. Let J be a relatively $\|\cdot\|$ -closed two-sided ideal of A_p . Then J is a minimal closed ideal of A_p if and only if \overline{J} , the closure of J in A, is a minimal closed ideal of A. If the latter condition holds (so that \overline{J} is a topologically simple H^* -algebra), then J is a topologically simple Hilbert algebra.

We use these results and Lemma 4.2 to obtain our structure theorem for A_p as a Hilbert algebra.

THEOREM 4.8 The Hilbert algebra $(A_p, || \cdot ||)$ is the direct topological sum of its minimal closed two-sided ideals, which are mutually orthogonal. Each of these is a topologically simple Hilbert algebra and is the p-class of a minimal closed two-sided ideal of A.

For the remainder of this section we consider the Banach *-algebras $(A_p, |\cdot|_p)$. Our aim in the following development is twofold: (1) to investigate the $|\cdot|_p$ -closed right ideals of A_p ; (2) to obtain a structure theorem for $(A_p, |\cdot|_p)$ analogous to Theorem 4.8.

LEMMA 4.9. Let I be any $||\cdot||$ -closed two-sided ideal of A. For any $a \in A$, let a_1 denote the orthogonal projection of a on I. Then $(1) \quad (a^*)_1 = (a_1)^*,$ $(2) \quad [a]_1 = [a_1].$

Proof. Let $a = a_1 + a_2$, where $a_2 \in I^{\perp}$, the orthogonal complement of I in A. Then $a^* = a_1^* = (a_1)^* + (a_2)^*$. (1) follows readily from the fact that I and I^{\perp} are closed under the involution. To establish (2), we first note that $a^*a = a_1^*a_1 + a_2^*a_2$. Then, letting $[a] = [a]_1 + [a]_2$, we have $a^*a = [a]^2 = [a]_1^2 + [a]_2^2$, and hence $[a]_1^2 = a_1^*a^1$, by the uniqueness of the decomposition. If we show that $[a]_1$ is positive, then $[a]_1 =$ $[a_1]$ by the definition of $[a_1]$. For any $x \in A$, let $x = x_1 + x_2$, where $x_1 \in I, x_2 \in I^{\perp}$. Then $([a]_1x, x) = ([a]_1x_1 + [a]_1x_2, x_1 + x_2) = ([a]_1x_1, x_1) =$ $([a]_1x_1 + [a]_2x_1, x_1) = ([a]x_1, x_1) \ge 0$. **PROPOSITION 4.10.** Let $\{J_r: \gamma \in \Gamma\}$ be a family of mutually orthogonal relatively $||\cdot||$ -closed two-sided ideals of A_p . Let $a_{\gamma} \in J_{\gamma}$ for each γ , and let $a = \Sigma a_{\gamma}$ (in the $||\cdot||$ -topology). Then $|a|_p^p = \Sigma |a_{\gamma}|_p^p$, and hence $a \in A_p$ if and only if $\Sigma |a_{\gamma}|_p^p < \infty$.

Proof. Clearly, each a_r is the orthogonal projection of a on J_r , and hence, by the preceding lemma, $[a] = \Sigma[a_r]$. Now for each γ , let $[a_r] = \Sigma_n \lambda_{r_n} e_{\tau_n}$ be the spectral representation of $[a_r]$ in the H^* algebra \overline{J}_r , the $|| \cdot ||$ -closure of J_r in A. Then $|a_r|_p^p = \Sigma_n \lambda_{r_n}^p ||e_{\tau_n}||^2$. Also, $[a] = \Sigma_r \Sigma_n \lambda_{\tau_n} e_{\tau_n}$, and since in this sum there cannot be infinitely many equal coefficients, the spectral represention of [a] is obtained by merely grouping the terms of the series having the same coefficient, and then rearranging the terms, if necessary. Hence $|a|_p^p = \Sigma_r \Sigma_n \lambda_{\tau_n}^p ||e_{\tau_n}||^2 = \Sigma_r |a_r|_p^p$.

REMARK 4.11. This proposition also holds for $2 \leq p < \infty$. Also, it is easily seen that $|a|_{\infty} = \sup_{r} |a_{r}|_{\infty}$.

LEMMA 4.12. Let $a \in A_p$ and let ε be any positive number. Then there exist projections e and f in A_p such that $|a - ae|_p < \varepsilon$ and $|a - fa|_p < \varepsilon$.

Proof. Let A_0 be the intersection of all maximal commutative *-subalgebras of A containing [a]. Then, as in Lemma 2.1, we have a representation $[a] = \Sigma \alpha_n p_n$ (each $\alpha_n \neq 0$), which, by grouping and rearranging of terms, yields the spectral representation of [a]; hence $|a|_p^p = \Sigma \alpha_n^p ||p_n||^2$. (Note that $[a] \in (A_0)_p$.) We may write $[a] = ([a] - \Sigma_{n=1}^k \alpha_n p_n) + (\sum_{n=1}^k \alpha_n p_n)$, where $\sum_{n=1}^k \alpha_n p_n$ belongs to the relatively $||\cdot||$ -closed two-sided ideal $\sum_{n=1}^k (A_0)_p p_n$ of $(A_0)_p$, and $([a] - \sum_{n=1}^k \alpha_n p_n)$ belongs to the orthogonal complement of this ideal in $(A_0)_p$. By Proposition 4.10, $|a|_p^p = |[a]|_p^p = |[a] - \sum_{n=1}^k \alpha_n p_n|_p^p + |\sum_{n=1}^k \alpha_n p_n|_p^p$. But this last term is $\sum_{n=1}^k \alpha_n^p ||p_n||^2$, which has the limit $|a|_p^p$ as $k \to \infty$. We therefore have $\lim_{k\to\infty} |[a] - \sum_{n=1}^k \alpha_n p_n|_p^p = 0 = \lim_{k\to\infty} |[a] - [a] \sum_{n=1}^k p_n|_p^p$.

Hence for sufficiently large k there is a projection $e = \sum_{n=1}^{k} p_n$ such that $|[a] - [a]e|_p < \varepsilon$. Taking W to be the partial isometry associated with a, we have, using Proposition 3.19, $|a - ae|_p = |W[a] - (W[a])e|_p = |W[a] - W([a]e)|_p \le ||W|| |[a] - [a]e|_p < \varepsilon$. There is likewise a projection f such that $|a^* - a^*f|_p < \varepsilon$; hence $|a - fa|_p = |(a - fa)^*|_p < \varepsilon$.

COROLLARY 4.13. For any $a \in A_p$, $a \in \overline{aA_p} \cap \overline{A_pa}$, where the closure is in the $|\cdot|_p$ -topology.

We remarked at the beginning of this section that A_p is a special

case of an *IP*-algebra, and now that we have established the result of Corollary 4.13, we immediately have the following from [12, Theorems 3.5 and 4.9].

COROLLARY 4.14. $(A_p, |\cdot|_p)$ has dense socle, and is the direct topological sum of its minimal closed two-sided ideals.

COROLLARY 4.15. $(A_p, |\cdot|_p)$ is a dual algebra.

A simple consequence of Corollary 4.15 is the following.

PROPOSITION 4.16. Let R be a right ideal of A_p . R is closed in the $|\cdot|_p$ -topology if and only if R is relatively closed in the $||\cdot||$ -topology.

Proof. Since $||a|| \leq |a|_p$ for every $a \in A_p$, by Proposition 3.12, it is clear that every relatively $|| \cdot ||$ -closed subset of A_p is $| \cdot |_p$ -closed. Moreover, if the right ideal R is $| \cdot |_p$ -closed, then it is an annihilator ideal, by Corollary 4.15, and therefore is relatively $|| \cdot ||$ -closed, by the $|| \cdot ||$ -continuity of multiplication.

REMARK 4.17. This result holds for $2 \leq p \leq \infty$. In this case, R is clearly $||\cdot||$ -closed if it is $|\cdot|_p$ -closed. But if R is a $||\cdot||$ -closed right ideal of $A_p(=A)$, we have $R = R^{\perp \perp} = \mathscr{L}(R^{\perp})^*$, by 4.4. By the $|\cdot|_p$ -continuity of multiplication, $\mathscr{L}(R^{\perp})$ is $|\cdot|_p$ -closed.

We combine Proposition 4.16 with Proposition 4.3 to obtain the following.

COROLLARY 4.18. $(A_p, |\cdot|_p)$ is a right complemented algebra (in the sense of [11]).

More can be said about the manner in which A_p is the direct topological sum of its minimal closed two-sided ideals. In order to do so, we obtain a converse of Proposition 4.10, which leads to our final structure theorem.

PROPOSITION 4.19. Let $\{J_r: \gamma \in \Gamma\}$ be a family of mutually orthogonal closed two-sided ideals of A_p . Let $a_r \in J_r$ for each γ , and suppose that $\Sigma |a_r|_p^p < \infty$. Then there exists $a \in A_p$ such that $a = \Sigma a_r$, where the sum may be taken in the $|\cdot|_p$ -topology or the $||\cdot||$ -topology.

Proof. Considering only the nonzero a_{γ} , which we denote as a_n , let $s_k = \sum_{n=1}^k a_n$. Then, by Proposition 4.10, for k > m we have $|s_k - s_m|_p^p = |\sum_{n=m+1}^k a_n|_p^p = \sum_{n=m+1}^k |a_n|_p^p \to 0$ as $k, m \to \infty$. The Cauchy sequence $\{s_k\}$ thus has a limit a in the Banach algebra $(A_p, |\cdot|_p)$, and a =

790

 $\Sigma a_n = \Sigma a_{\tau}$ in the $|\cdot|_{p}$ -topology. (A standard argument shows that the limit is independent of the order of summation.) By Proposition 3.12, the sum is the same in the $||\cdot||$ -topology.

THEOREM 4.20. The Banach *-algebra $(A_p, |\cdot|_p)$ is the p-direct sum of its minimal closed two-sided ideals J_{λ} . The J_{λ} are mutually orthogal and each is a topologically simple Banach *-algebra. A_p is the "pdirect sum" in that it consists precisely of all sums $\Sigma \alpha_{\lambda}, \alpha_{\lambda} \in J_{\lambda}$, such that $\Sigma |\alpha_{\lambda}|_p^p < \infty$, where $a = \Sigma \alpha_{\lambda}$ may be understood as a limit in either the $|\cdot|_p$ -topology or the $||\cdot||$ -topology, and $|\alpha|_p = (\Sigma |\alpha_{\lambda}|_p^p)^{1/p}$.

5. Relationship to other systems. If A is a topologically simple H^* -algebra, then there is a *-isomorphism $x \to X$ of A onto the Schmidt class σc of operators on the Hilbert space $H = l_2(\Gamma)$, where Γ is the index set of a maximal family $\{q_{\tau}\}$ of mutually orthogonal primitive projections in A [1, Theorem 4.3]. Under this isomorphism, $||x|| = \alpha \sigma(X)$, where $\sigma(X)$ denotes the Schmidt norm of the operator X and $\alpha \geq 1$ is the norm of each of the projections q_{τ} (actually, all primitive projections in A have the same norm [7, Corollary 5.9]). Now if x is any nonzero element of A and $[x] = \Sigma \lambda_n e_n$ is the spectral representation of [x], then we may replace the nonprimitive projections a new representation

$$(*) \qquad [x] = \Sigma \mu_n p_n ,$$

where $\mu_m \leq \mu_k$ if m > k. For a given coefficient μ_n in (*), we shall call the number of primitive projections having μ_n as coefficient the multiplicity of μ_n in this representation, denoted by $m(\mu_n)$. We have, for $0 , <math>|x|_p = (\Sigma \mu_n^p) |p_n||^2)^{1/p} = \alpha^{2/p} (\Sigma \mu_n^p)^{1/p}$. Also, $|x|_{\infty} = \mu_1$. Since the μ_n are the nonzero elements of the spectrum of [x], and since the corresponding operator [X] is compact, these numbers are the nonzero characteristic values of [X]. Now for each μ_n , let $M(\mu_n)$ denote the multiplicity of μ_n as a characteristic value of the operator [X]; that is, the dimension of the subspace of H spanned by the characteristic vectors of [X] corresponding to μ_n . We shall show that $m(\mu_n) = M(\mu_n)$.

LEMMA 5.1. Let p be a primitive projection in the topologically simple H^* -algebra A. Then the corresponding projection P in σc is one-dimensional on H.

Proof. If P is not one-dimensional, let P = Q + R, where Q and R are projections onto orthogonal nonzero subspaces of P(H). Letting

q and r be the corresponding elements of A, we see that q and r are orthogonal projections in A with p = q + r. Thus p is not primitive.

LEMMA 5.2. For any μ_n in (*), $m(\mu_n) = M(\mu_n)$.

Proof. Let p_{n_1}, \dots, p_{n_k} be the projections in (*) having coefficient μ_n . Then $m(\mu_n) = k$. Also, letting P_{n_1}, \dots, P_{n_k} be the corresponding projections in σc , we have, using the preceding lemma, $\dim (P_{n_1} + \dots + P_{n_k})(H) = k$; therefore $M(\mu_n) \ge k$. Suppose $M(\mu_n) > k$, and let h be a nonzero element of H such that $[X]h = \mu_n h$ and h is orthogonal to $(P_{n_1} + \dots + P_{n_k})(H)$. Let Q be the orthogonal projection onto the one-dimensional subspace of H spanned by $\{h\}$. $Q \in \sigma c$, and $[X]Q = \mu_n Q$. Now let q be the corresponding projection in A; then $[x]q = \mu_n q$. For $i = 1, \dots, k, p_{n_i}q = 0$ since $P_{n_i}Q = 0$; and for $m \neq n_1, \dots, n_k$, $p_m[x]q = \mu_m p_m q = \mu_n p_m q$, so that $p_m q = 0$, since $\mu_m \neq \mu_n$. Thus q is orthogonal to all the p_n , which means that [x]q = 0, a contradiction. We conclude that $m(\mu_n) = k = M(\mu_n)$.

Now we observe that the coefficients μ_n in (*) are the nonzero characteristic values of [X] enumerated according to their multiplicity $M(\mu_n)$. Thus, for $0 , <math>|X|_p = (\Sigma \mu_n^p)^{1/p}$ and also $|X|_{\infty} = \mu_1$, where $|\cdot|_p$ here denotes the c_p norm of X as an operator on H. Finally, we have $|x|_p = \alpha^{2/p} |X|_p$ for $0 , and therefore the mapping <math>x \rightarrow X$ is a bicontinuous isomorphism of A_p into $c_p(H)$. Since $c_2 = \sigma c$ [2, p. 1093] and $c_p \subset c_2$ for $0 , the isomorphism is onto <math>c_p$ for these values of p.

Now let A be any proper H^* -algebra, and let $\{I_{\lambda}: \lambda \in A\}$ be the family of minimal closed two-sided ideals of A. Each I_{λ} is a topologically simple H^* -algebra and A is the Hilbert space direct sum ΣI_{λ} . For each $\lambda \in \Lambda$, let Γ_{λ} be the index set of a maximal family $\{e_{\lambda_{\gamma}}: \gamma \in \Gamma_{\lambda}\}$ of mutually orthogonal primitive projections in I_{λ} , and let α_{λ} be the norm $||e_{\lambda_{\gamma}}||$ of each of the $e_{\lambda_{\gamma}}$ in I_{λ} . For each $x_{\lambda} \in I_{\lambda}$ let X_{λ} be the corresponding Schmidt class operator on $H_{\lambda} = l_2(\Gamma_{\lambda})$. Then, as we have noted above, $|x_{\lambda}|_{p} = \alpha_{\lambda}^{2/p} |X_{\lambda}|_{p}, 0 , where <math>|X_{\lambda}|_{p}$ is the c_{p} norm of the operator X_{λ} . Then, by Proposition 4.10, we have $|x|_{p} =$ $(\Sigma \,|\, x_{\lambda} \,|_{p}^{p})^{1/p} = (\Sigma lpha_{\lambda}^{2} \,|\, X_{\lambda} \,|_{p}^{p})^{1/p}$ for $0 , and, by 4.11, <math>|\, x \,|_{\infty} = \sup_{\lambda} |\, x_{\lambda} \,|_{p} = \sum_{j=1}^{n} || x_{j} \,|_{j} = \sum_{j=$ $\sup_{\lambda} |X_{\lambda}|$. Thus, again, as in Proposition 4.10, $x \in A_p$ if and only if each $x_{\lambda} \in (I_{\lambda})_{p} = I_{\lambda} \cap A_{p}$ and $\Sigma |x_{\lambda}|_{p}^{p} < \infty$. These conditions in turn imply that each corresponding operator $X_{\lambda} \in c_p(H_{\lambda})$ and $\Sigma \alpha_{\lambda}^2 |X_{\lambda}|_p^p < \infty$. For $1 \leq 1$ $p \leq 2$, it has been established that the last-mentioned implication is an equivalence; for these values of p, therefore, in the special situation in which each H_{λ} is finite-dimensional, we have shown that the algebras A_p are instances of the \mathcal{C}_p spaces studied in [3, pp. 70 ff.] and [5].

References

1. W. Ambrose, Structure theorems for a special class of Banach algebras, Trans. Amer. Math. Soc., 57 (1945), 364-386.

2. N. Dunford and J. T. Schwartz, *Linear Operators*, vol. II, New York, Interscience, 1963.

3. E. Hewitt and K. A. Ross, Abstract Harmonic Analysis, vol. II, Heidelberg, Springer-Verlag, 1970.

4. I. Kaplansky, Dual rings, Ann. of Math., 49 (1948), 689-701.

5. J. La Duke, On a certain generalization of l_p spaces, Pacific J. Math., **35** (1970), 155-168.

6. C. A. McCarthy, c_p, Israel J. Math., 5 (1967), 249-271.

7. M. A. Rieffel, Square-integrable representations of Hilbert algebras, J. Functional Analysis, **3** (1969). 265-300.

8. P. P. Saworotnow and J. C. Friedell, *Trace-class for an arbitrary H*-algebra*, Proc. Amer. Math. Soc., **26** (1970), 95-100.

9. P. P. Saworotnow, Trace-class and centralizers of an H*-algebra, Proc. Amer. Math. Soc., 26 (1970), 101-104.

10. R. Schatten, Norm Ideals of Completely Continuous Operators, Berlin, Springer-Verlag, 1960.

11. B. J. Tomiuk, Structure theory of complemented Banach algebras, Canad. J., Math. 14 (1962), 651-659.

12. B. Yood, Noncommutative Banach algebras and almost periodic functions, Illinois J. Math. 7 (1963), 305-321.

Received July 23, 1971 and in revised form November 23, 1971. This research was done while the author was at the University of Oregon on sabbatical leave from Le Moyne College. He gratefully acknowledges the direction and assistance of Professor Bertram Yood.

LE MOYNE COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

C. R. HOBBY

Stanford University Stanford, California 94305 J. DUGUNDJI

Department of Mathematics University of Southern California Los Angeles, California 90007

RICHARD ARENS University of California Los Angeles, California 90024

University of Washington Seattle, Washington 98105

ASSOCIATE EDITORS

E.F. BECKENBACH B.H. NEUMANN F. WOLF K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON

AMERICAN MATHEMATICAL SOCIETY NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The Pacific Journal of Mathematics is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics Vol. 42, No. 3 March, 1972

Catherine Bandle, <i>Extensions of an inequality by Pólya and Schiffer for vibrating membranes</i>	543
S. J. Bernau, Topologies on structure spaces of lattice groups	557
Woodrow Wilson Bledsoe and Charles Edward Wilks, On Borel product	
measures	569
Eggert Briem and Murali Rao, Normpreserving extensions in subspaces of $C(X)$	581
Alan Seymour Cover, Generalized continuation	589
Larry Jean Cummings, Transformations of symmetric tensors	603
Peter Michael Curran, Cohomology of finitely presented groups	615
James B. Derr and N. P. Mukherjee, <i>Generalized quasicenter and</i>	
hyperquasicenter of a finite group	621
Erik Maurice Ellentuck, Universal cosimple isols	629
Benny Dan Evans, Boundary respecting maps of 3-mainfolds	639
David F. Fraser, A probabilistic method for the rate of convergence to the	
Dirichlet problem	657
Raymond Taylor Hoobler, Cohomology in the finite topology and Brauer	
groups	667
Louis Roberts Hunt, Locally holomorphic sets and the Levi form	681
B. T. Y. Kwee, On absolute de la Vallée Poussin summability	689
Gérard Lallement, On nilpotency and residual finiteness in semigroups	693
George Edward Lang, Evaluation subgroups of factor spaces	701
Andy R. Magid, A separably closed ring with nonzero torsion pic	711
Billy E. Rhoades, <i>Commutants of some Hausdorff matrices</i>	715
Maxwell Alexander Rosenlicht, <i>Canonical forms for local derivations</i>	721
Cedric Felix Schubert, On a conjecture of L. B. Page	733
Reinhard Schultz, Composition constructions on diffeomorphisms of	
$S^p \times S^q$	739
J. P. Singhal and H. M. (Hari Mohan) Srivastava, A class of bilateral	
generating functions for certain classical polynomials	755
Richard Alan Slocum, Using brick partitionings to establish conditions	
which insure that a Peano continuum is a 2-cell, a 2-sphere or an annulus	763
James F. Smith, The p-classes of an H*-algebra	777
Jack Williamson, Meromorphic functions with negative zeros and positive	
poles and a theorem of Teichmuller	795
William Robin Zame, Algebras of analytic functions in the plane	811