

Pacific Journal of Mathematics

QUASI PROJECTIVES IN ABELIAN AND MODULE CATEGORIES

KULUMANI M. RANGASWAMY AND N. VANAJA

QUASI PROJECTIVES IN ABELIAN AND MODULE CATEGORIES

K. M. RANGASWAMY AND N. VANAJA

If R is a ring without zero divisors then it is shown that any torsion-free quasi-projective left R -module A is projective provided A is finitely generated or A is "big". It is proved that the universal existence of quasi-projective covers in an abelian category with enough projectives always implies that of the projective covers. Quasi-projective modules over Dedekind domains are described and as a biproduct we obtain an infinite family of quasi-projective modules Q such that no direct sum of infinite number of carbon copies of Q is quasi projective. Perfect rings are characterised by means of quasi-projectives. Finally the notion of weak quasi-projectives is introduced and weak quasi-projective modules over a Dedekind domain are investigated.

1. Introduction. An object A in a category \mathcal{A} is called quasi-projective [14] if given an epimorphism $A \xrightarrow{f} B$ and a morphism $g: A \rightarrow B$, there is $h: A \rightarrow A$ making the following diagram

$$\begin{array}{ccc} & A & \\ & \swarrow h & \downarrow g \\ A & \xrightarrow{f} & B \end{array}$$

commutative. This paper starts with the investigation of the quasi-projectives in an abelian category. Utilising a few basic lemmas, it is shown that the universal existence of the quasi-projective covers in an abelian category \mathcal{A} implies that of the projective covers, provided \mathcal{A} possesses enough projectives and this answers affirmatively a question of Faith [4] in a general form. Next we consider quasi-projectives in the category of modules. It turns out that "big" torsion-free quasi-projectives over rings without zero divisors are always projective. Artin semi-simple rings are characterised as those rings over which quasi-projectives and projectives coincide. In § 5, quasi-projectives over a Dedekind domain R are investigated: A quasi-projective R -module is either torsion or torsion-free. A torsion R -module is quasi-projective if and only if it is quasi-injective but not

injective. If R is a complete discrete valuation ring, then the torsion-free quasi-projective R -modules are just the free R -modules and the torsion-free modules of finite rank. Suppose R is a Dedekind domain which is not a complete discrete valuation ring and σ is the number of distinct prime ideals of R . If $\sigma \leq 2^{\aleph_0}$, then all the torsion-free R -quasi-projectives are projective. If $\sigma > 2^{\aleph_0}$, then a torsion-free quasi-projective R -module A is projective if either (i) $\text{rank } A \leq \aleph_0$ or (ii) $\text{rank } A > \sigma$. In the case when $\aleph_0 < \text{rank } A < \sigma$, A is torsionless, \aleph_1 -projective and contains a free summand F having the same rank as A . As a biproduct we at once get an infinite family of quasi-projective modules A such that no direct sum of infinite number of copies of A is quasi-projective. In § 6, Perfect rings are characterised as those rings R such that R -quasi-projectivity survives under direct limits. A weakened form of quasi-projectivity — called weak quasi-projectivity — is considered in the last section and weak quasi-projectives over a Dedekind domain are completely characterised.

2. Preliminaries. All the rings that we consider are associative and are assumed to possess an identity and all the modules unitary left modules. A sub-module S of an R -module M is called *fully invariant* if S is stable under every R -endomorphism of M . S is called a *small* submodule, if $S + T = M$ implies $T = M$ for any submodule T of M . A projective module P is called a *projective cover* of M if there is an epimorphism $P \rightarrow M$ whose kernel is small. A module M over an integral domain is called *reduced* if 0 is the only divisible submodule of M . By the *rank* of a torsion-free module M over a Dedekind domain R we shall mean the cardinality of a maximal R -independent subset of M . An R -module M is called *quasi-injective* if for any exact sequence $0 \rightarrow S \xrightarrow{i} M$, the induced sequence

$$\text{Hom}_R(M, M) \xrightarrow{i^*} \text{Hom}_R(S, M) \longrightarrow 0$$

is exact, where $i^*(f) = i \circ f$ for all f in $\text{Hom}_R(M, M)$. For the basic results in category theory, modules and abelian groups, the reader is referred to [5], [6], [10] and [11].

3. Quasi-projectivity in abelian categories. In this section, we examine the properties of quasi-projective objects in an abelian category. The main result shows that the universal existence of quasi-projective covers in an abelian category \mathcal{A} implies that of projective covers, provided \mathcal{A} possesses enough projectives.

NOTE. In conformity with our notation in the subsequent sections, a composite $f \circ g$ of two morphisms is obtained by applying f

first and then g .

LEMMA 3.1 [14]. *In an abelian category, any retract of a quasi-projective is quasi-projective.*

The following lemma gives a condition under which an object becomes projective.

LEMMA 3.2. *An object A in an abelian category is projective if and only if there exists an epimorphism $P \twoheadrightarrow A$ with P projective and $A \oplus P$ quasi-projective.*

Proof. We prove only the “if” part. Let $f: P \twoheadrightarrow A$ be the given epimorphism, $A \xrightarrow{i} A \oplus P \xrightarrow{j} A = 1_A$ and $P \xrightarrow{j'} A \oplus P \xrightarrow{j'} P = 1_P$. By the quasi-projectivity of $A \oplus P$, there exists $g: A \oplus P \rightarrow A \oplus P$ such that $A \oplus P \xrightarrow{j} A = A \oplus P \xrightarrow{g} A \oplus P \xrightarrow{j'} P \xrightarrow{f} A$. Then

$$1_A = ij = i(g \circ j' \circ f) = (i \circ g \circ j')f.$$

Thus A is a retract of P and hence projective.

Dualizing 3.2, we obtain

LEMMA 3.2'. *An object A in an abelian category is injective if and only if there is a monomorphism $A \rightarrow I$ with I injective and $A \oplus I$ quasi-injective.*

Next we examine the universal existence of quasi-projective covers.

DEFINITION 3.3. (i) An epimorphism f in a category is called a *minimal epimorphism* if, whenever $g \circ f$ is an epimorphism, g itself is an epimorphism.

(ii) $A \rightarrow X$ is called a *projective (quasi-projective) cover* in a category, if A is projective (quasi-projective) and f is a minimal epimorphism.

(iii) A category \mathcal{A} is called *perfect (quasi-perfect)* if every object in \mathcal{A} possesses a projective (quasi-projective) cover.

(iv) A category is said to possess *enough projectives*, if, to every object A , there is an epimorphism $P \rightarrow A$ with P projective.

REMARK. (i) For an axiomatic treatment of minimal epimorphisms see [1]. Observe that in the category of R -modules, an epimorphism $f: A \rightarrow B$ is minimal if and only if $\text{Ker } f$ is small in A .

(ii) The notion of a perfect category has been considered in [2], [3].

(iii) Our definition of a quasi-projective cover is slightly different from the one defined in [14] for modules. However, it is easy to see that for the category of modules over a ring R , the universal existence of quasi-projective covers according to the new definition is equivalent to the universal existence of quasi-projective covers according to the definition given in [14].

It is clear that a perfect abelian category is quasi-perfect. Conversely, is a quasi-perfect abelian category perfect? This is the category-theoretical formulation of a question raised by C. Faith [4]¹⁾. The following theorem answers this:

THEOREM 3.4. *An abelian category \mathcal{A} is perfect if and only if it is quasi-perfect and possesses enough projectives.*

Proof. IF part: Let $A \in \mathcal{A}$ and $P \xrightarrow{u} A$ an epimorphism with P projective. Let $g: Q' \rightarrow A \oplus P$ be a quasi-projective cover of $A \oplus P$. Consider the following commutative diagram

$$\begin{array}{ccccccccc}
 & & Q & \xrightarrow{i''} & Q' & & & & \\
 & & \downarrow g' & & \downarrow g & & & & \\
 0 & \longrightarrow & A & \xrightarrow{i} & A \oplus P & \xrightarrow{j'} & P & \longrightarrow & 0
 \end{array}$$

where the square is a pull-back and

$$A \xrightarrow{i} A \oplus P \xrightarrow{j} A = 1_A, P \xrightarrow{i'} A \oplus P \xrightarrow{j'} P = 1_P.$$

By Lemma 2.61 of [5],

$$0 \longrightarrow Q \xrightarrow{i''} Q' \xrightarrow{g \cdot j'} P \longrightarrow 0$$

is an exact sequence which splits since P is projective. Let $f: P \rightarrow Q'$ be such that $f \circ g \circ j' = 1_P$. Since g is epic and the square is a pull-back, g' is also epic. We claim g' is minimal. Let $h': C \rightarrow Q$ be such that $h' \circ g'$ is epic. Let $h = (h' \circ i'') \oplus f$. Consider the following commutative diagram

¹⁾ While this paper was being written we found out that this question has been recently answered independently by A. Koehler [12], K. R. Fuller, D. A. Hill and J. Golan for the category of R -modules.

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & C & \longrightarrow & C \oplus P & \longrightarrow & P & \longrightarrow & 0 \\
 & & \downarrow h' & & \downarrow h & & \downarrow 1_P & & \\
 0 & \longrightarrow & Q & \longrightarrow & Q' & \longrightarrow & P & \longrightarrow & 0 \\
 & & \downarrow g' & & \downarrow g & & \downarrow 1_P & & \\
 0 & \longrightarrow & A & \xrightarrow{i} & A \oplus P & \xrightarrow{j} & P & \longrightarrow & 0
 \end{array}$$

where the top row is split exact with the obvious maps. By the 5-lemma, $h \circ g$ is epic and since g is minimal, h is epic. Since

$$C \longrightarrow C \oplus P \longrightarrow P \longrightarrow 0$$

is exact, again by Lemma 2.61 of [5], the left top square is a pull-back. Since h is epic, h' is also epic. Thus g' is minimal epic. Since P is projective and $u: P \rightarrow A$, there exists $v: P \rightarrow Q$ such that $v \circ g' = u$. By the minimality of g' , v is an epimorphism. Then the quasi-projectivity of $Q \oplus P$ and the Lemma 3.2 imply that Q is projective. Thus $g': Q \rightarrow A$ is a projective cover of A and we conclude that the category is perfect.

REMARK 1. Theorem 3.4 is best possible in the sense that it fails to be true if \mathcal{A} is not an abelian category. To see this, let \mathcal{A}_b be the category of all the abelian groups and \mathcal{A} the full subcategory of \mathcal{A}_b consisting of all the cyclic groups. Then \mathcal{A} is not abelian. \mathcal{A} has enough projectives and is clearly quasi-perfect (every object in \mathcal{A} is quasi-projective). But \mathcal{A} is not perfect since the prime cyclic group $Z(p)$ possesses no projective cover in \mathcal{A} .

REMARK 2. A quasi-perfect abelian category need not possess enough projectives. The category \mathcal{F}_p of all finite abelian p -groups is one such. The quasi-projectives in \mathcal{F}_p are the direct sums of isomorphic cyclic p -groups [7]. \mathcal{F}_p is abelian and is readily seen to be quasi-perfect. But it possesses no non-trivial projectives.

4. **Quasi-projectives in the category of modules.** In this section we indicate some of the simple properties of quasi-projective modules over a ring. We also investigate when a quasi-projective module over a ring R without zero-divisors becomes projective. It turns out in a surprisingly simple way that the “big” torsion-free quasi-projectives over such R are projective. Some of the preliminary lemmas in this section hold in any abelian category but, for the sake

of convenience, we will consider only the module case. Lemmas 4.3 and 4.4 occur in [7], but are proved here for the same of completeness.

LEMMA 4.1. [14]. *If A is a quasi-projective R -module and S is fully invariant in A , then A/S is quasi-projective.*

COROLLARY. *Let I be a two sided ideal of a ring R . Then R/I is quasi-projective as an R -module.*

The converse of Lemma 4.1 is not always true. It holds, however, under some restriction on S , as indicated below.

LEMMA 4.2. *Let S be a small submodule of a quasi-projective module A . Then A/S is quasi-projective if and only if S is fully invariant in A .*

To prove this, replace the word, "projective" in the proof of proposition 2.2 of [14] by "quasi-projective".

The following lemma gives a condition when a submodule of a quasi-projective module becomes a summand.

LEMMA 4.3. *Let S be a submodule of a quasi-projective module A . Then S is a summand if and only if A/S is isomorphic to a summand of A .*

Proof. Let $A = B \oplus C$ and $f: B \rightarrow A/S$ be an isomorphism. Define $g: A \rightarrow A/S$ by $g|B = f$ and $g|C = 0^{2)}$. By the quasi-projectivity of A , g lifts to an endomorphism h of A such that $h \circ p = g$, where $p: A \rightarrow A/S$ is the natural map. Set $p' = f^{-1} \circ h$. Since $p' \circ p = 1_{A/S}$, the sequence $0 \rightarrow S \rightarrow A \rightarrow A/S \rightarrow 0$ splits and thus S is a summand of A .

Dualising 4.3, we obtain a corresponding statement for quasi-injectives.

LEMMA 4.3'. *Let S be a submodule of a quasi-injective module A . Then S will be a summand if and only if S is isomorphic to a summand of A .*

REMARK. Lemma 3.2 and 3.2' can also be easily deduced from 4.3 and 4.3' respectively.

LEMMA 4.4. *Let A be a quasi-projective module. Then the exact*

²⁾ $g|B$ denotes the restriction of the map g to B .

sequence $0 \rightarrow T \rightarrow S \xrightarrow{f} A \rightarrow 0$ splits, whenever S is a submodule of A .

Proof. Let $g: A \rightarrow A/T$ be an epimorphism such that $g|_S = f$. Let $h: A \rightarrow A/T$ be monic with $\text{Im } h = S/T$. Then there exists an endomorphism h' of A satisfying $h' \circ g = h$. Since $\text{Im } h' = S$, it is readily seen that h' is a split map of the sequence $0 \rightarrow T \rightarrow S \xrightarrow{f} A \rightarrow 0$. Hence the Lemma.

Dualising 3.4, we obtain an analogous property of quasi-injectives.

LEMMA 4.4'. *If A is quasi-injective, then the exact sequence $0 \rightarrow A \xrightarrow{i} X \rightarrow Y \rightarrow 0$ splits whenever X is a quotient of A .*

As an easy application of Lemma 4.4 we show that big torsion-free quasi-projectives over an integral domain are projective.

THEOREM 4.5. *Let R be a ring without zero divisors. Then any torsion-free quasi-projective R -module containing an R -independent subset of cardinality exceeding the cardinality of R is projective.*

We may assume, without loss in generality, that R is infinite (since otherwise R becomes a field). Let A be a quasi-projective torsion-free R -module and S a maximal R -independent subset with $|S| \geq |R|$. Let F be the (free) submodule generated by S . Then $|A| = |S| \cdot |R| = |S|$ and so A can be obtained as an epimorphic image of F . Since F is free, A is projective by Lemma 4.4.

REMARK. (i) From the proof of 4.5 it is clear that, if R has no zero divisors, then a torsion-free quasi-projective R -module A is projective exactly when $\bigoplus_m A$ is quasi-projective for every cardinal m .

(ii) K. H. Fuller and D. A. Hill (Notices, Amer. Math. Soc., 16 (1969) 961) show that if A is finitely generated quasi-projective, then $\bigoplus_m A$ is quasi-projective for any m . An immediate deduction from (i) above: *If R has no zero divisors, then a finitely generated torsion-free quasi-projective R -module is projective.*

COROLLARY 4.6. *A quasi-projective module over a ring without zero divisors is projective if and only if it is torsion-free and possesses a projective cover.*

We need only to prove the “if” part. Let A be torsion-free quasi-projective and $A \cong P/S$, P projective and S small. By Lemma 4.2, S is fully invariant in P . If m denotes the cardinality of R ,

then $\bigoplus_m A \cong (\bigoplus_m P)/(\bigoplus_m S)$ is quasi-projective, since $\bigoplus_m S$ is fully invariant in $\bigoplus_m P$. The projectivity of A then follows from Theorem 4.5.

REMARK. One can deduce that over a ring without zero divisors a quasi-projective module with a projective cover is either torsion or torsion-free. For, suppose A , P and S are as in the preceding proof and A contains a torsion-free element $a \neq 0$. If $m > \aleph_0 \cdot |R| \cdot |A|$, then $\bigoplus_m A$ is quasi-projective, has cardinality m and contains a free submodule F of rank m . By Lemma 4.4, $\bigoplus_m A$ and hence A is projective (and torsion-free).

The following theorem characterises Artin Semisimple rings by means of quasi-projectives.

THEOREM 4.7. *The following properties are equivalent for any ring R :*

- (i) R is Artin Semi-simple.
- (ii) The R -modules with a projective cover are precisely the quasi-projectives.
- (iii) Every quasi-projective R -module is projective.

Proof. Trivially (i) implies (ii).

Assume (ii). Let Q be quasi-projective. By assumption Q possesses a projective cover P . Then $P \oplus Q$ will have a projective cover and hence is quasi-projective by hypothesis. Lemma 3.2 then implies that Q is projective.

Assume (iii). Since any simple R -module is quasi-projective, it becomes projective by assumption. Then all the maximal left ideals of R are direct summands of the left R -module R and since R has 1, we conclude that R is Artinian Semi-simple. This completes the proof.

REMARK 1. Observe that if every R -module is quasi-projective then, by Lemma 3.2, R satisfies the condition (iii) above and hence R is Artinian Semi-simple.

REMARK 2. Johnson and Wong [9] showed that the quasi-injective modules over any ring R are exactly the fully invariant submodules of injective R -modules. A natural question is whether this can be dualised to quasi-projectives. Precisely, *must every quasi-projective R -module A be of the form P/S with P projective and S fully invariant in P ?* Jans and Wu [14] answered this in the affirmative under the assumption that A has a projective cover. In the general case,

the answer turns out to be in the negative. To see this, consider $M = \bigoplus (Z/pZ)$, where Z is the ring of integers, \bigoplus is a Z -module direct sum and p runs over the set of all primes in Z . Clearly M is a quasi-projective Z -module [7]. But M cannot be written as P/S , where P is a projective (hence free) abelian group and S fully invariant in P , since the only fully invariant subgroups of a free abelian group F are of the form nF , $n = 1, 2, \dots$.

REMARK 3. In the statement of the Theorem 4.7(ii), if we replace “precisely” by “necessarily”, we obtain a characterisation of Jacobson semi-simple rings: *A ring R is Jacobson semi-simple if and only if the R -modules possessing projective covers are necessarily quasi-projective.* To see this, assume the “if” part. Then, by Lemma 4.2, the small submodules of any projective R -module P are fully invariant in P . In particular, let $P = R_1 \oplus R_2$ with $R_i = R$ and let $J_i = J$, the Jacobson radical of R , for $i = 1, 2$. Now J_1 is small in R_1 and hence in P . But then J_1 would be fully invariant in P , an impossibility since J_1 can be mapped onto J_2 by an endomorphism of P . Thus $J_1 = 0$ and R is Jacobson Semi-simple. The converse follows on noting that if R is Jacobson Semi-simple, then 0 is the only small submodule of any projective R -module.

5. **Quasi-projectives over Dedekind domains.** In this section we propose to describe the quasi-projective modules over an arbitrary Dedekind domain R . First, observe that if A is any quasi-projective R -module, then any exact sequence $0 \rightarrow S \xrightarrow{i} A \xrightarrow{j} A/S \rightarrow 0$ yields the following two exact sequences.

$$\begin{aligned} 0 &\longrightarrow \text{Hom}_R(A, S) \xrightarrow{i'} \text{Hom}_R(A, A) \xrightarrow{j'} \text{Hom}_R(A, A/S) \longrightarrow 0 \\ 0 &\longrightarrow \text{Ext}_R^1(A, S) \xrightarrow{i''} \text{Ext}_R^1(A, A) \xrightarrow{j''} \text{Ext}_R(A, A/S) \longrightarrow 0. \end{aligned}$$

We first consider the torsion free quasi-projective modules. To avoid the trivial situations, the integral domains that we consider are not fields, unless explicitly stated.

LEMMA 5.1. *Let R be a Dedekind domain. Then the quotient field K of R is a quasi-projective R -module if and only if R is a complete discrete valuation ring.*

Proof. Suppose K is quasi-projective. Given any $f \in \text{Hom}_R(K/R, K/R)$, there exists a $f' \in \text{Hom}_R(K, K)$ such that $f' \circ j = j \circ f$ where j is the natural map from K onto K/R . Let $f'' = f'|_R$. Since $Rf' \subseteq R$, f'' is given by a multiplication by an element of R . It is readily

seen that the association $f \mapsto f''$ gives an isomorphism of $\text{Hom}_R(K/R, K/R)$ onto R . Now the exact sequence $0 \rightarrow R \rightarrow K \rightarrow K/R \rightarrow 0$ yields an exact sequence

$$\begin{aligned} \text{Hom}_R(K/R, K) = 0 &\longrightarrow \text{Hom}_R(K/R, K/R) \longrightarrow \text{Ext}_R^1(K/R, R) \\ &\longrightarrow \text{Ext}_R^1(K/R, K) = 0 \end{aligned}$$

(the first term is zero since K/R is torsion and K is torsion-free). Thus $R \cong \text{Hom}_R(K/R, K/R) \cong \text{Ext}_R^1(K/R, R)$ and the Corollary 7.9 of [13] implies that R is a complete discrete valuation ring.

Conversely, suppose R is a complete discrete valuation ring. Then any R -submodule S of K is isomorphic to R or K and hence, by Theorem 7.9 of [13], $\text{Ext}_R^1(K, S) = 0$. K is then clearly quasi-projective.

We shall first describe the torsion-free quasi-projectives over Dedekind domains which are not complete discrete valuation rings.

LEMMA 5.2. *Suppose R is a Dedekind domain which is not a complete discrete valuation ring. Then any torsion-free quasi-projective R -module A is torsionless.*

Proof. Let $0 \neq x \in A$ and S the pure submodule generated by x . Since R is not a complete discrete valuation ring, A (and therefore S) is reduced, by Lemma 3.1. Thus $S \neq PS$ for some prime ideal P of R . Then S/PS , being bounded and pure, is a summand of A/PS (Theorem 5 [11]). A nonzero cyclic summand of S/PS will be isomorphic to R/P and can be written as Ry/Py , for some $y \in S$. Let $g: S/PS \rightarrow Ry/Py$ be a nonzero map. Consider the following diagram

$$\begin{array}{ccc} & A & \\ & \swarrow h & \downarrow f' \circ g \\ A & \xrightarrow{f} & A/Py \longrightarrow 0 \end{array}$$

where $f': A \rightarrow S/PS$ is obtained via the projection $A/PS \rightarrow S/PS$ and f is the natural map. By the quasi-projectivity of A , there exists $h: A \rightarrow A$ making the diagram commutative. Now $A(h \circ f) = A(f' \circ g) \cong Ry/Py$, so that $Ah \cong Ry$. Thus $h: A \rightarrow Ry \cong R$ and $xh \neq 0$ since h does not vanish on the rank 1 submodule S . It follows that A is torsionless.

COROLLARY 5.3. *Let R be a Dedekind domain which is not a complete discrete valuation ring. Then any torsion-free R -module A is \aleph_1 -projective. Hence any torsion-free R -module of at most countable rank is projective.*

Proof. Let S be a submodule of A of rank 1. By Lemma 5.2, A is torsionless so that for each $a \neq 0$ in S , there exists $f: A \rightarrow R$ such that $af \neq 0$. Since S has rank 1 and $\text{im } f$ is torsion-free, $f|_S$ is mono. As R is hereditary, S is projective. By finite induction, it is clear that any submodule of A of finite rank is projective. Then a well-known step-wise argument (see for example Lemma 8.3.1 [13]) yields that any submodule of countable rank of A is projective.

In the following σ denotes cardinality of the set of all distinct prime ideals of R .

PROPOSITION 5.4. *Let R be a Dedekind domain. Then any torsion-free quasi-projective of rank $m \geq \sigma \aleph_0$ is projective.*

Proof. Let A be a torsion-free R -module of rank $m \geq \sigma \aleph_0$ and K be the quotient field of R . It is easy to see that $R(P^\infty)$ is countably generated. Now K/R is $\bigoplus_P R(P^\infty)$, where P runs over the set of distinct non-zero prime ideals of R and hence K has a generating set of cardinality $\sigma \aleph_0$. If D is an injective hull of A , then $D \cong \bigoplus_m K$ has a generating set of cardinality m . It is then readily seen that A itself is generated by m elements. Let F be a free submodule of A of rank m (for example F may be the submodule generated by a maximal R -independent subset of A). A can be got as an epimorphic image of F and hence by Lemma 4.4, A is a direct summand of F and hence projective.

Combining 5.3 and 5.4, we get the following.

THEOREM 5.5. *Let R be a Dedekind domain which is not a complete discrete valuation ring and $\sigma \leq \aleph_0$. Then a torsion-free R -module is quasi-projective if and only if it is projective.*

REMARK. If we assume the continuum hypothesis and use 5.3 and 5.4, then we can sharpen 5.5 to the following: *Let R be a Dedekind domain which is not a complete discrete valuation ring and $\sigma \leq 2^{\aleph_0}$. Then any torsion-free quasi-projective R -module is projective.*

Next we consider the case when $\sigma > 2^{\aleph_0}$.

PROPOSITION 5.6. *Let R be a Dedekind domain and A be a*

torsion-free quasi-projective R -module of infinite rank m . Then A contains a free summand of rank m .

Proof. Let P be any non-zero prime ideal of R . $R(P^\infty)$ is a countably generated injective R -module. If $Q = \bigoplus_m R(P^\infty)$, then, as R is Noetherian, Q is an injective R -module. Clearly Q has a generating set of cardinality m . Let F be the free-submodule generated by a maximal R -independent subset of A . Then Q can be obtained as a quotient of F , $Q \cong F/S$ for some submodule S . Consider the following diagram,

$$\begin{array}{ccc}
 & A & \\
 & \swarrow h & \downarrow f \\
 & & A/S \\
 A & \xrightarrow{f} & A/S = F/S \oplus T/S
 \end{array}$$

where $g: A/S \rightarrow F/S$ is a projection of A/S onto the injective summand F/S and f is the natural map. By the quasi-projectivity of A , there exists $h: A \rightarrow A$ such that $h \circ f = f \circ g$. It is clear that $Ah \subseteq F$ and since R is hereditary Ah is projective. As F/S is a direct sum of m copies of $R(P^\infty)$, it is clear that the rank of $Ah = m$. Thus $A = F' \oplus K$, where K is the kernel of h and F' is a projective module of infinite rank m and hence is free [11].

Combining 5.3, 5.4 and 5.6 we get,

THEOREM 5.7. *Let R be a Dedekind domain with $\sigma > 2^{\aleph_0}$. Then any torsion-free quasi-projective R -module A is projective if either (i) $\text{rank } A \leq \aleph_0$ or (ii) $\text{rank } A \geq \sigma$. In the case when $\aleph_0 < \text{rank } A < \sigma$, A is torsionless, \aleph_1 -projective and contains a free summand F having the same rank as A .*

The following theorem characterises torsion-free quasi-projectives over a complete discrete valuation ring.

THEOREM 5.8. *Suppose R is a complete discrete valuation ring. Then the torsion-free quasi-projective R -modules are just the free R -modules and the torsion-free R -modules of finite rank.*

Proof. By Kaplansky [10], any torsion-free R -module of finite rank is of the form $(\bigoplus_{i=1}^n K_i) \oplus (\bigoplus_{j=1}^m R_j)$ where each $R_j \cong R$ and each $K_i \cong K$, the quotient field of R . Thus if A is any finite rank torsion-free R -module and S is any submodule, then both are direct

sums of finite number of copies of K and R , so that

$$\text{Ext}_R^1(A, S) \cong \bigoplus_r \text{Ext}_R^1(K, R),$$

where r is finite. By Lemma 5.1, K is quasi-projective so that

$$\text{Ext}_R^1(K, R) = 0.$$

Thus $\text{Ext}_R^1(A, S) = 0$, whence $\text{Hom}_R(A, A) \xrightarrow{f'} \text{Hom}_R(A, A/S) \rightarrow 0$ is exact for every submodule S of A , where f' is induced by the natural map $f: A \rightarrow A/S$. The quasi-projectivity of A then follows. On the other hand if A is a torsion-free quasi-projective R -module of infinite rank, then by Proposition 5.4, A is projective and hence free.

COROLLARY 5.9. *If A is quasi-projective, then a direct sum $\bigoplus A$ of copies of A need not be quasi-projective.*

EXAMPLE. Suppose A is any torsion-free module of finite rank over a complete discrete valuation ring R such that A is not projective (for example $A = K$, the quotient field of R). Then any finite direct sum of copies A is quasi-projective but, by 5.8, no direct sum of infinite number of copies of A can be quasi-projective.

We shall now describe the torsion quasi-projectives over R .

THEOREM 5.10. *A torsion module A over a Dedekind domain R is quasi-projective if and only if each P -primary component A_P is a direct sum copies of the same cyclic module R/P^k for some fixed positive integer k depending on P .*

Proof. Since a P -primary module over R can be viewed as a module over the principal ideal domain R_P , and quasi-projectivity survives under this transition, we may assume that R itself is a principal ideal domain. Our proof would be sketchy since it is similar to the one given in [7]. Now $R(P^\infty)$ is not quasi-projective since otherwise, by Lemma 4.3, every submodule of $R(P^\infty)$ would be a summand. Thus a torsion quasi-projective R -module A is necessarily reduced. Again, by Lemma 4.3, A cannot contain a summand of the form $(R/P^{k_1}) \oplus (R/P^{k_2})$ with $k_1 > k_2$, since there is an epimorphism $R/(P^{k_2}) \rightarrow R/(P^{k_1})$ whose kernel is not a summand. Thus the basic submodules B_P (see [6]) of each P -primary component A_P are bounded and since the A_P are reduced, each A_P coincides with B_P which is clearly a direct sum of isomorphic cyclic modules. The "only if" part follows.

Conversely, if A is a direct sum $\bigoplus_m R/(P^k)$ of isomorphic cyclic modules, then $A \cong F/P^k F$, where F is free, say, $F = \bigoplus_m R$. Since

$P^k F$ is fully invariant in F , A is quasi-projective, by 4.1.

COROLLARY 5.11. *A torsion module A over a Dedekind domain R is quasi-projective if and only if A is quasi-injective but not injective.*

Proof. By Johnson and Wong [9], the quasi-injectives are precisely the fully invariant submodules of injective modules. The corollary then follows on noting that P -primary injective R -modules are direct sums of copies of $R(P^\infty)$ and their proper fully invariant submodules are direct sums of isomorphic cyclic P -primary modules.

The following theorem concludes our investigation of quasi-projectives over Dedekind domains.

THEOREM 5.12. *A quasi-projective module over a Dedekind domain is either torsion or torsion-free.*

Proof. Suppose A is a quasi-projective R -module with its maximal torsion submodule $A_t \neq 0$. Since $R(P^\infty)$ is not quasi-projective for any prime ideal P , A_t is reduced and thus A has torsion cyclic summands [11]. Let $A = (R/P^k) \oplus B$. Now if R is not a complete discrete valuation ring, B/B_t is torsion-free quasi-projective and hence is torsionless (5.2) so that B has a projective summand I of rank 1. If R is a complete discrete valuation ring, then as in the proof of 5.10, one can then show that $B_t = B_P$ is a bounded direct sum of isomorphic cyclic modules, where P is the unique nonzero prime ideal of R . Hence $B = B_P \oplus B/B_P$, so B/B_P is a torsion-free quasi-projective R -module and hence contains a summand isomorphic to R or K , the quotient field of R (5.8). Thus, in either case, A has a summand of the form $(R/P^k) \oplus C$, where $C \cong K$, the quotient field of R or $C \cong I$, an ideal of R . Choose a submodule S of C such that $S \cong R$ or $S \cong IP^k$ according as $C \cong K$ or $C \cong I$. Then there exists a nonzero morphism $g: R/P^k \rightarrow C/S$. Consider the following diagram.

$$\begin{array}{ccc}
 (R/P^k) \oplus C & & \\
 \downarrow g' & & \\
 (R/P^k) \oplus C & \xrightarrow{f'} & (R/P^k) \oplus (C/S)
 \end{array}$$

where $f' = \begin{pmatrix} 1 & 0 \\ 0 & f \end{pmatrix}$, f being the natural map and $g' = \begin{pmatrix} 0 & g \\ 0 & 0 \end{pmatrix}$, where g is any nonzero homomorphism $R/P^k \rightarrow C/S$. This g' cannot be lifted to an endomorphism h of $(R/P^k) \oplus C$ satisfying $h \circ f = g'$, a

contradiction. We thus conclude that A is either torsion or torsion-free.

6. Perfect rings. In this section perfect rings are characterised by means of quasi-projective R -modules.

THEOREM 6.1. *Let R be any ring. Then the following properties are equivalent.*

- (i) R is left perfect.
- (ii) A direct limit of quasi-projective left R -modules is quasi-projective.
- (iii) A direct limit of finitely generated quasi-projectives over R is quasi-projective.
- (iv) Any flat left R -module is quasi-projective.³⁾

Proof. Let $Q = \varinjlim Q_i$, $i \in I$ where I is a directed set and the Q_i 's are quasi-projective R -modules. To each $i \in I$, there exists, by hypothesis, an exact sequence $0 \rightarrow K_i \xrightarrow{u_i} P_i \xrightarrow{v_i} Q_i \rightarrow 0$ where P_i is projective and K_i is small in P_i . Now $\{P_i\}_{i \in I}$ and $\{K_i\}_{i \in I}$ can be made into directed systems in a natural way so that we get a directed system of exact sequences. Let $K = \varinjlim K_i$ and $P = \varinjlim P_i$. Suppose for each $i \in I$ $\alpha_i: P_i \rightarrow P$ and $\beta_i: K_i \rightarrow K$ are the natural maps associated with the direct limits. Since the direct limit commutes with exact sequences, $0 \rightarrow K \xrightarrow{u} P \xrightarrow{v} Q \rightarrow 0$ is exact. We have the following commutative diagram:

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & K_i & \xrightarrow{u_i} & P_i & \xrightarrow{v_i} & Q_i & \longrightarrow & 0 \\
 & & \downarrow \beta_i & & \downarrow \alpha_i & & \downarrow & & \\
 0 & \longrightarrow & K & \xrightarrow{u} & P & \xrightarrow{v} & Q & \longrightarrow & 0.
 \end{array}$$

We claim that Ku is fully invariant in P . Let $f \in \text{End}_R(P)$ and $k \in K$. As R is perfect, P is a direct sum of cyclic projective R -modules [12]. Let P' be a finitely generated summand of P containing $(k)u$ and let $P \xrightarrow{g} P'$ be the natural projection. As $(P')f$ is finitely generated, we can choose a $j \in I$ and a $k_j \in K_j$ such that $(P_j)\alpha_j \supset (P')f$ and $(k_j)\beta_j = k$. Consider the following diagram:

³⁾ In a private communication Dr. J. Golan has indicated that he has also proved the equivalence of (iv) and (i).

$$\begin{array}{ccc}
 & P_j & \\
 & \swarrow h & \downarrow \alpha_j \circ g \circ f \\
 P_j & \longrightarrow & (P_j)\alpha_j
 \end{array}$$

where h exists by the projectivity of P_j . As $(K_j)u_j$ is fully invariant in P_j (by 4.2), $(k_j)u_j h \in (K_j)u_j$. Now

$$\begin{aligned}
 (k)u \circ f &= (k)u \circ g \circ f \text{ (as } g \mid P' = 1_{P'}) = (k_j)\beta_j \circ u \circ g \circ f = (k_j)u_j \circ \alpha_j \circ g \circ f \\
 &= (k_j)u_j \circ h \circ \alpha_j \in (K_j)u_j \circ \alpha_j = (K_j)\beta_j \circ u \cong (K)u.
 \end{aligned}$$

Thus $(K)u$ is fully invariant in P whence $Q \cong P/(K)u$ is quasi-projective.

Clearly (ii) \Rightarrow (iii) and, since a flat module is a direct limit of finitely generated projectives, (iii) implies (iv).

Assume (iv). Let A be flat and P projective such that $A \cong P/S$. Since $A \oplus P$ is flat, it is quasi-projective, by hypothesis. Then Lemma 3.2 implies that A is projective. Thus a direct limit of projective left R -modules is projective and so R is left perfect, by theorem P of [2]. This proves (i).

REMARK. If R is left perfect and A is a quasi-projective left R -module, then a direct sum of any number of carbon copies of A is again quasi-projective. This property, however, does not characterize the perfect rings. Indeed, the investigations made in § 5 show that if R is a countable Dedekind domain which is not a complete discrete valuation ring and A is a quasi-projective R -module, then $\bigoplus_m A$ is quasi-projective for any cardinal number m .

7. Generalization. In this section, we consider a weakened form of quasi-projectivity called w .quasi-projectives. The w .quasi-projective abelian groups were considered in [8]. We give a description of w . quasi-projectives over a Dedekind domain. It is also shown that w .quasi-perfect abelian categories with enough projectives are perfect.

DEFINITION. An object A in a category \mathcal{A} is called *weak quasi-projective* (for short, w . quasi-projective) if for any epimorphism $f: A \rightarrow B$ and any $g: A/B \rightarrow A/B$, there is a $g': A \rightarrow A$ making the following diagram

$$\begin{array}{ccc}
 A & \xrightarrow{f} & A/B \\
 \downarrow g' & & \downarrow g \\
 A & \xrightarrow{f} & A/B
 \end{array}$$

commutative.

It is clear that any quasi-projective is weak quasi-projective. But the converse is not true. The abelian group $Z(P^\infty)$ is *w.* quasi-projective, eventhough it is not a quasi-projective Z -module.

We start with the following lemma which gives a criterion for quasi-projectivity. The proof is straight forward and hence is omitted.

LEMMA 7.1. *An R -module A is quasi-projective if and only if $A \oplus A$ is weak quasi-projective.*

REMARK. It is clear from 7.1 that, unlike the quasi-projective case, if A is *w.* quasi-projective then $A \oplus A$ need not be *w.* quasi-projective.

The next lemma can be obtained by modifying the arguments of 3.2.

LEMMA 7.2. [8]. *If $A \oplus B$ is *w.* quasi-projective and there is an epimorphism $f: A \rightarrow B$, then B will be isomorphic to a summand of A .*

One can define a weak quasi-perfect category in the obvious manner. Using Lemma 7.1 and proceeding exactly as in the proof of Theorem 3.4, we obtain.

THEOREM 7.3. *A weak quasi-perfect abelian category with enough projectives is perfect.*

If we suitably modify the preceding investigation of the quasi-projectives over a Dedekind domain and make use of Lemma 7.2 we can obtain the following theorem whose proof is omitted.

THEOREM 7.4. *Let R be a Dedekind domain.*

(i) *A torsion R -module A is weak quasi-projective if and only if*

each P -primary component A_P is either quasi-projective or $A_P \cong R(P^\infty)$.

(ii) If the number σ of prime ideals of R is $\leq 2^{\aleph_0}$ then the torsion-free weak quasi-projectives are just the (torsion-free) quasi-projectives. If $\sigma > 2^{\aleph_0}$, then a torsionfree weak quasi-projective R -module A is projective if either A has rank $\leq \aleph_0$ or (ii) rank $A > \sigma$. If $\aleph_0 < \text{rank } A < \sigma$, A is \aleph_1 -projective and contains a free summand F whose rank is equal to rank A .

(iii) A properly mixed R -module A is weak quasi-projective if and only if $A \cong B \oplus C$ where B is reduced torsion-free quasi-projective of finite rank and C is an injective submodule of K/R , where K is the quotient field of R .

The authors are indebted to the referee for pointing out a few inaccuracies and for offering many suggestions for improvement.

REFERENCES

1. B. Banaschewski, *Projective Covers in Categories of Topological Spaces and Topological Algebras*, McMaster University, 1968.
2. H. Bass, *Finitistic dimension and a homological generalisation of semiprimary rings*, Trans. Amer. Math. Soc., **95** (1960), 466-488.
3. S. Eilenberg, *Homological dimension and Syzygies*, Ann. of Maths., **64** (1956), 328-336.
4. C. Faith, Review 3817, Math. Reviews, Vol. **36** (1968).
5. P. Freyd, *Abelian Categories*, Harper and Row, New York, (1964).
6. L. Fuchs, *Abelian Groups*, Pergamon Press, New York, 1960.
7. L. Fuchs and K. M. Rangaswamy, *Quasi-projective abelian groups*, Bull. Soc. Math. France, **98** (1970), 5-8.
8. S. Janakiraman, Ph. D. thesis, Madurai University (1970).
9. R. E. Johnson and R. T. Wong, *Quasi-injective modules and irreducible rings*, J. London Math. Soc., **36** (1961), 260-268.
10. I. Kaplansky, *Infinite Abelian Groups*, Ann Arbor, (1956).
11. ———, *Modules over Dedekind domains and valuation rings*, Trans. Amer. Math. Soc., **72** (1952), 327-340.
12. A. Koehler, Ph. D. thesis, Indiana University, (1969).
13. R. J. Nunke, *Modules of extensions over a Dedekind ring*, Illinois J. Math. **3** (1959), 224-242.
14. L. E. T. Wu and J. P. Jans, *On quasi-projectives*, Illinois J. Math. **11** (1967), 439-448.

Received August 1, 1970 and in revised form October 12, 1971. Most of the results in this paper were presented at a seminar held under the chairmanship of Professor N. Jacobson during the Algebra Conference at New Delhi, in April, 1969.

MADURAI UNIVERSITY

AND

THE AUSTRALIAN NATIONAL UNIVERSITY

CANBERRA, A.C.T. 2600, AUSTRALIA

AND

THE UNIVERSITY OF BOMBAY

SANTA CRUZ (EAST), BOMBAY-28, INDIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 43, No. 1

March, 1972

Alexander (Smbat) Abian, <i>The use of mitotic ordinals in cardinal arithmetic</i>	1
Helen Elizabeth Adams, <i>Filtrations and valuations on rings</i>	7
Benno Artmann, <i>Geometric aspects of primary lattices</i>	15
Marilyn Breen, <i>Determining a polytope by Radon partitions</i>	27
David S. Browder, <i>Derived algebras in L_1 of a compact group</i>	39
Aiden A. Bruen, <i>Unimbeddable nets of small deficiency</i>	51
Michael Howard Clapp and Raymond Frank Dickman, <i>Unicoherent compactifications</i>	55
Heron S. Collins and Robert A. Fontenot, <i>Approximate identities and the strict topology</i>	63
R. J. Gazik, <i>Convergence in spaces of subsets</i>	81
Joan Geramita, <i>Automorphisms on cylindrical semigroups</i>	93
Kenneth R. Goodearl, <i>Distributing tensor product over direct product</i>	107
Julien O. Hennefeld, <i>The non-conjugacy of certain algebras of operators</i>	111
C. Ward Henson, <i>The nonstandard hulls of a uniform space</i>	115
M. Jeanette Huebener, <i>Complementation in the lattice of regular topologies</i>	139
Dennis Lee Johnson, <i>The diophantine problem $Y^2 - X^3 = A$ in a polynomial ring</i>	151
Albert Joseph Karam, <i>Strong Lie ideals</i>	157
Soon-Kyu Kim, <i>On low dimensional minimal sets</i>	171
Thomas Latimer Kriete, III and Marvin Rosenblum, <i>A Phragmén-Lindelöf theorem with applications to $\mathcal{M}(u, v)$ functions</i>	175
William A. Lampe, <i>Notes on related structures of a universal algebra</i>	189
Theodore Windle Palmer, <i>The reducing ideal is a radical</i>	207
Kulumani M. Rangaswamy and N. Vanaja, <i>Quasi projectives in abelian and module categories</i>	221
Ghulam M. Shah, <i>On the univalence of some analytic functions</i>	239
Joseph Earl Valentine and Stanley G. Wayment, <i>Criteria for Banach spaces</i>	251
Jerry Eugene Vaughan, <i>Linearly stratifiable spaces</i>	253
Zbigniew Zielezny, <i>On spaces of distributions strongly regular with respect to partial differential operators</i>	267