

Pacific Journal of Mathematics

CRITERIA FOR BANACH SPACES

JOSEPH EARL VALENTINE AND STANLEY G. WAYMENT

CRITERIA FOR BANACH SPACES

J. E. VALENTINE AND S. G. WAYMENT

It is well known in euclidean geometry that the quadrilateral obtained from an arbitrary quadrilateral by joining its midpoints is a parallelogram. The purpose of this paper is to show that a complete metric space with a unique metric line joining any pair of its distinct points is a Banach space if and only if it has the above mentioned property.

Let $p, q, r,$ and s be distinct points in a Banach space such that no three are linear let $m_1, m_2, m_3,$ and m_4 be the midpoints of the algebraic segments joining p and q, q and r, r and $s,$ and s and $p,$ respectively. It is well known that $m_3 - m_2 = m_4 - m_1$ and $m_2 - m_1 = m_3 - m_4$. In Euclidean space one usually refers to this result by saying that the midpoints m_1, m_2, m_3, m_4 form a parallelogram. If the Banach space does not have unique segments joining pairs of distinct points, then the restriction that the $\{m_i\}$ be midpoints of algebraic segments is easily seen to be necessary. We shall say that the metric space M satisfies the quadrilateral midpoint postulate provided that if p, q, r, s are points of M such that no three are linear and if m_1, m_2, m_3, m_4 are the respective midpoints, then $m_1m_2 = m_3m_4$ and $m_2m_3 = m_1m_4$. Hereafter we shall assume that M is a complete metric space with a unique metric line joining any pair of its distinct points and show that the Quadrilateral Midpoint Postulate characterizes the class of Banach spaces among such metric spaces.

The technique will be to show that a complete metric space with a unique metric line joining any pair of its distinct points satisfies the Quadrilateral Midpoint Postulate if and only if it satisfies the Young Postulate which may be stated as follows.

The Young Postulate. If p, q, r are points of a metric space M and q' and r' are the midpoints of p and $q,$ and p and $r,$ respectively, then $q'r' = qr/2$.

The result will then follow, for Andalafte and Blumenthal [1] have shown that a complete metric space with a unique metric line joining any pair of its distinct points is a Banach space if and only if it satisfies the Young Postulate.

That the Young Postulate implies the Quadrilateral Midpoint Postulate is almost immediate. For if a complete metric space with a metric line joining any pair of its distinct points satisfies the Young Postulate, then it is a Banach space and consequently satisfies the Quadrilateral Midpoint Postulate.

Suppose M satisfies the Quadrilateral Midpoint Postulate and p, q, r , are non-linear points of M with m_1, m_2 the midpoints of p and q, q and r , respectively.

LEMMA 1. *There exists a number k , depending only on p and r , such that if q, m_1, m_2 are as above, then $m_1 m_2 = kpr$.*

Proof. Let s be a point such that no three of p, q, r, s are collinear, and let m_3, m_4 be the midpoints of the segments joining r and s, s and p , respectively. Let $k = m_3 m_4 / pr$. Then since M satisfies the quadrilateral midpoint property, $m_1 m_2 = m_3 m_4 = kpr$. We see immediately that k does not depend on q .

LEMMA 2. *The k in Lemma 1 is $1/2$.*

Proof. Let $\{x_i\}$ be a sequence of points tending to x on the segment between p and r with $p \neq x \neq r$ and such that for each i we have p, x_i, r non-collinear. Let $\{p_i\}$ and $\{r_i\}$ be the sequences such that p_i and r_i are the midpoints of the segments determined by p and x_i, r and x_i, r and x_i , respectively. Then $\lim p_i x_i = 1/2 \lim px_i = 1/2 px$ and similarly $\lim r_i x_i = 1/2 rx$. This, along with the triangle inequality $p_i x_i + x_i r_i \geq p_i r_i = kpr$, implies $k \leq 1/2$. However, the inequality $pr \leq pp_i + p_i r_i + r_i r = pp_i + kpr + r_i r$ and the aforementioned limits imply $k \geq 1/2$. Hence k is $1/2$.

THEOREM. *A complete metric space with a unique line joining any two of its distinct points is a normed linear space (Banach Space) if and only if it satisfies the Quadrilateral Midpoint Postulate.*

Proof. We have shown that the Quadrilateral Midpoint Postulate implies the Young Postulate; that is, if p' and r' are midpoints of p and q , and q and r , respectively, then $p'r' = (1/2)pr$. Thus an application of the Andalafte-Blumenthal result [1] completes the proof.

REFERENCES

1. E. Z. Andalafte and L. M. Blumenthal, *Metric characterization of banach and euclidean spaces*, Fund. Math., LV (1964), 23-55.

Received August 16, 1971.

UTAH STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 43, No. 1

March, 1972

Alexander (Smbat) Abian, <i>The use of mitotic ordinals in cardinal arithmetic</i>	1
Helen Elizabeth Adams, <i>Filtrations and valuations on rings</i>	7
Benno Artmann, <i>Geometric aspects of primary lattices</i>	15
Marilyn Breen, <i>Determining a polytope by Radon partitions</i>	27
David S. Browder, <i>Derived algebras in L_1 of a compact group</i>	39
Aiden A. Bruen, <i>Unimbeddable nets of small deficiency</i>	51
Michael Howard Clapp and Raymond Frank Dickman, <i>Unicoherent compactifications</i>	55
Heron S. Collins and Robert A. Fontenot, <i>Approximate identities and the strict topology</i>	63
R. J. Gazik, <i>Convergence in spaces of subsets</i>	81
Joan Geramita, <i>Automorphisms on cylindrical semigroups</i>	93
Kenneth R. Goodearl, <i>Distributing tensor product over direct product</i>	107
Julien O. Hennefeld, <i>The non-conjugacy of certain algebras of operators</i>	111
C. Ward Henson, <i>The nonstandard hulls of a uniform space</i>	115
M. Jeanette Huebener, <i>Complementation in the lattice of regular topologies</i>	139
Dennis Lee Johnson, <i>The diophantine problem $Y^2 - X^3 = A$ in a polynomial ring</i>	151
Albert Joseph Karam, <i>Strong Lie ideals</i>	157
Soon-Kyu Kim, <i>On low dimensional minimal sets</i>	171
Thomas Latimer Kriete, III and Marvin Rosenblum, <i>A Phragmén-Lindelöf theorem with applications to $\mathcal{M}(u, v)$ functions</i>	175
William A. Lampe, <i>Notes on related structures of a universal algebra</i>	189
Theodore Windle Palmer, <i>The reducing ideal is a radical</i>	207
Kulumani M. Rangaswamy and N. Vanaja, <i>Quasi projectives in abelian and module categories</i>	221
Ghulam M. Shah, <i>On the univalence of some analytic functions</i>	239
Joseph Earl Valentine and Stanley G. Wayment, <i>Criteria for Banach spaces</i>	251
Jerry Eugene Vaughan, <i>Linearly stratifiable spaces</i>	253
Zbigniew Zielezny, <i>On spaces of distributions strongly regular with respect to partial differential operators</i>	267