

# Pacific Journal of Mathematics

**ZEROS OF SUMS OF SERIES WITH HADAMARD GAPS**

LINDA RUTH SONS

## ZEROS OF SUMS OF SERIES WITH HADAMARD GAPS

L. R. SONS

If  $f$  is a function of the complex variable  $z$  in the unit disc and the power series expansion for  $f$  about zero can be expressed as a finite sum of series with Hadamard gaps, then  $f(z)$  assumes every finite value infinitely often provided the coefficients in the power series expansion of  $f$  do not tend to zero and the average value of  $(\log^+ 1/|f(re^{i\theta})|)^p$  does not grow too rapidly as  $r \rightarrow 1^-$  for some  $p > 1$ .

1. Introduction and statement of results. Let  $f$  be a function analytic in  $|z| < 1$  for which

$$(1) \quad f(z) = c_0 + \sum_{k=1}^{\infty} c_k z^{n_k}$$

where  $\{n_k\}$  is a sequence of positive integers for which

$$(2) \quad \frac{n_{k+1}}{n_k} \geq q > 1, \quad (k \geq 1).$$

The series in (1) is said to have Hadamard gaps.

If  $q$  is greater than about 100, G. and M. Weiss [9] proved  $f(z)$  assumes every finite value infinitely often provided

$$(3) \quad \sum_{k=0}^{\infty} |c_k| = \infty.$$

If  $q > 1$ , W. H. J. Fuchs [2] showed  $f(z)$  assumes every finite value infinitely often provided

$$(4) \quad \limsup_{k \rightarrow \infty} |c_k| > 0.$$

In [3] Fuchs has extended his result to show that  $f$  assumes every finite value infinitely often in each sector

$$S = \{z | \alpha < \arg z < \beta \text{ and } |z| < 1\}$$

where  $\alpha$  and  $\beta$  are fixed real numbers.

The original result of Fuchs may also be extended as follows:

**THEOREM 1.** *Let  $\{n_k\}$  be a sequence of positive integers for which (2) holds. Let  $l$  be a fixed positive integer, and let  $n_k^{(i)}$  for  $i=1, 2, \dots, l$  be integers for which*

$$n_{k-1} < n_k^{(1)} < n_k^{(l-1)} < \dots < n_k^{(1)} < n_k.$$

Suppose  $f$  is a function analytic in  $|z| < 1$  for which

$$(5) \quad \begin{aligned} f(z) &= a_0 + \sum_{k=1}^{\infty} (a_{n_k^{(l)}} z^{n_k^{(l)}} + a_{n_k^{(l-1)}} z^{n_k^{(l-1)}} + \cdots + a_{n_k^{(1)}} z^{n_k^{(1)}} + a_{n_k} z^{n_k}), \\ &= \sum_{k=0}^{\infty} c_k z^k. \end{aligned}$$

Suppose (4) holds and for some  $p > 1$  there exists a constant  $C$  with  $0 < C < +\infty$  such that

$$(6) \quad \frac{1}{2\pi} \int_0^{2\pi} (\log^+ |f(re^{i\theta})|)^p d\theta \leq C \left( \frac{1}{2\pi} \int_0^{2\pi} (\log^+ |f(re^{i\theta})|) d\theta \right)^p$$

for a sequence of values of  $r$  approaching one. Then  $f(z)$  assumes every finite value infinitely often in  $|z| < 1$ .

Two immediate corollaries of Theorem 1 are:

**COROLLARY 1.** Assume the hypothesis of Theorem 1 with  $n_k^{(\alpha)} = n_k - \alpha$  for  $k = 1, 2, 3, \dots$  and  $0 < \alpha \leq l$ . Then  $f(z)$  assumes every finite value infinitely often in  $|z| < 1$ .

**COROLLARY 2.** Let  $f$  be a function analytic in  $|z| < 1$  for which

$$f(z) = \sum_{k=0}^{\infty} c_k z^k = f_0(z) + f_1(z) + \cdots + f_l(z)$$

where for each  $i$ ,  $f_i(z)$  has a power series expansion about zero with Hadamard gaps. If (4) holds and for some  $p > 1$  there exists a constant  $C$  with  $0 < C < +\infty$  such that (6) holds for a sequence of values of  $r$  approaching one, then  $f(z)$  assumes every finite value infinitely often in  $|z| < 1$ .

Corollary 1 is a special case of Theorem 1 and extends a result of C. Pommerenke [6] who showed that functions of the type of Corollary 1 without the assumption (6) must assume every value at least once. (G. Schmeisser [7] has recently extended the method in [6] to show the Pommerenke-type series assume every value infinitely often). Corollary 2 follows from Theorem 1 by noticing that  $f(z)$  can be rewritten, if necessary, to be in the form (5).

For functions of the form (5) for which

$$\limsup_{r \rightarrow 1^-} \frac{\log M(r)}{-\log(1-r)} > 2(2+l)$$

where  $M(r)$  denotes the maximum modulus of  $f(z)$  on  $|z| = r$ , we remark in [8] that

$$\limsup_{r \rightarrow 1^-} \frac{n(r)}{-\log(1-r)} > 0,$$

where  $n(r)$  denotes the number of zeros of  $f$  in  $|z| \leq r$ . It seems probable that functions of the type of Theorem 1 also assume every finite value infinitely often in each sector

$$\{z | \alpha < \arg z < \beta \quad \text{and} \quad |z| < 1\}$$

where  $\alpha$  and  $\beta$  are fixed real numbers. It has been shown by P. Erdos and A. Renyi [1] that if  $\{n_k\}$  is an increasing sequence of natural numbers satisfying

$$\liminf_{(k-j) \rightarrow \infty} (n_k - n_j)^{1/(k-j)} = 1,$$

then, for any sequence  $\{\omega_k\}$  of natural numbers for which

$$\lim_{k \rightarrow \infty} \omega_k = +\infty$$

there exists a sequence  $\{m_k\}$  of natural numbers such that

$$0 \leq m_k - n_k < \omega_k$$

and a function  $g$ , analytic in  $|z| < 1$  with the power series expansion

$$g(z) = \sum_{k=0}^{\infty} b_k z^{m_k},$$

where the  $b_k$  are positive, such that  $g(z)$  is unbounded in  $|z| < 1$ , but bounded in the domain  $|z| < 1, |\arg z| > \varepsilon$ , for any  $\varepsilon > 0$ .

If  $f$  is an analytic function in  $|z| < 1$ , D. Gaier and W. Meyer-König [5] have defined the radius  $R_\varphi$  defined by  $z = re^{i\varphi}$ ,  $0 \leq r < 1$ , *singular* for  $f$  if  $f(z)$  is unbounded in any sector  $|z| < 1, \varphi - \varepsilon < \arg z < \varphi + \varepsilon$  with  $\varepsilon > 0$ . They showed that if  $f$  is unbounded in  $|z| < 1$  and the power series expansion for  $f$  about zero has Hadamard gaps, then every radius is singular for  $f$ . We have

**THEOREM 2.** *Suppose  $f$  is a function which is analytic in  $|z| < 1$  and has the power series expansion (5). Suppose*

$$\limsup_{r \rightarrow 1^-} \{\max |c_k| r^k\} = \infty.$$

*Then each radius  $R_\varphi (0 \leq \varphi < 2\pi)$  is singular for  $f$ .*

In section two the necessary lemmas are stated and the theorems are proved, while section three contains the proof of the essential lemma which enables us to use the idea of G. H. Hardy and J. E. Littlewood of accentuating the dominance of the largest term in the series (5) by repeated differentiation (c.f. Fuchs [2]).

**2. Proofs of the Theorems.** We need three lemmas:

**LEMMA 1 (Fuchs [2]).** *Let  $g$  be a function analytic in  $|z| < R$ . If, for some positive integer  $p$*

$$|g^{(p)}(z)| \leq M \quad (|z| < R),$$

and

$$|g^{(p)}(0)| \geq A > 0,$$

then  $g(z)$  assumes in  $|z| < R$  every value  $w$  lying in the disc

$$|w - g(0)| < KR^p A^{p+1} M^{-p}$$

where  $K$  is a positive number depending only on  $p$ .

**LEMMA 2 (Gaier [4]).** *Let  $E$  be a closed subset of  $\{z \mid |z| = 1\}$  and assume that  $E$  has measure  $2\pi\gamma$  where  $0 < \gamma < 1$ . If  $p$  is a polynomial with  $N$  terms, then*

$$\max_{|z|=1} |p(z)| \leq C_N(\gamma) \cdot \max_{z \in E} |p(z)|$$

where

$$(7) \quad \log C_N(\gamma) = \left( \frac{\gamma}{1-\gamma} \gamma^{-N} - \frac{1}{1-\gamma} \right) \log 3.$$

**LEMMA 3.** *Assume the hypothesis of Theorem 1. Let  $p, \nu$ , and  $\alpha$  be positive integers where  $0 \leq \alpha \leq l$ . For  $k = 1, 2, 3, \dots$  let  $n_k^{(0)} = n_k$ . Define*

$$S_0 = \exp\{-p/n_\nu^{(\alpha)}\}, \quad S_1 = \exp\left\{-\frac{P}{2n_\nu^{(\alpha)}}\left(1 + \frac{\log q}{q-1}\right)\right\},$$

and

$$W_{k,(\beta)} = n_k^{(\beta)}(n_k^{(\beta)} - 1)(n_k^{(\beta)} - 2) \dots (n_k^{(\beta)} - p + 1) S_k^{n_k^{(\beta)}}$$

where  $S_0 < S < S_1$ ;  $0 \leq \beta \leq l$ ; and  $k = 1, 2, 3, \dots$ . Then for a fixed  $\gamma$  with  $0 < \gamma < 1$  there exists an integer  $p_0$  depending on  $q$  and  $l$  such that for  $p > p_0$  and  $\nu > \nu_0(p)$

$$\sum_{k \neq \nu-1, \nu, \nu+1} (W_{k,(l)} + W_{k,(l-1)} + \dots + W_{k,(0)}) < \frac{1}{4} (C_{3l+3}(\gamma))^{-1} W_{\nu,(\alpha)},$$

where  $C_{3l+3}(\gamma)$  is defined by (7).

*Proof of Theorem 1.* We note that it suffices to show  $f(z)$  assumes zero infinitely often in  $|z| < 1$ . So suppose  $f(z)$  is zero only a finite number of times in  $|z| < 1$  and denote these zeros by  $z_1, z_2, \dots, z_j$ . Then  $N(r, 1/f) = O(1)$ , and it follows from the first fundamental theorem of Nevanlinna theory that

$$(8) \quad m(r, f) = m(r, 1/f) + O(1) .$$

For  $0 < r < 1$ ,

$$|f(re^{i\theta})| > \lambda_q \left( \sum_{k=0}^{\infty} |c_k|^2 r^{2k} \right)^{1/2} > 0$$

on a set of  $\theta$  of measure not less than  $\mu_q > 0$  [10, p. 216<sup>1</sup>], so

$$(9) \quad m(r, f) \rightarrow \infty$$

as  $r$  approaches one by condition (4).

For a value  $r'$  for which (6) holds, let  $\mathcal{E}(r')$  denote the set of  $\theta$  in  $[0, 2\pi]$  at which

$$(10) \quad \log^+ \frac{1}{|f(r'e^{i\theta})|} > \frac{1}{2} m(r', 1/f) .$$

Denote the measure of  $\mathcal{E}(r')$  by  $|\mathcal{E}(r')|$ . Then using Hölder's inequality and (6)

$$\begin{aligned} \pi m(r', 1/f) &\leq \int_{\mathcal{E}(r')} \log^+ \frac{1}{|f(r'e^{i\theta})|} d\theta \\ &\leq \left( \int_{\mathcal{E}(r')} \left( \log^+ \frac{1}{|f(r'e^{i\theta})|} \right)^p d\theta \right)^{1/p} |\mathcal{E}(r')|^{1/q} \\ &\leq (2\pi C)^{1/p} m(r', 1/f) |\mathcal{E}(r')|^{1/q} . \end{aligned}$$

Thus,

$$(\pi/(2\pi C)^{1/p})^q \leq |\mathcal{E}(r')| .$$

Define  $\gamma$  by

$$2\pi\gamma = (\pi/(2\pi C)^{1/p})^q .$$

Let  $\rho = \max_{1 \leq k \leq j} |z_k|$ , and let

$$U = \limsup_{k \rightarrow \infty} |c_k| .$$

If  $U < \infty$ , let  $N$  be the least integer such that

$$|c_k| < \frac{3}{2} U$$

for  $k > N$ . If  $U = \infty$ , let  $N = 0$ .

Define

$$\mu(r) = \sup_{k > N} |c_k| r^k , \quad (0 \leq r < 1) .$$

Let  $V = V(r)$  be the largest integer such that

---

<sup>1</sup> Theorem 8.20 on page 215 easily extends to finite sums of series with Hadamard gaps, and so Theorem 8.25 on page 216 does also.

$$|c_\nu| r^\nu > \frac{1}{2} \mu(r) .$$

If  $U = \infty$ , we see

$$|c_\nu| r^\nu > 1 , \quad (r > r_0) ,$$

and also  $V(r) \rightarrow \infty$  as  $r \rightarrow 1^-$ . If  $U < \infty$ , we see

$$|c_\nu| r^\nu > \frac{1}{3} U$$

and again  $V(r) \rightarrow \infty$  as  $r \rightarrow 1^-$  since there are infinitely many integers  $k$  with

$$|c_k| > \frac{3}{4} U > \frac{1}{2} \mu(r) , \quad (r < 1) .$$

Using the notation of Lemmas 1 and 3, we choose  $p \geq \max(N, p_0)$  and choose  $r$  so close to one that

$$\rho < rS_0 = r \exp \{-p/n_\nu^{(\alpha)}\}$$

where  $n_\nu^{(\alpha)} = V(r)$ ,  $n_{\nu-1}^{(l)} > 2p$ , and  $\nu > \nu_0(p)$ . We may assume  $r(S_0 S_1)^{1/2}$  is a value  $r'$  for which (6) holds. Let

$$T(z) = \sum_{k=\nu-1}^{\nu+1} (a_{n_k^{(l)}} z^{n_k^{(l)}} + \dots + a_{n_k} z^{n_k}) .$$

By Lemma 3

$$\begin{aligned} \sum_{k \neq \nu-1, \nu, \nu+1} \left( \sum_{i=0}^l |a_{n_k^{(i)}}| r^{n_k^{(i)}} W_{k, (i)} \right) &\leq \mu(r) \sum_{k \neq \nu-1, \nu, \nu+1} (W_{k, (l)} + \dots + W_{k, (0)}) , \\ &\leq 2 |a_{n_\nu^{(\alpha)}}| r^{n_\nu^{(\alpha)}} \cdot \frac{1}{4} (C_{3l+3}(\gamma))^{-1} W_{\nu, (\alpha)} . \end{aligned}$$

Hence,

$$f^{(p)}(rSe^{i\theta}) = T^{(p)}(rSe^{i\theta}) + E(rSe^{i\theta})$$

where

$$|E(rSe^{i\theta})| < \frac{1}{2} (C_{3l+3}(\gamma))^{-1} n_\nu^{(\alpha)} (n_\nu^{(\alpha)} - 1) \dots (n_\nu^{(\alpha)} - p + 1) |a_{n_\nu^{(\alpha)}}| (rS)^{n_\nu^{(\alpha)} - p} .$$

Consequently,

$$|f^{(p)}(rSe^{i\theta})| \leq C_1(p, l) (n_\nu^{(\alpha)})^p |a_{n_\nu^{(\alpha)}}| (rS_1)^{n_\nu^{(\alpha)}} ,$$

and using Lemma 2 on the polynomial  $T^{(p)}(rSe^{i\theta})$  where  $rS$  is a value  $r'$  for which (6) is valid we find

$$|f^{(p)}(rSe^{i\theta})| \geq C_2(p, l, \gamma) (n_\nu^{(\alpha)})^p |a_{n_\nu^{(\alpha)}}| (rS_0)^{n_\nu^{(\alpha)}}$$

for values of  $\theta$  in  $\mathcal{E}(rs)$ .

Therefore if  $r' = r(S_0S_1)^{1/2}$  is a value for which (6) holds, then we may apply Lemma 1 to

$$g(\zeta) = f(r(S_0S_1)^{1/2}e^{\zeta+i\theta})$$

with

$$R = \frac{1}{4} \left( 1 + \frac{\log q}{q-1} \right) \frac{p}{n_\nu^{(\alpha)}}$$

where  $p$  and  $\nu$  satisfy the hypotheses of Lemma 3. Then

$$\begin{aligned} R^p A^{p+1} M^{-p} &> C_3(p, q, l, \gamma) |a_{n_\nu^{(\alpha)}}| (rS_0)^{n_\nu^{(\alpha)}} (S_0/S_1)^{pn_\nu^{(\alpha)}}, \\ &> C_4(p, q, l, \gamma) \mu(r), \\ &> C_5(p, q, l, \gamma, U). \end{aligned}$$

Thus  $f(z)$  takes every value  $w$  in the disc

$$|w - f(r(S_0S_1)^{1/2}e^{i\theta})| < C_6(p, q, l, \gamma, U).$$

But by (10) we note that

$$|f(r(S_0S_1)^{1/2}e^{i\theta})| < \exp\left(-\frac{1}{2}m(r(S_0S_1)^{1/2}, 1/f)\right),$$

and because of (8) and (9) we conclude that when  $r$  is near enough to one

$$|f(r(S_0S_1)^{1/2}e^{i\theta})| < C_6(p, q, l, \gamma, U).$$

Thus  $f(z)$  will assume the value zero at points arbitrarily near  $|z| = 1$  which contradicts our earlier assumption and proves the theorem.

*Proof of Theorem 2.* Suppose there is some radius  $R_\varphi$  which is not singular for  $f$ , and so there exists an  $\varepsilon > 0$  such that  $|f(z)|$  is bounded in the sector  $\mathcal{S} = \{z | \varphi - \varepsilon < \arg z < \varphi + \varepsilon, |z| < 1\}$ . Then for each complex number  $a$ ,  $f(z) - a$  is also bounded in  $\mathcal{S}$ . Thus taking  $2\pi\gamma = 2\varepsilon$ , the argument of Theorem 1 shows  $f$  assumes  $a$  infinitely often in  $\mathcal{S}$ . Since  $a$  is arbitrary,  $|f|$  is unbounded in  $\mathcal{S}$ , and therefore  $R_\varphi$  is singular for  $f$ .

**3. Proof of Lemma 3.** If  $n_k^{(\beta)} < p$ , then  $W_{k,(\beta)} = 0$ . Turning to  $p \leq n_k^{(\beta)} < n_\nu^{(\alpha)}$ , we first observe that for fixed  $\beta$  with  $0 \leq \beta \leq l$ ,

$$\frac{n_{k+2}^{(\beta)}}{n_k^{(\beta)}} \geq q > 1, \quad (k = 1, 2, 3, \dots).$$

Assume  $W_{k+2,(\beta)} \neq 0$  and  $n_{k+2}^{(\beta)} \leq n_\nu^{(\alpha)}$ . Then



$$\begin{aligned}
 (11) \quad \frac{W_{k,(\beta)}}{W_{k+2,(\beta)}} &\leq \left(\frac{n_k^{(\beta)}}{n_{k+2}^{(\beta)}}\right)^p (S_0)^{-n_{k+2}^{(\beta)}+n_k^{(\beta)}} \leq \left(\frac{n_k^{(\beta)}}{n_{k+2}^{(\beta)}}\right)^p \exp\left\{p\left(1 - \frac{n_k^{(\beta)}}{n_{k+2}^{(\beta)}}\right)\right\}, \\
 &< \sup_{0 < t < 1/q} \exp\{p(1 - t + \log t)\} \\
 &\leq \exp\{p(1 - q) - \log q\}.
 \end{aligned}$$

Hence the right-hand side of (11) is less than  $A^{-1} = (1 + 16(l + 1)C_{3l+3}(\gamma))^{-1}$  provided

$$p > \frac{\log A}{\log q - 1 + q^{-1}} = p_1.$$

Proceeding in a similar manner, we may also show

$$\frac{W_{\nu-3,(\beta)}}{W_{\nu,(\alpha)}} \leq \frac{1}{A} \quad \text{and} \quad \frac{W_{\nu-2,(\beta)}}{W_{\nu,(\alpha)}} \leq \frac{1}{A}$$

when  $p > p_1$ . Consequently for  $W_{k,(\beta)} \neq 0$  where  $k \leq \nu - 3$  and  $k + 2n = \nu - 1$ , we see

$$(12) \quad \frac{W_{k,(\beta)}}{W_{k+2,(\beta)}} \cdot \frac{W_{k+2,(\beta)}}{W_{k+4,(\beta)}} \dots \frac{W_{\nu-5,(\beta)}}{W_{\nu-3,(\beta)}} \frac{W_{\nu-3,(\beta)}}{W_{\nu,(\alpha)}} \leq \left(\frac{1}{A}\right)^n, \quad (0 \leq \beta \leq l),$$

and for  $W_{k,(\beta)} \neq 0$  where  $k \leq \nu - 2$  and  $k + 2n = \nu$  we see

$$(13) \quad \frac{W_{k,(\beta)}}{W_{k+2,(\beta)}} \cdot \frac{W_{k+2,(\beta)}}{W_{k+4,(\beta)}} \dots \frac{W_{\nu-4,(\beta)}}{W_{\nu-2,(\beta)}} \cdot \frac{W_{\nu-2,(\beta)}}{W_{\nu,(\alpha)}} \leq \left(\frac{1}{A}\right)^n, \quad (0 \leq \beta \leq l).$$

Using (12) and (13) provided  $p > p_1$ , we get

$$\begin{aligned}
 (14) \quad \sum_{k < \nu-1} (W_{k,(l)} + W_{k,(l-1)} + \dots + W_{k,(0)}) &< \frac{2(l+1)}{A} \sum_{n=0}^{\infty} \left(\frac{1}{A}\right)^n W_{\nu,(\alpha)}, \\
 &\leq \frac{2(l+1)}{A-1} W_{\nu,(\alpha)}.
 \end{aligned}$$

Now for  $k \geq \nu > \nu_0(p)$  and  $x$  any integer with  $0 < x \leq p - 1$ , we have simultaneously

$$\frac{n_{k+2}^{(\beta)} - x}{n_k^{(\beta)} - x} < 2^{1/p} \left(\frac{n_{k+2}^{(\beta)}}{n_k^{(\beta)}}\right), \quad \frac{n_{\nu+2}^{(\beta)} - x}{n_{\nu}^{(\alpha)} - x} < 2^{1/p} \left(\frac{n_{\nu+2}^{(\beta)}}{n_{\nu}^{(\alpha)}}\right),$$

and

$$\frac{n_{\nu+3}^{(\beta)} - x}{n_{\nu}^{(\alpha)} - x} < 2^{1/p} \left(\frac{n_{\nu+3}^{(\beta)}}{n_{\nu}^{(\alpha)}}\right).$$

Then when  $n_k^{(\beta)} \geq n_{\nu}^{(\alpha)}$ ,

$$\begin{aligned}
 \frac{W_{k+2,(\beta)}}{W_{k,(\beta)}} &< 2 \left(\frac{n_{k+2}^{(\beta)}}{n_k^{(\beta)}}\right)^p S_{k+2}^{n_{k+2}^{(\beta)}-n_k^{(\beta)}}, \\
 &< 2t^p (S_1^{n_k^{(\beta)}})^{(t-1)},
 \end{aligned}$$

$$\leq 2t^p \left\{ \exp \left( -\frac{1}{2} \left( 1 + \frac{\log q}{q-1} \right) \right) \right\}^{p(t-1)} = \theta(t)$$

where  $t = n_{k+2}^{(\beta)} / n_k^{(\beta)} \geq q$ . For  $t \geq q$ ,  $\theta(t) \leq \theta(q)$ , so

$$(15) \quad \frac{W_{k+2,(\beta)}}{W_{k,(\beta)}} \leq 2 \exp \left\{ -\frac{1}{2} p(q-1-\log q) \right\},$$

when  $n_k^{(\beta)} \geq n_\nu^{(\alpha)}$ . Hence the right-hand side of (15) is less than  $1/A$  provided

$$p > \frac{2 \log(2A)}{q-1-\log q} = p_2.$$

Proceeding in a similar manner we may also show

$$\frac{W_{\nu+2,(\beta)}}{W_{\nu,(\alpha)}} \leq \frac{1}{A} \quad \text{and} \quad \frac{W_{\nu+3,(\beta)}}{W_{\nu,(\alpha)}} \leq \frac{1}{A},$$

when  $p > p_2$ . Consequently for  $k \geq \nu + 2 \geq \nu_0(p) + 2$  and  $k = \nu + 2n$ , we see

$$(16) \quad \frac{W_{k,(\beta)}}{W_{k-2,(\beta)}} \cdot \frac{W_{k-2,(\beta)}}{W_{k-4,(\beta)}} \cdots \frac{W_{\nu+4,(\beta)}}{W_{\nu+2,(\beta)}} \frac{W_{\nu+2,(\beta)}}{W_{\nu,(\alpha)}} \leq \left( \frac{1}{A} \right)^n, \quad (0 \leq \beta \leq l),$$

and for  $k \geq \nu + 3 > \nu_0(p) + 3$  and  $k = \nu + 2n + 1$ , we see

$$(17) \quad \frac{W_{k,(\beta)}}{W_{k-2,(\beta)}} \cdot \frac{W_{k-2,(\beta)}}{W_{k-4,(\beta)}} \cdots \frac{W_{\nu+5,(\beta)}}{W_{\nu+3,(\beta)}} \frac{W_{\nu+3,(\beta)}}{W_{\nu,(\alpha)}} \leq \left( \frac{1}{A} \right)^n, \quad (0 \leq \beta \leq l).$$

Using (16) and (17) provided  $p > p_2$ , we get

$$(18) \quad \sum_{k > \nu+1} (W_{k,(l)} + W_{k,(l-1)} + \cdots + W_{k,(0)}) < \frac{2(l+1)}{A} \sum_{n=0}^{\infty} \left( \frac{1}{A} \right)^n W_{\nu,(\alpha)}, \\ \leq \frac{2(l+1)}{A-1} W_{\nu,(\alpha)}.$$

Combining (14) and (18) we now have the lemma provided  $p_0$  is the maximum of  $p_1$  and  $p_2$  (and remembering that  $A = 1 + 16(l+1)C_{3l+3}(\gamma)$ ).

The authoress is grateful to the referee for his helpfulness.

#### REFERENCES

1. P. Erdős and A. Rényi, *On singular radii of power series*, Publications Math. Inst. Hungarian Acad. Sciences, **3** (1959), 159-169.
2. W. H. J. Fuchs, *On the zeros of power series with Hadamard gaps*, Nagoya Math. J., **29** (1967), 167-174.
3. ———, *Topics in Nevanlinna theory*, Proceedings of the NRL conference on classical function theory, (1970), 1-32.

4. D. Gaier, *Bemerkungen zum Turánschen Lemma*, Abh. Math. Sem. Univ. Hamburg, **35** (1970), 1-7.
5. D. Gaier and W. Meyer-König, *Singulare Radian bei Potenzreihen*, Jahresbericht Deutschen Math. Ver., **59** (1956), 36-48.
6. C. Pommerenke, *Lacunary power series and univalent functions*, Michigan Math. J., **11** (1964), 219-233.
7. G. Schmeisser, *Zur Theorie der Potenzreihen mit Lücken*, Math. Z., **126** (1972), 40-46.
8. L. R. Sons, *Zeros of power series with large gaps*, Indiana Univ. Math. J., **22** (1972), 199-205.
9. G. Weiss and M. Weiss, *On the Picard property of lacunary power series*, Studia Math., **22** (1963), 221-245.
10. A. Zygmund, *Trigonometric Series*, Vol. 1, Cambridge University Press.

Received September 2, 1970 and in revised form August 4, 1972. Research supported by NSF grant GP-16548.

IMPERIAL COLLEGE OF SCIENCE AND TECHNOLOGY  
AND  
NORTHERN ILLINOIS UNIVERSITY

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON

Stanford University  
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

C. R. HOBBY

University of Washington  
Seattle, Washington 98105

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

# Pacific Journal of Mathematics

Vol. 43, No. 2

April, 1972

Arne P. Baartz and Gary Glenn Miller, <i>Souslin's conjecture as a problem on the real line</i> . . . . .	277
Joseph Barback, <i>On solutions in the regressive isols</i> . . . . .	283
Barry H. Dayton, <i>Homotopy and algebraic K-theory</i> . . . . .	297
William Richard Derrick, <i>Weighted convergence in length</i> . . . . .	307
M. V. Deshpande and N. E. Joshi, <i>Collectively compact and semi-compact sets of linear operators in topological vector spaces</i> . . . . .	317
Samuel Ebenstein, <i>Some <math>H^p</math> spaces which are uncomplemented in <math>L^p</math></i> . . . . .	327
David Fremlin, <i>On the completion of locally solid vector lattices</i> . . . . .	341
Herbert Paul Halpern, <i>Essential central spectrum and range for elements of a von Neumann algebra</i> . . . . .	349
G. D. Johnson, <i>Superadditivity intervals and Boas' test</i> . . . . .	381
Norman Lloyd Johnson, <i>Derivation in infinite planes</i> . . . . .	387
V. M. Klassen, <i>The disappearing closed set property</i> . . . . .	403
B. Kuttner and B. N. Sahney, <i>On the absolute matrix summability of Fourier series</i> . . . . .	407
George Maxwell, <i>Algebras of normal matrices</i> . . . . .	421
Kelly Denis McKennon, <i>Multipliers of type <math>(p, p)</math></i> . . . . .	429
James Miller, <i>Sequences of quasi-subordinate functions</i> . . . . .	437
Leonhard Miller, <i>The Hasse-Witt-matrix of special projective varieties</i> . . . . .	443
Michael Cannon Mooney, <i>A theorem on bounded analytic functions</i> . . . . .	457
M. Ann Piech, <i>Differential equations on abstract Wiener space</i> . . . . .	465
Robert Piziak, <i>Sesquilinear forms in infinite dimensions</i> . . . . .	475
Muril Lynn Robertson, <i>The equation <math>y'(t) = F(t, y(g(t)))</math></i> . . . . .	483
Leland Edward Rogers, <i>Continua in which only semi-aposyndetic subcontinua separate</i> . . . . .	493
Linda Preiss Rothschild, <i>Bi-invariant pseudo-local operators on Lie groups</i> . . . . .	503
Raymond Earl Smithson and L. E. Ward, <i>The fixed point property for arcwise connected spaces: a correction</i> . . . . .	511
Linda Ruth Sons, <i>Zeros of sums of series with Hadamard gaps</i> . . . . .	515
Arne Stray, <i>Interpolation sets for uniform algebras</i> . . . . .	525
Alessandro Figà-Talamanca and John Frederick Price, <i>Applications of random Fourier series over compact groups to Fourier multipliers</i> . . . . .	531