

# Pacific Journal of Mathematics

**THE CONVEX CONE OF  $n$ -MONOTONE FUNCTIONS**

ROY MARTIN RAKESTRAW

# THE CONVEX CONE OF $n$ -MONOTONE FUNCTIONS

R. M. RAKESTRAW

A reformulation of the Krein-Milman Theorem is used to obtain an integral representation of each function in a certain class of real monotonic functions defined on  $[0, 1]$ .

Let  $\{i_1, i_2, i_3, \dots\}$  denote a fixed sequence all of whose terms are either 0 or 1, and let  $M_1$  be the set of real non-negative functions  $f$  on  $[0, 1]$  such that

$$(-1)^{(i_1)} \Delta_h^1 f(x) = (-1)^{(i_1)} [f(x+h) - f(x)] \geq 0,$$

$h > 0$ , for  $[x, x+h] \subset [0, 1]$ . Let  $M_n$ ,  $n > 1$ , be the set of functions belonging to  $M_{n-1}$  such that

$$(-1)^{(i_n)} \Delta_h^n f(x) = (-1)^{(i_n)} [\Delta_h^{n-1} f(x+h) - \Delta_h^{n-1} f(x)] \geq 0$$

for  $[x, x+nh] \subset [0, 1]$ . If  $f \in M_n$ , then  $f$  is said to be an  $n$ -monotone function. Since the sum of two  $n$ -monotone functions is in  $M_n$  and since a nonnegative real multiple of an  $n$ -monotone function is an  $n$ -monotone function, the set  $M_n$  is a convex cone. It is the purpose of this paper to give the extremal elements (i.e., the generators of extreme rays) of this cone, and to show that for the  $n$ -monotone functions an integral representation in terms of extremal elements is possible.

A portion of this work appears in the author's Ph. D. dissertation written at Oklahoma State University under the direction of Professor E. K. McLachlan at which time the author was an NDEA Graduate Fellow. The proof of Proposition 3 was suggested by the referee. The author gratefully acknowledges the guidance given by Professor McLachlan and the assistance of the referee's comments.

1. Extremal elements of  $M_n$ . Let  $f$  be a function in  $M_1$  which assumes exactly one positive value in  $[0, 1]$ . If  $f = f_1 + f_2$ , where  $f_1$  and  $f_2 \in M_1$ , then  $f_1$  and  $f_2$  are zero where  $f$  is zero and  $f_1$  and  $f_2$  are constant where  $f$  is constant. Therefore,  $f_1$  and  $f_2$  are proportional to  $f$  and  $f$  is an extremal element of  $M_1$ . On the other hand, if  $f$  assumes at least two positive values in  $[0, 1]$ , then a nonproportional decomposition can be given by taking

$$f_1(x) = \min \{f(x), (1/2) [f(0) + f(1)]\}$$

and  $f_2 = f - f_1$ . Therefore, the extremal elements of  $M_1$  are precisely the functions in  $M_1$  which assume exactly one positive value in  $[0, 1]$ .

Let  $f \in M_n$ ,  $n > 1$ , and let  $\alpha_0 = 0$  if  $i_1 = 0$  and  $\alpha_0 = 1$  if  $i_1 = 1$ . If  $f(\alpha_0) > 0$  and  $f$  is not constant, then take  $f_1 = f(\alpha_0)$  and  $f_2 = f - f_1$ .

In so doing,  $f_1$  and  $f_2 \in M_n$  and  $f_1$  and  $f_2$  are not proportional to  $f$ . Therefore, the only extremal elements  $f$  of  $M_n$  with  $f(a_0) > 0$  are the positive constant functions.

Let  $f \in M_n$ ,  $n > 1$ , and define  $a'_0 = 1 - a_0$ , if  $i_2 = 0$  and  $a'_0 = a_0$  if  $i_2 = 1$ , where  $a_0$  is defined above. It can be shown that if  $f \in M_n$ , then  $f$  must be continuous on  $[0, 1]$  except at  $a'_0$  [9, p. 148]. It follows that the only extremal elements of  $M_1$  that are in  $M_n$  are those which are continuous on  $[0, 1]$  except, possibly, at  $a'_0$ , and these functions are again extremal elements of  $M_n$ .

If  $i_2 = 0$ ,  $f \in M_n$ ,  $n > 1$ ,  $f$  is not constant on  $(0, 1)$  and  $f$  is discontinuous at  $a'_0 = 1 - a_0$ , then take  $f_1(x) = 0$  for  $x \in [0, 1]$  and  $x \neq a'_0$ ,

$$f_1(a'_0) = f(a'_0) - \lim_{x \rightarrow a'_0} f(x) > 0$$

and  $f_2 = f - f_1$ . In so doing,  $f_1$  and  $f_2 \in M_n$  and  $f_1$  and  $f_2$  are not proportional to  $f$ . Hence, whenever  $i_2 = 0$ , the only extremal elements of  $M_n$  that are discontinuous at  $a'_0 = 1 - a_0$  are the functions which are positive at  $a'_0$  and zero elsewhere on  $[0, 1]$ .

On the other hand, if  $i_2 = 1$ ,  $f \in M_n$ ,  $n > 1$ ,  $f$  is not constant on  $(0, 1)$  and  $f$  is discontinuous at  $a'_0 = a_0$ , then let

$$f_1(x) = \lim_{x \rightarrow a'_0} f(x) > 0,$$

$x \in [0, 1]$  and  $x \neq a'_0$ ,  $f_1(a'_0) = 0$  and  $f_2 = f - f_1$ . Then  $f_1$  and  $f_2$  are in  $M_n$  and  $f_1$  and  $f_2$  are not proportional to  $f$ . Therefore, whenever  $i_2 = 1$ , the only extremal elements of  $M_n$  that are discontinuous at  $a'_0 = a_0$  are the functions which are zero at  $a'_0$  and equal to a positive constant elsewhere on  $[0, 1]$ .

Consequently, the extremal elements of  $M_n$ ,  $n > 1$ , which are not extremal elements of  $M_1$  must be zero at  $a_0$  and continuous on  $[0, 1]$ . It will be shown that these extremal elements of  $M_n$  are indefinite integrals of the extremal elements of a cone which is similar to  $M_1$ . This cone is given in Definitions 1 and 2.

**DEFINITION 1.** If  $g$  is a real function monotonic on  $(0, 1)$  and  $n > 1$ , then define the (possibly extended real-valued) function  $I(g, n - 1; \cdot)$  by the equation

$$I(g, n - 1; x) = \int_{a_0}^x \int_{a_1}^{t_1} \cdots \int_{a_{n-3}}^{t_{n-3}} \int_{a_{n-2}}^{t_{n-2}} g(t) dt dt_{n-2} \cdots dt_2 dt_1$$

for  $x \in (0, 1)$ , where  $a_0 = (1/2) [1 - (-1)^{(i_1)}]$  and

$$a_j = (1/2) [1 - (-1)^{(i_j + i_{j+1})}], 1 \leq j \leq n - 2.$$

DEFINITION 2. Let  $K_n$ ,  $n > 1$ , denote the convex cone of real functions  $g$  on  $(0, 1)$  such that

- (a)  $g$  is right-continuous;
- (b)  $(-1)^{(i_{n-1})} g(x) \geq 0$ , for  $x \in (0, 1)$ ;
- (c)  $(-1)^{(i_n)} \Delta_h^1 g(x) \geq 0$ , for  $0 < x < x + h < 1$ ;
- (d)  $I(g, n - 1; x)$  is finite, for  $x \in (0, 1)$ ; and
- (e) limit  $I(g, n - 1; x)$  exists and is finite.  
 $x \rightarrow 1 - a_0$

Note. If  $g \in K_n$ ,  $n > 1$ , then  $I(g, n - 1; \cdot)$  will denote the function which is the continuous extension to  $[0, 1]$  of the function given in Definition 1.

DEFINITION 3. Let  $a$  and  $b$  be two distinct numbers in the interval  $[0, 1]$  and define the function  $\chi_{(a,b)}$  on  $(0, 1)$  by

$$\begin{aligned}\chi_{(a,b)}(x) &= 1, \text{ if } x \text{ is between } a \text{ and } b \text{ or } 0 < x = \min\{a, b\}; \\ \chi_{(a,b)}(x) &= 0, \text{ otherwise.}\end{aligned}$$

DEFINITION 4. If  $m$  is a nonzero real number,  $\xi \in [0, 1]$  and  $n > 1$ , then define the function  $e(m, \xi, n - 1; \cdot)$  by the equation

$$e(m, \xi, n - 1; x) = mI(\chi_{(\xi, 1 - a_{n-1})}, n - 1; x)$$

for  $0 \leq x \leq 1$ , where  $a_{n-1} = (1/2)[1 - (-1)^{(i_{n-1} + i_n)}]$ .

The principal theorem of this section can now be stated and the remainder of the section will be devoted to its proof. The key results are Lemma 3 and Proposition 2.

THEOREM 1. *The extremal elements of  $M_1$  are the functions in  $M_1$  which assume exactly one positive value in  $[0, 1]$ . The positive constant functions and the extremal elements of  $M_1$  which are discontinuous at  $a'_0 = (1/2)[1 + (-1)^{(i_1 + i_2)}]$  are extremal elements of  $M_n$ ,  $n > 1$ . The functions  $e(m, \xi, n - 1; \cdot)$ , where  $(-1)^{(i_{n-1})} m > 0$  and  $\xi \in (0, 1)$  or  $\xi = a_{n-1}$  are extremal elements of  $M_n$ ,  $n > 1$ . There are no other extremal elements of  $M_2$ . The only other extremal elements of  $M_n$ ,  $n > 2$ , are those functions  $e(m, a_k, k; \cdot)$ , where  $(-1)^{(i_k)} m > 0$  and  $1 \leq k \leq n - 2$ .*

In the same manner that the extremal elements of  $M_1$  were found, it can be shown that the extremal elements of  $K_n$  are precisely those functions in  $K_n$  which assume exactly one nonzero value in  $(0, 1)$ . Before determining the extremal elements of  $M_n$ , it is shown in the following three lemmas how the  $n$ -monotone functions are related to the functions in  $K_n$ , where  $n > 1$ .

LEMMA 1. If  $f \in M_n$ , then  $f_+^{(n-1)} \in K_n$ , where  $n > 1$ .

*Proof.* Since  $(-1)^{(i_n)} \Delta_h^n f(x) \geq 0$  for  $0 \leq x < x + nh \leq 1$ , then  $f^{(n-2)}$  exists and is continuous on  $(0, 1)$  and  $(-1)^{(i_n)} f^{(n-2)}$  is convex [1]. Therefore  $(-1)^{(i_n)} f^{(n-2)}$  has a right-continuous, nondecreasing right-hand derivative [4, p. 10]. It follows that  $(-1)^{(i_n)} \Delta_h f_+^{(n-1)}(x) \geq 0$  for  $0 < x + h < 1$ . If  $f \in M_n$ , then  $(-1)^{(i_{n-1})} \Delta_h^{n-1} f(x) \geq 0$  for  $0 \leq x < x + (n-1)h \leq 1$ , which implies that

$$(-1)^{(i_{n-1})} \Delta_{\delta_1}^1 \Delta_{\delta_2}^1 \cdots \Delta_{\delta_{n-1}}^1 f(x) \geq 0$$

for  $0 \leq x < x + \delta_1 + \delta_2 + \cdots + \delta_{n-1} \leq 1$  [1]. It then follows that  $(-1)^{(i_{n-1})} f_+^{(n-1)}(x) \geq 0$  for  $0 < x < 1$ , since  $f_+^{(n-1)}$  exists on  $(0, 1)$ . It remains to show that

$$\lim_{x \rightarrow 1-a_0} I(f_+^{(n-1)}, n-1; x)$$

exists and is finite and this proof will be by induction on  $n$ .

If  $f \in M_2$ , then

$$f(x) = \int_{a_0}^x f'_+(t) dt + \lim_{x \rightarrow a_0} f(x),$$

which implies that

$$\lim_{x \rightarrow 1-a_0} I(f'_+, 1; x) = \lim_{x \rightarrow 1-a_0} f(x) - \lim_{x \rightarrow a_0} f(x)$$

and this latter limit exists and is finite since  $f$  is monotonic on  $[0, 1]$  [4, Theorem 1.1]. Now assume that  $f \in M_n$  implies that

$$\lim_{x \rightarrow 1-a_0} I(f_+^{(n-1)}, n-1; x)$$

exists and is finite and let  $f \in M_{n+1}$ . Then  $f \in M_n$  and it follows from the first part of the proof that  $(-1)^{(i_{n-1})} f^{(n-1)}$  is nonnegative and monotonic on  $(0, 1)$  and

$$\begin{aligned} (-1)^{(i_{n-1})} f^{(n-1)}(a_{n-1}) &= \lim_{x \rightarrow a_{n-1}} (-1)^{(i_{n-1})} f^{(n-1)}(x) \\ &= \inf \{(-1)^{(i_{n-1})} f^{(n-1)}(x) : 0 < x < 1\}. \end{aligned}$$

Therefore,

$$\begin{aligned} &\lim_{x \rightarrow 1-a_0} I(f_+^{(n)}, n; x) \\ &= \lim_{x \rightarrow 1-a_0} I(f^{(n-1)} - f^{(n-1)}(a_{n-1}), n-1; x) \\ &= \lim_{x \rightarrow 1-a_0} I(f^{(n-1)}, n-1; x) - f^{(n-1)}(a_{n-1}) I(1, n-1; x) \end{aligned}$$

exists and is finite by the induction hypothesis.

LEMMA 2. If  $g \in K_n$ , then  $I(g, n-1; \cdot) \in M_n$ , where  $n > 1$ .

*Proof.* The proof will be by induction on  $n$ . If  $g \in K_2$ , then

$$I(g, 1; x) = \int_{a_0}^x g(t) dt$$

for  $x \in [0, 1]$ , and since  $(-1)^{(i_1)}g(t) \geq 0$ ,  $t \in (0, 1)$ , and

$$a_0 = (1/2) [1 - (-1)^{(i_1)}],$$

then  $I(g, 1; x) \geq 0$ . If  $0 \leq x < x+h \leq 1$ , then

$$(-1)^{(i_1)} \Delta_h^1 I(g, 1; x) = \int_x^{x+h} (-1)^{(i_1)} g(t) dt \geq 0.$$

Since  $(-1)^{(i_2)}g$  is nondecreasing, then  $I((-1)^{(i_2)}g, 1; \cdot)$  is convex [4, p. 13]. It follows that  $(-1)^{(i_2)} \Delta_h^2 I(g, 1; x) \geq 0$  for  $0 \leq x < x+2h \leq 1$ , and hence,  $I(g, 1; \cdot) \in M_2$ . Assume that  $I(g, n-1; \cdot) \in M_n$  for  $g \in K_n$  and  $n > 1$ . If  $g \in K_{n+1}$ , then let

$$f(x) = \int_{a_{n-1}}^x g(t) dt,$$

for  $x \in (0, 1)$ . Since  $(-1)^{(i_n)}g$  is nonnegative and

$$a_{n-1} = (1/2) [1 - (-1)^{(i_{n-1}+i_n)}],$$

it is easily seen that  $f \in K_n$  and it follows from the induction hypothesis that  $I(g, n; \cdot) = I(f, n-1; \cdot) \in M_n$ . By a repeated application of the mean value theorem for a Riemann integral, it can be shown that

$$\Delta_h^{n-1} I(g, n; x) = h^{n-1} f(\xi)$$

for  $0 \leq x < \xi < x + (n-1)h \leq 1$ . Since  $(-1)^{(i_{n+1})}g$  is nondecreasing, then  $(-1)^{(i_{n+1})}f$  is convex on  $(0, 1)$  [4, p. 13]. It follows that

$$\begin{aligned} (-1)^{(i_{n+1})} \Delta_h^{n+1} I(g, n; x) &= (-1)^{(i_{n+1})} \Delta_h^2 \Delta_h^{n-1} I(g, n; x) \\ &= (-1)^{(i_{n+1})} \Delta_h^2 f(\xi) \geq 0 \end{aligned}$$

for  $0 \leq x < x + (n+1)h \leq 1$ , and this inequality, together with the fact that  $I(g, n; \cdot) \in M_n$  implies that  $I(g, n; \cdot) \in M_{n+1}$ .

In the proofs that follow,  $f^{(k)}(a_k)$  should be interpreted as

$$f^{(k)}(a_k) = \lim_{x \rightarrow a_k} f^{(k)}(x),$$

where  $f \in M_n$ ,  $n > 2$ , and  $1 \leq k \leq n-2$ . Since  $f^{(k)} \in K_{k+1}$ , this limit will always exist and be finite. It is a consequence of Lemmas 1 and

2 that  $f = I(f_+^{(n-1)}, n-1; \cdot)$  whenever  $f \in M_n$ ,  $n > 1$ , and  $f^{(k)}(a_k) = 0$  for  $0 \leq k \leq n-2$ . It is shown in the following lemma that extremal elements of  $M_n$  can be obtained directly from the extremal elements of  $K_n$ .

**LEMMA 3.** *If  $g \in K_n$  and  $f = I(g, n-1; \cdot)$ , then  $f$  is an extremal element of  $M_n$  if, and only if,  $g$  is an extremal element of  $K_n$ , where  $n > 1$ .*

*Proof.* Suppose that  $f$  is an extremal element of  $M_n$ . If  $g_1$  and  $g_2 \in K_n$  such that  $g = g_1 + g_2$ , then

$$\begin{aligned} f &= I(g, n-1; \cdot) = I(g_1 + g_2, n-1; \cdot) \\ &= I(g_1, n-1; \cdot) + I(g_2, n-1; \cdot). \end{aligned}$$

If  $f_j = I(g_j, n-1; \cdot)$ ,  $j = 1, 2$ , then  $f_1$  and  $f_2 \in M_n$  and  $f = f_1 + f_2$ . Since  $f$  is an extremal element of  $M_n$ , there are numbers  $\lambda_j \geq 0$  such that  $f_j = \lambda_j f$ ,  $j = 1, 2$ , which implies that  $g_j = \lambda_j f_+^{(n-1)} = \lambda_j g$ ,  $j = 1, 2$ , and  $g$  is therefore an extremal element of  $K_n$ .

Conversely, if  $g$  is an extremal element of  $K_n$  and  $f_1$  and  $f_2 \in M_n$  such that  $f = f_1 + f_2$ , then  $g_1$  and  $g_2 \in K_n$  and  $g_1 + g_2 = f_+^{(n-1)} = g$ , where  $g_j$  is the  $(n-1)$ th right derivative of  $f_j$ ,  $j = 1, 2$ . This implies there are constants  $\lambda_j \geq 0$ ,  $j = 1, 2$ , such that  $g_j = \lambda_j g$ . It is evident from the definition of  $f$  that  $f^{(k)}(a_k) = 0$ , where  $0 \leq k \leq n-2$ . This, together with the fact that  $f_j^{(k)} \in K_{k+1}$  for  $1 \leq k \leq n-2$ , implies that  $f_j^{(k)}(a_k) = 0$ ,  $j = 1, 2$  and  $0 \leq k \leq n-2$ .

Hence,

$$f_j = I(g_j, n-1; \cdot) = I(\lambda_j g, n-1; \cdot) = \lambda_j I(g, n-1; \cdot) = \lambda_j f$$

for  $j = 1, 2$ , and  $f$  is therefore an extremal element of  $M_n$ .

**PROPOSITION 1.** *The function  $e(m, \xi, n-1; \cdot)$  is an extremal element of  $M_n$ ,  $n > 1$ , where  $(-1)^{(i_{n-1})} m > 0$  and  $\xi \in (0, 1)$  or  $\xi = a_{n-1}$ .*

*Proof.* Since  $m\chi_{(\xi, 1-a_{n-1})}$  is an extremal element of  $K_n$  whenever  $(-1)^{(i_{n-1})} m > 0$  and  $\xi \in (0, 1)$  or  $\xi = a_{n-1}$ , and

$$e(m, \xi, n-1; \cdot) = I(m\chi_{(\xi, 1-a_{n-1})}, n-1; \cdot),$$

the result follows immediately from Lemma 3.

**PROPOSITION 2.** *The function  $e(m, a_k, k; \cdot)$  is an extremal element of  $M_n$ ,  $n > 2$ , where  $(-1)^{(i_k)} m > 0$  and  $1 \leq k \leq n-2$ .*

*Proof.* Since  $M_n$  is a subcone of  $M_{k+1}$  and  $e(m, a_k, k; \cdot)$  is an extremal element of  $M_{k+1}$ , it is sufficient to show that

$$e(m, a_k, k; \cdot) \in M_n.$$

If  $f = e(m, a_k, k; \cdot)$ , then  $f = I(f^{(k)}, k; \cdot)$ , where

$$f^{(k)}(x) = m\chi_{(a_k, 1-a_k)}(x) = m\chi_{(0,1)}(x) = m$$

for  $0 < x < 1$ . Since  $f^{(k)}$  is constant on  $(0, 1)$ , it follows from a repeated application of the mean value theorem for a Riemann integral that

$$\Delta_h^{k+1} f(x) = \Delta_h^1 \Delta_h^k f(x) = h^k \Delta_h^1 f^{(k)}(\xi) = 0$$

for  $0 \leq x < x + (k+1)h \leq 1$ , where  $x < \xi < x + kh$  and thus,  $\Delta_h^p f(x) = 0$  for  $0 \leq x < x + ph \leq 1$  and  $p \geq k+1$ . Hence,  $f \in M_n$ , for every  $n$ , which implies that  $f$  is an extremal element of  $M_p$ , for  $p \geq k+1$ .

It will follow, as a consequence of the next three lemmas, that no other functions in  $M_n$  are extremal elements of  $M_n$ ,  $n > 2$ .

**LEMMA 4.** *Let  $f \in M_n$ ,  $n > 2$ , such that  $f(a_0) = 0$ ,  $f$  is continuous on  $[0, 1]$  and  $f \neq e(m, a_k, k; \cdot)$  for  $(-1)^{(i_k)} m > 0$  and  $1 \leq k \leq n-2$ . If there is an integer  $k$  such that  $1 \leq k \leq n-2$  and  $f^{(k)}(a_k) \neq 0$ , then  $f$  is not an extremal element of  $M_n$ .*

*Proof.* Let  $k$  denote the smallest integer such that  $f^{(k)}(a_k) \neq 0$ . Then  $f \in M_n \subset M_{k+2}$  implies that  $f_+^{(k+1)} \in K_{k+2}$ , and it follows from Lemma 2 that  $I(f_+^{(k+1)}, k+1; \cdot) \in M_{k+2}$ . Since  $f(a_0) = 0$  and  $f^{(p)}(a_p) = 0$  for  $1 \leq p < k$ , then

$$I(f_+^{(k+1)}, k+1; \cdot) = I(f^{(k)}, k; \cdot) - f^{(k)}(a_k) I(1, k; \cdot) = f - e(m, a_k, k; \cdot)$$

where  $m = f^{(k)}(a_k)$ . Since

$$\Delta_h^p e(m, a_k, k; x) = 0$$

for  $0 \leq x < x + ph \leq 1$  and  $k+1 \leq p \leq n$  and  $f \in M_n$ , it follows that

$$(-1)^{(i_p)} \Delta_h^p I(f_+^{(k+1)}, k+1; x) = (-1)^{(i_p)} \Delta_h^p f(x) \geq 0$$

for  $0 \leq x < x + ph \leq 1$  and  $k+1 \leq p \leq n$ . Hence,

$$f - e(m, a_k, k; \cdot) \in M_n,$$

where  $m = f^{(k)}(a_k)$ , and a nonproportional decomposition of  $f$  can be given by taking  $f_1 = e(m, a_k, k; \cdot)$  and  $f_2 = f - f_1$ . Thus  $f$  is not an extremal element.

**LEMMA 5.** *Let  $f \in M_n$ ,  $n > 2$ , such that  $f \neq 0$ ,  $f(a_0) = 0$ ,  $f$  is*



continuous on  $[0, 1]$  and  $f \neq e(m, a_k, k; \cdot)$  for  $(-1)^{(i_k)} m > 0$  and  $1 \leq k \leq n - 2$ . If  $f_+^{(n-1)} = 0$  on  $(0, 1)$ , then  $f$  is not an extremal element of  $M_n$ .

*Proof.* If  $f_+^{(n-1)} = 0$ , then there is a positive integer  $k \leq n - 2$  such that  $f^{(k)} \neq 0$  and  $f^{(k)}$  is constant on  $(0, 1)$ . Thus,  $f^{(k)}(a_k) \neq 0$  and it follows from Lemma 4 that  $f$  is not an extremal element.

It follows from Lemmas 4 and 5 that if  $f$  is an extremal element of  $M_n$ ,  $n > 2$  such that  $f(a_0) = 0$ ,  $f$  is continuous on  $[0, 1]$  and either  $f_+^{(n-1)} = 0$  or  $f^{(k)}(a_k) \neq 0$  for some  $k$ ,  $1 \leq k \leq n - 2$ , then  $f = e(m, a_k, k; \cdot)$ , where  $(-1)^{(i_k)} m > 0$  and  $1 \leq k \leq n - 2$ .

LEMMA 6. Let  $f \in M_n$ ,  $n \geq 2$ , such that  $f$  is continuous on  $[0, 1]$ ,  $f_+^{(n-1)} \neq 0$  and  $f^{(k)}(a_k) = 0$  for  $0 \leq k \leq n - 2$ . If  $f$  is an extremal element of  $M_n$ , then  $f = e(m, \xi, n - 1; \cdot)$ , where  $(-1)^{(i_{n-1})} m > 0$  and  $\xi \in (0, 1)$  or  $\xi = a_{n-1}$ .

*Proof.* Since  $f^{(k)}(a_k) = 0$  for  $0 \leq k \leq n - 2$ , then

$$f = I(f_+^{(n-1)}, n - 1; \cdot)$$

and it follows from Lemma 3 that  $f_+^{(n-1)}$  is an extremal element of  $K_n$ . Thus,  $f_+^{(n-1)} = m\chi_{(\xi, 1-a_{n-1})}$  for  $(-1)^{(i_{n-1})} m > 0$  and  $\xi \in (0, 1)$  or  $\xi = a_{n-1}$ , which implies that  $f = I(f_+^{(n-1)}, n - 1; \cdot) = e(m, \xi, n - 1; \cdot)$ . This completes the proof of Theorem 1.

2. Integral representations. The set of functions  $M_n - M_n$ ,  $n \geq 1$ , forms the smallest linear space containing the convex cone  $M_n$ . With the topology of simple convergence,  $M_n - M_n$  is a Hausdorff locally convex space such that for each  $x \in [0, 1]$ , the linear functional  $L_x$  defined by  $L_x(f) = f(x)$  is continuous.

PROPOSITION 3. The set  $M_n$  is closed in  $M_n - M_n$  for  $n \geq 1$ .

*Proof.* The linear functional  $F$  defined on  $M_n - M_n$  by  $F(f) = \Delta_h^n f(x)$ , for  $[x, x + nh] \subset [0, 1]$ , is continuous in the topology of simple convergence. By definition,  $M_n$  is the intersection of a collection of closed half-spaces corresponding to such functionals.

Since  $M_n$  is closed and every  $n$ -monotone function  $f$  is nonnegative and bounded by  $f(1 - a_0)$ , Tychonoff's theorem implies that the normalized  $n$ -monotone functions, namely

$$C_n = \{f \in M_n: f(1 - a_0) = 1\},$$

form a compact base for  $M_n$ ,  $n \geq 1$ . Thus, every nonzero  $n$ -monotone function can be uniquely expressed as a positive multiple of some  $f$  in  $C_n$  and  $f$  is an extreme point of the convex set  $C_n$  if, and only if,  $f$  is an extremal element of  $M_n$  which lies in  $C_n$ .

DEFINITION 5. For  $n \geq 2$ , let  $m_\xi$  denote the number which satisfies the equation  $e(m_\xi, \xi, n-1; 1-a_0) = 1$ , where  $\xi \in (0, 1)$  or  $\xi = a_{n-1}$ . For  $n > 2$ , let  $m_k$  denote the constant which satisfies the equation  $e(m_k, a_k, k; 1-a_0) = 1$ , where  $1 \leq k \leq n-2$ . Let  $\text{ext } C_n$  denote the set of extreme points of  $C_n$ ,  $n \geq 1$ , and let  $e(m_0, a_0, 0; \cdot)$  denote the unique function in  $\text{ext } C_n$ ,  $n \geq 2$ , which is discontinuous at  $a'_0 = (1/2)[1 + (-1)^{(i_1+i_2)}]$ ; that is,  $e(m_0, a_0, 0; x) = (1/2)[1 - (-1)^{(i_2)}]$  for  $0 < x < 1$ ,  $e(m_0, a_0, 0; a_0) = 0$  and  $e(m_0, a_0, 0; 1-a_0) = 1$ .

The principal theorem of this section can now be stated and the remainder of the section will be devoted to its proof.

THEOREM 2. To each  $f \in C_n$ ,  $n \geq 2$ , there correspond unique non-negative regular Borel measures  $\nu$  and  $\mu$  on  $[0, 1]$  and

$$\{e(m_k, a_k, k; \cdot): 0 \leq k \leq n-2\},$$

respectively, such that

$$\nu([0, 1]) + f(a_0) + \sum_{\substack{k=0 \\ k \neq k_0}}^{n-2} \mu[e(m_k, a_k, k; \cdot)] = 1$$

and

$$f(x) = \int_0^1 e(m_\xi, \xi, n-1; x) d\nu(\xi) + f(a_0) + \sum_{\substack{k=0 \\ k \neq k_0}}^{n-2} \alpha_k e(m_k, a_k, k; x)$$

for each  $x \in [0, 1]$ , where  $\alpha_k = \mu[e(m_k, a_k, k; \cdot)]$  for each  $k$  and

$$e(m_{1-a_{n-1}}, 1-a_{n-1}, n-1; \cdot) = e(m_{k_0}, a_{k_0}, k_0; \cdot)$$

denotes the function which is the pointwise limit of the functions  $e(m_\xi, \xi, n-1; \cdot)$  as  $\xi$  approaches  $1-a_{n-1}$ . Thus, each  $n$ -monotone function is a scalar multiple of such a representation.

Theorem 2 will be proved by using an integral reformulation of the Krein-Milman theorem. In order to apply this result, it must first be demonstrated that  $\text{ext } C_n$  is closed.

PROPOSITION 4. The set of extreme points of  $C_n$  is closed in  $C_n$ ,  $n \geq 2$ .

*Proof.* Since  $C_n$  with the relative topology is a subspace of a first countable space, it will suffice to show that if  $\{f_i\}$  is a sequence of functions in  $\text{ext } C_n$  which converges pointwise to the function  $f$ , then  $f \in \text{ext } C_n$  [3, p. 164]. Since all except a finite number of the functions in  $\text{ext } C_n$  are of the form  $e(m_{\xi}, \xi, n-1; \cdot)$ , where  $\xi \in (0, 1)$  or  $\xi = a_{n-1}$ , it can be assumed without loss of generality that  $f_i = e(m_{\xi_i}, \xi_i, n-1; \cdot)$  for each  $i$ .

If  $a_0 = a_1 = \dots = a_{n-1}$ , then the function in  $C_n$  are convex and

$$f_i(x) = \left( \frac{x - \xi_i}{1 - a_0 - \xi_i} \right)^{n-1} \chi_{(\xi_i, 1-a_0)}(x)$$

for  $x \in (0, 1)$ . If the sequence  $\{\xi_i\}$  of real numbers converges to  $1 - a_0$ , then it is easily seen that

$$\lim_{i \rightarrow \infty} f_i(x) = 0$$

for  $x \in (0, 1)$  or  $x = a_0$ . Since the topology of simple convergence is a Hausdorff topology, it follows that  $f(1 - a_0) = 1$  and  $f(x) = 0$ , otherwise, which implies that  $f = e(m_0, a_0, 0; \cdot)$  and  $f \in \text{ext } C_n$ . On the other hand, if  $\{\xi_i\}$  does not converge to  $1 - a_0$ , then there is a real number  $\xi_0 \neq 1 - a_0$  and a subsequence  $\{\xi_j\}$  of  $\{\xi_i\}$  such that  $\{\xi_j\}$  converges to  $\xi_0$ . Hence,

$$\begin{aligned} \lim_{j \rightarrow \infty} f_j(x) &= \lim_{j \rightarrow \infty} \left( \frac{x - \xi_j}{1 - a_0 - \xi_j} \right)^{n-1} \chi_{(\xi_j, 1-a_0)}(x) \\ &= \left( \frac{x - \xi_0}{1 - a_0 - \xi_0} \right)^{n-1} \chi_{(\xi_0, 1-a_0)}(x) \\ &= e(m_{\xi_0}, \xi_0, n-1; x) \end{aligned}$$

for each  $x \in (0, 1)$ . Therefore, since the topology is a Hausdorff topology,  $f = e(m_{\xi_0}, \xi_0, n-1; \cdot)$  and it follows that  $f \in \text{ext } C_n$ .

If  $a_1 = a_2 = \dots = a_{n-1}$  and  $a_0 \neq a_{n-1}$ , then the functions in  $C_n$  are concave and

$$f_i(x) = 1 - \left( \frac{x - \xi_i}{a_0 - \xi_i} \right)^{n-1} \chi_{(\xi_i, a_0)}(x)$$

for  $x \in (0, 1)$ . If the sequence  $\{\xi_i\}$  converges to  $a_0$ , then

$$\lim_{i \rightarrow \infty} f_i(x) = 1$$

for  $x \in (0, 1)$  or  $x = 1 - a_0$  and  $f = e(m_0, a_0, 0; \cdot)$ . On the other hand, if there is a subsequence  $\{\xi_j\}$  of  $\{\xi_i\}$  which converges to  $\xi_0 \neq a_0$ , then

$$\begin{aligned}\lim_{j \rightarrow \infty} f_j(x) &= \lim_{j \rightarrow \infty} \left[ 1 - \left( \frac{x - \xi_j}{a_0 - \xi_j} \right)^{n-1} \chi_{(\xi_j, a_0)}(x) \right] \\ &= 1 - \left( \frac{x - \xi_0}{a_0 - \xi_0} \right)^{n-1} \chi_{(\xi_0, a_0)}(x) = e(m_{\xi_0}, \xi_0, n-1; x)\end{aligned}$$

for each  $x \in (0, 1)$  and  $f = e(m_{\xi_0}, \xi_0, n-1; \cdot)$ . In either case, it follows that  $f \in \text{ext } C_n$ .

If there are exactly  $p > 0$  integers  $k_1, \dots, k_p$  such that

$$1 \leq k_1 < k_2 < \dots < k_p \leq n-2$$

and  $a_{k_j} \neq a_{n-1}$ ,  $1 \leq j \leq p$ , and  $a_0 = a_{n-1}$ , then

$$\begin{aligned}f_i(x) &= m_{\xi_i} \left[ \frac{(x - \xi_i)^{n-1}}{(n-1)!} \chi_{(\xi_i, 1-a_0)}(x) \right. \\ &\quad \left. + \sum_{r=1}^p (-1)^r \sum_{j_r=r}^p \dots \sum_{j_1=1}^{j_2-1} \frac{(1 - a_0 - \xi_i)^{n-k_{j_r}-1} (1 - 2a_0)^{k_{j_r}-k_{j_1}} (x - a_0)^{k_{j_1}}}{(n-k_{j_1}-1)! (k_{j_r}-k_{j_{r-1}})! \dots (k_{j_2}-k_{j_1})! (k_{j_1})!} \right]\end{aligned}$$

for  $x \in (0, 1)$ , where

$$\begin{aligned}m_{\xi_i}^{-1} &= \frac{(1 - a_0 - \xi_i)^{n-1}}{(n-1)!} \\ &\quad + \sum_{r=1}^p (-1)^r \sum_{j_r=r}^p \dots \sum_{j_1=1}^{j_2-1} \frac{(1 - a_0 - \xi_i)^{n-k_{j_r}-1} (1 - 2a_0)^{k_{j_r}-k_{j_1}}}{(n-k_{j_r}-1)! (k_{j_r}-k_{j_{r-1}})! \dots (k_{j_2}-k_{j_1})! (k_{j_1})!}.\end{aligned}$$

If there is a subsequence  $\{\xi_j\}$  of  $\{\xi_i\}$  which converges to  $\xi_0 \neq 1 - a_0$ , then it is easily seen that

$$f(x) = \lim_{j \rightarrow \infty} f_j(x) = e(m_{\xi_0}, \xi_0, n-1; x)$$

for each  $x \in (0, 1)$ . On the other hand, if  $\{\xi_i\}$  converges to  $1 - a_0$ , then

$$\begin{aligned}\lim_{i \rightarrow \infty} f_i(x) &= m_{k_p} \left[ \frac{(x - a_0)^{(k_p)}}{(k_p)!} \right. \\ &\quad \left. + \sum_{r=1}^{p-1} (-1)^r \sum_{j_r=r}^{p-1} \dots \sum_{j_1=1}^{j_2-1} \frac{(1 - 2a_0)^{k_p-k_{j_1}} (x - a_0)^{k_{j_1}}}{(k_p-k_{j_r})! (k_{j_r}-k_{j_{r-1}})! \dots (k_{j_2}-k_{j_1})! (k_{j_1})!} \right] \\ &= e(m_{k_p}, a_{k_p}, k_p; x)\end{aligned}$$

for  $x \in (0, 1)$ , where

$$\begin{aligned}m_{k_p}^{-1} &= \frac{(1 - 2a_0)^{(k_p)}}{(k_p)!} \\ &\quad + \sum_{r=1}^{p-1} (-1)^r \sum_{j_r=r}^{p-1} \dots \sum_{j_1=1}^{j_2-1} \frac{(1 - 2a_0)^{(k_p)}}{(k_p-k_{j_r})! (k_{j_r}-k_{j_{r-1}})! \dots (k_{j_2}-k_{j_1})! (k_{j_1})!}.\end{aligned}$$

In either case, it follows that  $f \in \text{ext } C_n$ .

Finally if there are exactly  $p > 0$  integers  $k_1, \dots, k_p$  such that  $1 \leq k_1 < k_2 < \dots < k_p \leq n - 2$  and  $a_{k_j} \neq a_{n-1}$ ,  $1 \leq j \leq p$  and  $a_0 \neq a_{n-1}$ , then

$$\begin{aligned} & f_i(x) \\ &= m_{\xi_i} \left[ \frac{(a_0 - \xi_i)^{n-1}}{(n-1)!} - \frac{(x - \xi_i)^{n-1}}{(n-1)!} \chi_{(\xi_i, a_0)}(x) \right. \\ & \quad + \sum_{r=1}^p (-1)^r \sum_{j_r=r}^p \dots \sum_{j_1=1}^{j_2-1} \frac{(a_0 - \xi_i)^{n-k_{j_r}-1} (2a_0 - 1)^{k_{j_r}}}{(n-k_{j_r}-1)! (k_{j_r}-k_{j_{r-1}})! \dots (k_{j_2}-k_{j_1})! (k_{j_1})!} \\ & \quad \left. - \sum_{r=1}^p (-1)^r \sum_{j_r=r}^p \dots \sum_{j_1=1}^{j_2-1} \frac{(a_0 - \xi_i)^{n-k_{j_r}-1} (2a_0 - 1)^{k_{j_r}-k_{j_1}} (x-1+a_0)^{k_{j_1}}}{(n-k_{j_r}-1)! (k_{j_r}-k_{j_{r-1}})! \dots (k_{j_2}-k_{j_1})! (k_{j_1})!} \right] \end{aligned}$$

for  $x \in (0, 1)$ , where

$$\begin{aligned} & m_{\xi_i}^{-1} \\ &= \frac{(a_0 - \xi_i)^{n-1}}{(n-1)!} \\ & \quad + \sum_{r=1}^p (-1)^r \sum_{j_r=r}^p \dots \sum_{j_1=1}^{j_2-1} \frac{(a_0 - \xi_i)^{n-k_{j_r}-1} (2a_0 - 1)^{k_{j_r}}}{(n-k_{j_r}-1)! (k_{j_r}-k_{j_{r-1}})! \dots (k_{j_2}-k_{j_1})! (k_{j_1})!} . \end{aligned}$$

If there is a subsequence  $\{\xi_{j_i}\}$  of  $\{\xi_i\}$  which converges to  $\xi_0 \neq a_0$ , then it is evident that

$$f(x) = \lim_{j \rightarrow \infty} f_j(x) = e(m_{\xi_0}, \xi_0, n-1; x)$$

for each  $x \in (0, 1)$ . On the other hand, if  $\{\xi_i\}$  converges to  $a_0$ , then

$$\begin{aligned} & \lim_{i \rightarrow \infty} f_i(x) \\ &= m_{k_p} \left[ \frac{(2a_0 - 1)^{(k_p)}}{(k_p)!} - \frac{(x - 1 + a_0)^{(k_p)}}{(k_p)!} \right. \\ & \quad + \sum_{r=1}^{p-1} (-1)^r \sum_{j_r=r}^{p-1} \dots \sum_{j_1=1}^{j_2-1} \frac{(2a_0 - 1)^{k_p-k_{j_1}} [(2a_0 - 1)^{k_{j_1}} - (x-1+a_0)^{k_{j_1}}]}{(k_p-k_{j_r})! (k_{j_r}-k_{j_{r-1}})! \dots (k_{j_2}-k_{j_1})! (k_{j_1})!} \left. \right] \\ &= e(m_{k_p}, a_{k_p}, k_p; x) \end{aligned}$$

for  $x \in (0, 1)$ , where

$$\begin{aligned} & m_{k_p}^{-1} \\ &= \frac{(2a_0 - 1)^{(k_p)}}{(k_p)!} \\ & \quad + \sum_{r=1}^{p-1} (-1)^r \sum_{j_r=r}^{p-1} \dots \sum_{j_1=1}^{j_2-1} \frac{(2a_0 - 1)^{(k_p)}}{(k_p-k_{j_r})! (k_{j_r}-k_{j_{r-1}})! \dots (k_{j_2}-k_{j_1})! (k_{j_1})!} . \end{aligned}$$

In either case it follows that  $f \in \text{ext } C_n$  and this completes the proof.

DEFINITION 6. Let  $e_0$  denote the function in  $\text{ext } C_n$  which is identically one and let  $e(m_{1-a_{n-1}}, 1-a_{n-1}, n-1; \cdot)$  be the function defined by

$$e(m_{1-a_{n-1}}, 1-a_{n-1}, n-1; x) = \lim_{\xi \rightarrow 1-a_{n-1}} e(m_\xi, \xi, n-1; x)$$

for  $0 \leq x \leq 1$  and  $n > 1$ . Finally, let

$$e(m_{k_0}, a_{k_0}, k_0; \cdot) = e(m_{1-a_{n-1}}, 1-a_{n-1}, n-1; \cdot)$$

and notice that  $k_0 = 0$  if  $a_1 = a_2 = \dots = a_{n-1}$  or  $k_0$  is the largest positive integer such that  $a_{k_0} \neq a_{n-1}$ .

If the mapping  $\phi: [0, 1] \rightarrow \text{ext } C_n$ ,  $n \geq 2$ , is defined by

$$\phi(\xi) = e(m_\xi, \xi, n-1; \cdot) \quad \text{for } 0 \leq \xi \leq 1,$$

then it follows from the proof of Proposition 4 that  $\phi$  is continuous. If  $E = \phi([0, 1])$ , then  $\phi$  is a homeomorphism from  $[0, 1]$  onto  $E$ , since  $[0, 1]$  is a compact space and  $E$  is a Hausdorff space. By the Krein-Milman representation theorem, to each  $f$  in  $C_n$  there corresponds a regular Borel probability measure  $\mu$  on  $\text{ext } C_n$  such that

$$L(f) = \int_{\text{ext } C_n} L d\mu$$

for each continuous linear functional  $L$  on  $M_n - M_n$ , since both  $C_n$  and  $\text{ext } C_n$  are compact subsets of  $M_n - M_n$ ,  $n \geq 2$ . For  $0 \leq x \leq 1$ , the evaluation functional  $L_x$  defined by  $L_x(f) = f(x)$  is continuous on  $M_n - M_n$ , so that

$$\begin{aligned} (1) \quad f(x) &= \int_{\text{ext } C_n} L_x d\mu \\ &= \int_E L_x d\mu + \mu(e_0) + \sum_{\substack{k=0 \\ k \neq k_0}}^{n-2} e(m_k, a_k, k; x) \mu[e(m_k, a_k, k; \cdot)] \end{aligned}$$

for each  $x \in [0, 1]$ . Define  $\nu$  on each Borel subset  $B$  of  $[0, 1]$  by

$$\nu(B) = \mu[\phi(B)]; \text{ i.e., } \nu = \mu\phi.$$

Since  $L_x[\phi(\xi)] = e(m_\xi, \xi, n-1; x)$ , then

$$\int_E L_x d\mu = \int_{\phi^{-1}(E)} L_x \phi d(\mu\phi) = \int_0^1 e(m_\xi, \xi, n-1; x) d\nu(\xi)$$

for  $0 \leq x \leq 1$ . Finally, by observing that  $\mu(e_0) = f(a_0)$ , since  $e_0$  is the only function in  $\text{ext } C_n$  which is positive at  $a_0$ , Equation (1) can be written as

$$f(x) = \int_0^1 e(m_\xi, \xi, n-1; x) d\nu(\xi) \\ + f(a_0) + \sum_{\substack{k=0 \\ k \neq k_0}}^{n-2} e(m_k, a_k, k; x) \mu[e(m_k, a_k, k; \cdot)] .$$

It remains to prove that  $\mu$  is unique. Since  $\mu$  is supported by  $\text{ext } C_n$ , then  $\mu$  is a maximal measure in Choquet's ordering [6, pp. 24, 70]. Thus, by the Choquet-Meyer uniqueness theorem, it suffices to prove that  $C_n$  is a simplex [6, p. 66].

LEMMA 7. Suppose  $f \in M_n - M_n$  and  $n \geq 2$ . Then there is a function  $g \in K_n$  such that  $g - f_+^{(n-1)} \in K_n$  and if  $h$  is any function in  $K_n$  such that  $h - f_+^{(n-1)} \in K_n$ , then it must follow that  $h - g \in K_n$ .

*Proof.* First assume that  $i_{n-1} = i_n = 0$ . Since  $f_+^{(n-1)} \in K_n - K_n$ , then  $f_+^{(n-1)}$  is of bounded variation on every interval  $[0, x]$ , where  $0 < x < 1$ . Define  $g(x) = f_+^{(n-1)}(0) + P_0^x(f_+^{(n-1)})$ , where  $P_0^x(f_+^{(n-1)})$  denotes the positive variation of  $f_+^{(n-1)}$  over  $[0, x]$ ,  $0 \leq x < 1$  [8, p. 85]. Then both  $g$  and  $g - f_+^{(n-1)}$  are nonnegative, nondecreasing and right-continuous on  $[0, 1)$ . If  $h \in K_n$  such that  $h - f_+^{(n-1)} \in K_n$ , then it follows that  $h - g$  is nonnegative, nondecreasing and right-continuous on  $[0, 1)$ . Therefore,

$$0 \leq \lim_{x \rightarrow 1-a_0} I(h - g, n-1; x) \leq \lim_{x \rightarrow 1-a_0} I(h, n-1; x) ,$$

which implies that both  $g$  and  $h - g$  are in  $K_n$ .

If  $i_{n-1}$  and  $i_n$  are not both zero, then define

$$y = (1/2) [1 - (-1)^{(i_{n-1} + i_n)} (1 - 2x)]$$

and

$$F(x) = (-1)^{(i_{n-1})} f_+^{(n-1)}(y) \quad \text{for } 0 \leq x < 1 .$$

Let  $G(x) = F(0) + P_0^x(F)$  for  $0 \leq x < 1$  and define  $g(x) = (-1)^{(i_{n-1})} G(y)$ . Then  $g$  and  $g - f_+^{(n-1)} \in K_n$  and it follows from the first part of the proof that if  $h$  and  $h - f_+^{(n-1)} \in K_n$ , then  $h - g \in K_n$ .

DEFINITION 7. If  $u$  is a function in  $M_n - M_n$ ,  $n \geq 2$ , then define the functions  $u_k$ ,  $0 \leq k \leq n-2$ , by

$$u_0(x) = u(a_0) \quad \text{and} \\ u_k(x) = I(u^{(k)}(a_k), k; x) \quad \text{for } 1 \leq k \leq n-2$$

where  $x \in [0, 1]$ .

LEMMA 8. Suppose  $f \in M_n - M_n$  and  $n \geq 2$ . Then there is a

function  $g \in M_n$  such that  $g - f \in M_n$  and if  $h$  is any  $n$ -monotone function such that  $h - f \in M_n$ , then it must follow that  $h - g \in M_n$ .

*Proof.* First assume that  $f^{(k)}(a_k) = 0$  for  $0 \leq k \leq n - 2$  and let  $g_+^{(n-1)}$  denote the function in  $K_n$  guaranteed by Lemma 7. Define  $g = I(g_+^{(n-1)}, n - 1; \cdot)$ ; then  $g \in M_n$  and

$$g - f = I(g_+^{(n-1)} - f_+^{(n-1)}, n - 1; \cdot) \in M_n.$$

If  $h$  is an  $n$ -monotone function such that  $h - f \in M_n$ , then  $h_+^{(n-1)}$  and  $h_+^{(n-1)} - f_+^{(n-1)} \in K_n$  and it follows that  $h_+^{(n-1)} - g_+^{(n-1)} \in K_n$ . If  $h^{(k)}(a_k) = 0$  for  $0 \leq k \leq n - 2$ , then

$$h - g = I(h_+^{(n-1)} - g_+^{(n-1)}, n - 1; \cdot) \in M_n.$$

If there is some integer  $p$  such that  $0 \leq p \leq n - 2$  and  $h^{(p)}(a_p) \neq 0$ , then let

$$\bar{h} = h - \sum_{k=0}^{n-2} h_k,$$

where  $h_0 = h(a_0)$  and  $h_k = I(h^{(k)}(a_k), k; \cdot)$  for  $1 \leq k \leq n - 2$ . Then  $\bar{h}^{(k)}(a_k) = 0$  for  $0 \leq k \leq n - 2$  and  $\bar{h}$  and  $\bar{h} - f \in M_n$ , since  $h$  and  $h - f \in M_n$  (cf. proof of Lemma 4). It follows that  $\bar{h} - g \in M_n$  which implies that

$$h - g = \bar{h} - g + \sum_{k=0}^{n-2} h_k \in M_n$$

since  $h_k$  is an  $n$ -monotone function for  $0 \leq k \leq n - 2$ .

On the other hand, if there is a nonnegative integer  $p \leq n - 2$  such that  $f^{(p)}(a_p) \neq 0$ , then let

$$\bar{f} = f - \sum_{k=0}^{n-2} f_k$$

where  $f_k$  is given by Definition 7. Since  $\bar{f} \in M_n - M_n$  and  $\bar{f}^{(k)}(a_k) = 0$  for  $0 \leq k \leq n - 2$ , it follows from the first part of the proof that there is an  $n$ -monotone function  $\bar{g}$  such that  $\bar{g} - \bar{f} \in M_n$  and if  $h$  is an  $n$ -monotone function such that  $h - f \in M_n$ , then  $h - \bar{g} \in M_n$ . Let  $k_j, 0 \leq j \leq p < n - 1$ , denote those integers for which

$$(-1)^{i_{k_j}} f^{(k_j)}(a_{k_j}) > 0$$

and define

$$g = \bar{g} + \sum_{j=0}^p f_{k_j}.$$

Then  $g \in M_n$  since



$$f_{k_j} = I(f^{(k_j)}(a_{k_j}), k_j; \cdot) = e(f^{(k_j)}(a_{k_j}), a_{k_j}, k_j; \cdot) \in M_n$$

for  $0 \leq j \leq p$ , and

$$g - f = \bar{g} + \sum_{j=0}^p f_{k_j} - f = \bar{g} - \bar{f} - \sum_{k \neq k_j} f_k \in M_n$$

since  $-f_k \in M_n$  if  $k \neq k_j$ . Suppose that  $h$  is an  $n$ -monotone function such that  $h - f \in M_n$ . Then

$$h - f - \sum_{k=0}^{n-2} (h - f)_k \in M_n$$

which implies that

$$h - f - \sum_{k \neq k_j} (h - f)_k = h - f - \sum_{k=0}^{n-2} (h - f)_k + \sum_{j=0}^p (h - f)_{k_j} \in M_n$$

since  $(h - f)_{k_j} \in M_n$  (cf. proof of Lemma 4). Since  $h_k$  is an  $n$ -monotone function for  $0 \leq k \leq n - 2$ , then

$$\begin{aligned} h - f + \sum_{k \neq k_j} f_k &= h - f - \sum_{k \neq k_j} (h_k - f_k) + \sum_{k \neq k_j} h_k \\ &= h - f - \sum_{k \neq k_j} (h - f)_k + \sum_{k \neq k_j} h_k \in M_n. \end{aligned}$$

Therefore,

$$h - \sum_{j=0}^p f_{k_j} - \bar{f} = h - f + \sum_{k \neq k_j} f_k \in M_n$$

and  $h - \sum_{j=0}^p f_{k_j} \in M_n$  since  $h - \sum_{j=0}^p h_{k_j} \in M_n$  and

$$h - \sum_{j=0}^p f_{k_j} = h - \sum_{j=0}^p h_{k_j} + \sum_{j=0}^p (h_{k_j} - f_{k_j}) = h - \sum_{j=0}^p h_{k_j} + \sum_{j=0}^p (h - f)_{k_j}.$$

It follows that  $h - \sum_{j=0}^p f_{k_j} - \bar{g} \in M_n$ , which implies that  $h - g \in M_n$ .

If the function  $g$  of Lemma 8 is denoted by  $f \vee 0$ , then the least upper bound of two functions  $f_1$  and  $f_2 \in M_n - M_n$  can be given by  $f_1 + (f_2 - f_1) \vee 0$  and therefore  $M_n - M_n$  is a vector lattice. Thus,  $C_n$  is a simplex and the proof of Theorem 2 is complete.

**3. REMARKS.** If  $i_2 = 0$ , then  $C_2$  is the set of functions  $f$  which are monotonic and convex on  $[0, 1]$  such that  $\max \{f(x) : 0 \leq x \leq 1\} = 1$ . If  $i_1 = 0$ , then the  $C_2$  functions are nondecreasing and  $e(m_\xi, \xi, 1; x) = 0$ ,  $x \in [0, \xi]$  and  $(x - \xi)/(1 - \xi)$  for  $x \in [\xi, 1]$ , where  $0 \leq \xi < 1$ . Thus, to each  $f \in C_2$  there corresponds a unique nonnegative regular Borel measure  $\nu$  on  $[0, 1]$  such that

$$f(x) = f(0) + \int_0^x \frac{x - \xi}{1 - \xi} d\nu(\xi)$$

for  $0 < x < 1$ . On the other hand, if  $i_1 = 1$ , then these functions are nonincreasing and  $e(m_\xi, \xi, 1; x) = 1 - (x/\xi)$ ,  $x \in [0, \xi]$  and 0 for  $x \in [\xi, 1]$ , where  $0 < \xi \leq 1$ . It follows from Theorem 2 that to each  $f$  in  $C_2$  there corresponds a unique nonnegative regular Borel measure  $\nu$  on  $[0, 1]$  such that

$$f(x) = f(1) + \int_x^1 [1 - (x/\xi)] d\nu(\xi)$$

for  $0 < x < 1$ .

If  $i_k = 0$  for every  $k \leq n$ , then  $e(m_\xi, \xi, n-1; x) = 0$ ,  $x \in [0, \xi]$  and  $[(x-\xi)/(1-\xi)]^{n-1}$  for  $x \in [\xi, 1]$ , where  $0 \leq \xi < 1$ , and

$$e(m_k, 0, k; x) = x^k$$

for  $x \in [0, 1]$ , where  $1 \leq k \leq n-2$ . Thus, for each function  $f$  in  $C_n$ , there exist unique nonnegative real numbers  $\alpha_1, \dots, \alpha_{n-2}$  and a unique nonnegative regular Borel measure  $\nu$  on  $[0, 1]$  such that

$$f(x) = f(0) + \sum_{k=1}^{n-2} \alpha_k x^k + \int_0^x \left( \frac{x-\xi}{1-\xi} \right)^{n-1} d\nu(\xi)$$

for  $0 < x < 1$ . In this case, the intersection of the  $M_n$  cones is the class of absolutely monotonic functions on  $[0, 1]$ . It is well known that if  $f \in C_n$  for every  $n$ , then

$$f(x) = \sum_{n=0}^{\infty} f^{(n)}(0) (x^n/n!)$$

for  $0 \leq x < 1$ . For a discussion of these cones see [5].

Lastly, if  $i_k = (1/2)[1 + (-1)^k]$  for  $1 \leq k \leq n$ , then

$$e(m_\xi, \xi, n-1; x) = 1 - [1 - (x/\xi)]^{n-1},$$

$x \in [0, \xi]$  and 1 for  $x \in [\xi, 1]$ , where  $0 < \xi \leq 1$ , and

$$e(m_k, 1, k; x) = 1 - (1-x)^k$$

for  $x \in [0, 1]$ , where  $1 \leq k \leq n-2$ . It follows from Theorem 2 that for each function  $f$  in  $C_n$ , there exist unique nonnegative real numbers  $\alpha_1, \dots, \alpha_{n-2}$  and a unique nonnegative regular Borel measure  $\nu$  on  $[0, 1]$  such that

$$f(x) = 1 - \sum_{k=1}^{n-2} \alpha_k (1-x)^k - \int_x^1 [1 - (x/\xi)]^{n-1} d\nu(\xi)$$

for  $0 < x < 1$ . In this case, the  $C_n$  functions were called alternating of order  $n$  by Choquet [2, p. 170]. It can be shown that if  $f \in C_n$  for every  $n$ , then

$$f(x) = \sum_{n=0}^{\infty} f^{(n)}(1) [(x-1)^n/n!]$$

for  $0 < x \leq 1$ . For a proof of this fact together with a discussion of these cones see [7].

#### REFERENCES

1. R. P. Boas, Jr. and D. V. Widder, *Functions with positive differences*, Duke Math. J. **7** (1940), 496-503.
2. G. Choquet, *Theory of capacities*, Annales de l'Institut Fourier, **5** (1953 and 1954), 131-296.
3. J. L. Kelley, *General Topology*, D. Van Nostrand, Princeton, New Jersey, 1955.
4. M. A. Krasnosel'skii and Ya. B. Rutickii, *Convex Functions and Orlicz Spaces* (Translation, AEC-tr-4311), State Publishing House of Physico-Mathematical Literature, Moscow, 1958.
5. E. K. McLachlan, *Extremal elements of the convex cone  $B_n$  of functions*, Pacific J. Math. **14** (1964), 987-993.
6. R. R. Phelps, *Lectures on Choquet's Theorem*, Van Nostrand Mathematical Studies **7** (1966).
7. R. M. Rakestraw, *Extremal elements of the convex cone  $A_n$  of functions*, Pacific J. Math., **34** (1970), 489-498.
8. H. L. Royden, *Real Analysis*, Macmillan, New York, N. Y., 1963.
9. D. V. Widder, *The Laplace Transform*, Princeton Mathematical Series 6 (1946).

Received March 15, 1971.

UNIVERSITY OF MISSOURI

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON

Stanford University  
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

C. R. HOBBY

University of Washington  
Seattle, Washington 98105

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

# Pacific Journal of Mathematics

Vol. 43, No. 3

May, 1972

Max K. Agoston, <i>An obstruction to finding a fixed point free map on a manifold</i> . . . .	543
Nadim A. Assad and William A. Kirk, <i>Fixed point theorems for set-valued mappings of contractive type</i> . . . . .	553
John Winston Bunce, <i>Characterizations of amenable and strongly amenable <math>C^*</math>-algebras</i> . . . . .	563
Erik Maurice Ellentuck and Alfred Berry Manaster, <i>The decidability of a class of AE sentence in the isols</i> . . . . .	573
U. Haussmann, <i>The inversion theorem and Plancherel's theorem in a Banach space</i> . . . . .	585
Peter Lawrence Falb and U. Haussmann, <i>Bochner's theorem in infinite dimensions</i> . . . . .	601
Peter Fletcher and William Lindgren, <i>Quasi-uniformities with a transitive base</i> . . . .	619
Dennis Garbanati and Robert Charles Thompson, <i>Classes of unimodular abelian group matrices</i> . . . . .	633
Kenneth Hardy and R. Grant Woods, <i>On <math>c</math>-realcompact spaces and locally bounded normal functions</i> . . . . .	647
Manfred Knebusch, Alex I. Rosenberg and Roger P. Ware, <i>Grothendieck and Witt rings of hermitian forms over Dedekind rings</i> . . . . .	657
George M. Lewis, <i>Cut loci of points at infinity</i> . . . . .	675
Jerome Irving Malitz and William Nelson Reinhardt, <i>A complete countable <math>L_{\omega_1}^Q</math> theory with maximal models of many cardinalities</i> . . . . .	691
Wilfred Dennis Pepe and William P. Ziemer, <i>Slices, multiplicity, and Lebesgue area</i> . . . . .	701
Keith Pierce, <i>Amalgamating abelian ordered groups</i> . . . . .	711
Stephen James Pride, <i>Residual properties of free groups</i> . . . . .	725
Roy Martin Rakestraw, <i>The convex cone of <math>n</math>-monotone functions</i> . . . . .	735
T. Schwartzbauer, <i>Entropy and approximation of measure preserving transformations</i> . . . . .	753
Peter F. Stebe, <i>Invariant functions of an iterative process for maximization of a polynomial</i> . . . . .	765
Kondagunta Sundaresan and Wojbor Woyczynski, <i><math>L</math>-orthogonally scattered measures</i> . . . . .	785
Kyle David Wallace, <i><math>C_\lambda</math>-groups and <math>\lambda</math>-basic subgroups</i> . . . . .	799
Barnet Mordecai Weinstock, <i>Approximation by holomorphic functions on certain product sets in <math>C^n</math></i> . . . . .	811
Donald Steven Passman, <i>Corrections to: "Isomorphic groups and group rings"</i> . . . .	823
Don David Porter, <i>Correction to: "Symplectic bordism, Stiefel-Whitney numbers, and a Novikov resolution"</i> . . . . .	825
John Ben Butler, Jr., <i>Correction to: "Almost smooth perturbations of self-adjoint operators"</i> . . . . .	825
Constantine G. Lascarides, <i>Correction to: "A study of certain sequence spaces of Maddox and a generalization of a theorem of Iyer"</i> . . . . .	826
George A. Elliott, <i>Correction to: "An extension of some results of Takesaki in the reduction theory of von Neumann algebras"</i> . . . . .	826
James Daniel Halpern, <i>Correction to: "On a question of Tarski and a maximal theorem of Kurepa"</i> . . . . .	827