

# Pacific Journal of Mathematics

## **THE INDEX OF CONVEXITY AND THE VISIBILITY FUNCTION**

GERALD A. BEER

# THE INDEX OF CONVEXITY AND THE VISIBILITY FUNCTION

GERALD BEER

If the integral of the visibility function for a set  $E$  is normalized, one arrives at the Index of convexity of  $E$ , a measure of the relative convexity of  $E$  in terms of the average "area seen" by a variable point of  $E$ . As the visibility function is upper semicontinuous on a compact set in  $E_n$ , the Index is upper semicontinuous on the class of all compact sets in  $E_n$  with an appropriate metric. We also investigate natural generalizations of convex and starshaped sets in terms of the visibility function.

DEFINITION. The visibility function assigns to each point  $x$  of a fixed measurable set  $E$  in a Euclidean space  $E_n$  the Lebesgue outer measure of  $S(x)$ , the set  $\{y: rx + (1 - r)y \in E \text{ for each } r \text{ in } [0, 1]\}$  and zero to each point of  $E_n/E$ .

The study of the visibility function is the study of the dynamic properties of the "star" of a variable point  $x$  of  $E$ ; in effect, the techniques used may be described as "starshaped analysis."

If  $E$  has positive finite Lebesgue measure, a natural representation of the relative convexity of  $E$  is the following Index of convexity:

$$I(E) = \int_E \frac{m(S(x))}{m(E)^2} dm(x)$$

providing this expression makes sense. We show that the Index is upper semicontinuous on the family of all compact sets in  $E_n$  with an appropriate metric. Finally, we investigate the natural generalizations of convex and starshaped sets in terms of the visibility function and establish desired decomposition theorems for these objects in the compact case.

2. Preliminaries. We essentially use the same terminology as in [3]. We denote ordinary Lebesgue measure in  $E_n$  by either  $m$  or  $m_n$  (if more than one measure is under discussion).  $B_r(x)$  will denote the closed  $r$ -ball about a point  $x$ ;  $\text{conv ker } E$  and  $\text{conv } E$  will denote the convex kernel of  $E$  and convex hull of  $E$ , respectively. The interior of a set  $E$  relative to the smallest flat containing  $E$  is given by  $\text{intv } E$ . Finally,  $xy$  will denote the line segment joining  $x$  to  $y$  and  $L(x, y)$  will denote the line determined by  $x$  and  $y$ . We begin with this obvious fact: If  $E$  is a closed (open) set in  $E_n$ , then  $S(x)$  is closed (open) for each  $x$  in  $E$ . Let us designate the visibility

function for a fixed set by the letter  $v$ .

### 3. Continuity properties of the visibility function.

**THEOREM 1.** *If  $O \subset E_n$  is open, then the visibility function associated with  $O$  is lower semicontinuous.*

*Proof.* Let  $x \in O$  and let  $\{x_n\} \rightarrow x$ . Clearly  $S(x) \subset \bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} S(x_n)$ , for if  $p \in S(x)$ , there exists a neighborhood about  $x$  which sees  $p$  via  $O$ . Hence  $v(x) = m(S(x)) \leq m(\bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} S(x_n)) = \lim_{k \rightarrow \infty} m(\bigcap_{n=k}^{\infty} S(x_n)) \leq \varliminf_{k \rightarrow \infty} m(S(x_k)) = \varliminf_{k \rightarrow \infty} v(x_k)$ . Since  $v(x) = 0$  when  $x \notin O$  and  $O$  is open,  $v$  is lower semicontinuous.

**THEOREM 2.** *Let  $K \subset E_n$  be compact. Then the visibility function associated with  $K$  is upper semicontinuous.*

*Proof.* Again let  $x \in K$  and  $\{x_n\} \rightarrow x$ . The compactness of  $K$  implies that  $S(x) \supset \bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} S(x_n)$ . We have  $v(x) = m(S(x)) \geq m(\bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} S(x_n)) = \lim_{k \rightarrow \infty} m(\bigcup_{n=k}^{\infty} S(x_n)) \geq \overline{\lim}_{k \rightarrow \infty} m(S(x_k))$ . The result now follows as before.

**COROLLARY.** *The visibility function for a compact set when restricted to the set attains a maximum.*

**COROLLARY.** *The visibility function for a closed set is a Borel function.*

*Proof.* Let  $E$  be closed and  $x \in E$  be arbitrary. Then  $S(x) = \bigcup_{n=1}^{\infty} S(x) \cap B_n(0)$  so that  $v$  is the limit of upper semicontinuous functions.

**EXAMPLE.** A closed set  $E$  on which  $v$  is not upper semicontinuous. Let  $E$  be the following set in the plane:

$$(0, 1) \cup \text{cl} \bigcup_{n=1}^{\infty} \text{conv} ((1/n, 1) \cup \{(r, 0) : r \geq n\}) .$$

Then  $v(0, 1) = 0$  but  $v(1/n, 1) = \infty$  for all  $n$ .

**DEFINITION.** Let  $K$  be a compact set in  $E_n$ . The  $\varepsilon$ -parallel set of  $K$ , denoted by  $K_\varepsilon$ , is the compact set  $\bigcup_{x \in K} B_\varepsilon(x)$ .

Let  $S_{n-1}$  denote the unit sphere in  $E_n$ . Every  $x$  in  $E_n/\{0\}$  can be written uniquely as  $r \cdot \theta$  where  $r > 0$  and  $\theta \in S_{n-1}$ . If  $\sigma_{n-1}$  denotes the standard surface measure for  $S_{n-1}$  and if  $f$  is a positive Borel

function on  $E_n$ , recall that

$$\int_{E_n} f dm_n = \int_0^\infty \int_{S_{n-1}} r^{n-1} f(r \cdot \theta) d\sigma_{n-1}(\theta) dr .$$

LEMMA. *Let  $f$  be a nonnegative Borel function defined on  $S_{n-1}$ . If  $x_\theta = f(\theta) \cdot \theta$  for each  $\theta \in S_{n-1}$ , then  $\bigcup_\theta x_\theta$  is a set of  $n$ -dimensional measure zero.*

*Proof.* By Lusin's Theorem,  $\forall n$  there exists a continuous function  $f_n$  on  $S_{n-1}$  such that  $f_n = f$  on a compact set whose surface measure differs from that of the unit sphere by less than  $1/n$ . Hence  $E = \bigcup_{n=1}^\infty \{x_\theta : f(\theta) = f_n(\theta)\}$  is an  $F_\sigma$ -set differing from  $\bigcup_\theta x_\theta$  by a set of measure zero so that  $\bigcup_\theta x_\theta$  is measurable. Since the characteristic function of  $E$  is Borel we have

$$m(\bigcup_\theta x_\theta) = m(E) = \int_{S_{n-1}} \int_0^\infty r^{n-1} \chi_E(r \cdot \theta) dr d\sigma_{n-1}(\theta) = 0$$

as each ray emanating from the origin intersects  $E$  in at most one point.

The following corollary is indispensable in studying the visibility function.

COROLLARY. *Let  $E$  be a compact set in  $E_n$ . If  $x \in E$ , the set of endpoints of all segments in  $S(x)$  with one end point  $x$  forms a measurable set and has measure zero.*

*Proof.* We may assume  $x = 0$ . Let  $p_\theta$  denote the endpoint in the  $\theta$  direction (which may be 0). If we apply the previous lemma to the upper semicontinuous function  $f(\theta) = |p_\theta - 0|$ , the result follows.

#### 4. The index of convexity.

DEFINITION. Let  $E \subset E_n$  be a measurable set with measurable visibility function  $v$ , and suppose  $m(E) < \infty$ . If  $m(E) > 0$ , the Index of convexity of  $E$ ,  $I(E)$ , is given by

$$\int_E \frac{v}{m(E)^2} dm .$$

If  $m(E) = 0$ , we agree to let  $I(E)$  be 1.

Clearly  $0 \leq I(E) \leq 1$  for any set  $E$  for which the Index makes sense, and if  $E$  has a nonempty interior, the Index is nonzero. It is evident that the class of sets of Index 1 is closed under countable

intersections and contains the bounded convex sets as a subclass.

**THEOREM 3.** *Suppose  $F_1 \subset E_n$  and  $F_2 \subset E_1$  are two sets of positive measure for which the Index is defined. Then  $I(F_1 \times F_2) = I(F_1)I(F_2)$ .*

*Proof.* Let  $v$  be the visibility function for  $F_1 \times F_2$ . Clearly  $S(x) \times S(y) = S(x, y)$  for every  $(x, y) \in F_1 \times F_2$  so that  $v(x, y) = v_1(x)v_2(y)$ , where  $v_i$  is the visibility function for  $F_i$ ,  $i = 1, 2$ . The result now follows in the obvious way using Fubini's Theorem.

From the results of Theorems 1 and 2, it is immediate that the Index makes sense for compact and bounded open sets.

If  $\{E^n\}$  is a sequence of compact sets which converge to a compact set  $E$  in the Hausdorff metric, we might expect some relationship between  $\{I(E^n)\}$  and  $I(E)$ . However, we are working in the wrong metric space as the two following planar examples show.

(1) Let  $E = \{z: |z| \leq 1\} \cup \{z: |z - 2| \leq 1\}$ . Let  $E^n = \{z: |z - 2| \leq 1\} \cup \{z: |z| \leq 1, \text{Im } z = k/n \text{ for some integer } k\}$ . Then  $E^n \rightarrow E$ ,  $I(E^n) = 1$  for every  $n$ , but  $I(E) = 1/2$ .

(2) Let  $E = \{z: 0 \leq \text{Im } z \leq 1, |\text{Re } z| \leq 1\}$  and let  $E^n = \{z: 0 \leq \text{Im } z \leq 1, 1/n \leq |\text{Re } z| \leq 1\}$ . Then  $E^n \rightarrow E$ ,  $I(E^n) = 1/2$  for every  $n$ , but  $I(E) = 1$ .

What goes wrong is our improper generalization of convergence for convex sets, for if convex sets converge in the Hausdorff metric, they also converge in measure. We define a new metric  $\bar{d}$  on the class of compact sets in  $n$ -space. If  $A$  and  $B$  are two such sets, define  $\bar{d}(A, B) = \sup \{d(A, B), m(A \Delta B)\}$  where  $d$  denotes Hausdorff distance.

Although this metric space is not complete, we can still generalize that part of Blaschke's theorem which says that the limit of convex sets is convex.

**THEOREM 4.** *The Index of convexity is upper semicontinuous on the metric space  $\bar{d}$  of compact sets in  $n$ -space.*

*Proof.* We show that if  $K_n$  are compact and converge to  $K(\bar{d})$ , then  $I(K) \geq \limsup I(K_n)$ . We may assume  $m(K) > 0$  by our convention that  $I(K) = 1$  if  $m(K) = 0$ .

Let  $K' = \{x \in K: x \in K_n \text{ for infinitely many } n\}$ . We claim  $m(K') = m(K)$ . We must show  $m\{x: x \in K \cap K_n^c \text{ for all but finitely many } n\} = 0$ . Suppose  $m(\bigcup_{k=1}^{\infty} \bigcap_{n>k} K_n^c \cap K) > \delta > 0$ . Then there exists  $N$  such that  $m(\bigcap_{n>N} K_n^c \cap K) > \delta/2$ . Hence  $m(K_n^c \cap K) > \delta/2$  for  $n > N$ . But  $K_n^c \cap K \subset (K \Delta K_n)$ , a contradiction to  $m(K \Delta K_n) \rightarrow 0$ .

Let  $v$  denote the visibility function for  $K$  and let  $v_n$  denote the

visibility function for  $K_n$ .

We claim for each  $x$  in  $K'$ ,  $v(x) \geq \limsup v_n(x)$ . Let  $S_n(x) = \{y: yx \subset K_n\}$ . We need only show that for every  $\varepsilon > 0$ , there exists  $N$  such that  $n > N \Rightarrow (S(x))_\varepsilon \supset S_n(x)$  whenever  $x \in K_n$ . Suppose not. Then there exists a sequence of segments  $\{y_{n_l}x\}$  such that for each  $l$ ,  $y_{n_l}x \subset K_{n_l}$ , but  $y_{n_l}$  is not contained in  $(S(x))_\varepsilon$ . Passing to a subsequence we may assume that  $\{y_{n_l}x\}$  converges in the Hausdorff metric to a subset  $D \subset S(x)$ . Hence if  $l$  is sufficiently large  $y_{n_l}x \subset D_\varepsilon \subset (S(x))_\varepsilon$ , a contradiction.

We now have by Fatou's lemma

$$\begin{aligned} I(K) &= \int_{K'} \frac{v}{m(K')^2} dm \\ &\geq \int_{K'} \limsup \frac{v_n}{m(K_n)^2} dm \\ &\geq \limsup \int_{K'} \frac{v_n}{m(K_n)^2} dm \\ &= \limsup \int_{K' \cap K_n} \frac{v_n}{m(K_n)^2} dm + \lim \int_{K_n \setminus K'} \frac{v_n}{m(K_n)^2} dm \\ &= \limsup I(K_n) . \end{aligned}$$

Our second example shows that strict inequality may indeed occur.

**COROLLARY.** *If  $I(K_n) = 1$  for each  $n$  and if  $K_n \rightarrow K$  in the  $\bar{d}$  metric, then  $I(K) = 1$ .*

**5. Pseudo kernels.** Sets of Index 1 are the obvious generalization of convex sets in terms of "visibility theory." Analogously, a set  $E$  is starshaped in a more general sense if there is a point  $x$  of  $E$  satisfying  $m(E) = m(S(x))$ . In this section we classify in the compact case sets of Index 1 and "pseudo starshaped" sets.

**DEFINITION.** Let  $E$  be a measurable set in  $E_n$ . The pseudo kernel of  $E$  is the set  $\{x \in E: v(x) = m(E)\}$ . We denote this set by  $P \ker E$ .  $E$  is pseudo starshaped if  $P \ker E \neq \emptyset$ .

**LEMMA.** *If  $E \subset E_n$  is open,  $P \ker E$  is a  $G_\delta$ -set. If  $K \subset E_n$  is compact,  $P \ker K$  is compact.*

*Proof.* If  $E$  is open,  $P \ker E = \bigcap_{n=1}^\infty \{x \in E: v(x) > m(E) - 1/n\}$ . If  $K$  is compact and  $m(K) = 0$  the assertion is obviously true; otherwise, Theorem 2 implies that  $\{x \in K: v(x) \geq m(E)\}$  is compact.

LEMMA. Let  $K$  be a compact set in  $E_n$ . If  $\text{conv ker } K$  has a nonempty interior, then  $\text{conv ker } K = P \text{ ker } K$ .

*Proof.* Let  $y \in P \text{ ker } K / \text{conv ker } K$ . Since  $K^c$  is open,  $y$  must see every point of  $\text{int conv } (x \cup \text{conv ker } K)$  where  $x \in K$  is arbitrary. Since  $K$  is compact,  $x \in S(y)$  for every  $x \in K$ .

EXAMPLE. A compact set  $K$  in  $E_3$  satisfying  $\dim(\text{conv ker } K) = 2$  but  $\text{conv ker } K \neq P \text{ ker } K$ . Let  $K = \{(x, y, z): x^2 + y^2 + z^2 \leq 1\} \cup \{(x, y, z): x^2 + y^2 \leq 2, z = 0\}$ .

The following property of the visibility function peculiar to compact sets is particularly illuminating in this section.

THEOREM 5. Let  $E$  and  $F$  be compact sets in  $E_n$ . If  $m(E \Delta F) = 0$ , then the visibility functions for the two sets are equal.

*Proof.* We first show the two functions agree on  $E \cap F$ . W.l.o.g., we may assume  $0$  to be an arbitrary point of  $E \cap F$ . Let  $S_F(0)$  be those points of  $F$  which  $0$  sees via  $F$  and let  $S_E(0)$  be those points of  $E$   $0$  sees via  $E$ . Suppose  $m(S_E(0)) > m(S_F(0))$  so that certain segments in  $S_E(0)$  contain segments in  $F^c$ .

Consider the function  $f$  defined on  $S_{n-1}$  given by  $f(\theta) = \sup \{|x - y|: xy \subset F^c \cap R_\theta \cap S_E(0)\}$  where  $R_\theta$  denotes the ray from  $0$  through  $\theta$ . We claim  $f$  is a Borel function. To see this, note that  $f(\theta) = \inf f_n(\theta)$  where  $f_n(\theta) = \sup \{|x - y|: xy \subset F^c \cap R_\theta \cap (S_E(0))_{1/n}\}$ . However, for every  $n$  the  $1/n$ -parallel set about  $S_E(0)$  is a compact starshaped set whose convex kernel has a nonempty interior. Hence, the boundary is a continuous function of  $\theta$  which implies  $f_n$  is lower semicontinuous for every  $n$ , so that  $f$  is a Borel function. Now we must have  $\sigma_{n-1}(f^{-1}(0, \infty)) > 0$ . Hence there exists  $\delta > 0$  such that  $\sigma_{n-1} \{ \theta: \text{there exists } xy \subset F^c \cap R_\theta \cap S_E(0), |x - y| > \delta \} > 0$ . Since this set, which we denote by  $M_\delta$ , is a Borel set, the set of all points in  $F^c \cap S_E(0)$  which project radially onto  $M_\delta$  is a Borel set. Denoting this last set by  $M$  we have

$$\begin{aligned} \int \chi_M dm_n &= \int_{M_\delta} \int_0^\infty r^{n-1} \chi_M(r \cdot \theta) dr d\sigma_{n-1}(\theta) \\ &\geq \frac{\delta^n}{n} \sigma_{n-1}(M_\delta) > 0. \end{aligned}$$

But  $M \subset S_E(0) \cap F^c \subset E \cap F^c$ . Hence we have a contradiction to  $m(E \Delta F) = 0$ . It is easy to show that both visibility functions are zero at any point  $x$  in  $E \Delta F$  since  $E \cap F$  is compact.

Intuitively we should only expect that  $I(P \text{ ker } K) = 1$  and not

that  $P \ker K$  is convex. It is surprising that the following theorem holds.

**THEOREM 6.** *Let  $K$  be any compact set in  $E_n$  with  $m(K) > 0$ . Then  $P \ker K$  is convex.*

*Proof.* Again we may assume that  $0$  is an arbitrary element of  $P \ker K$ , and let  $y$  be any other element of  $P \ker K$ . Since  $m(S(0)) > 0$ , every neighborhood of  $0$  must contain a subset of  $K$  of positive measure so that  $y$  must see some point in each neighborhood via  $K$ . The compactness of  $K$  implies  $y$  sees  $0$  via  $K$ .

We claim  $0y \subset P \ker K$ . Consider the Borel set  $S(0) \cap S(y)^c$ , which has measure zero. We must have for almost every  $\theta$  on  $S_{n-1}$

$$\int_0^\infty r^{n-1} \chi_{S(0) \cap S(y)^c}(r \cdot \theta) dr = 0.$$

Since  $R_\theta \cap S(0) \cap S(y)^c$  is either empty or contains open segments, we have  $R_\theta \cap S(0) \cap S(y)^c = \emptyset$  for almost all  $\theta$ . This means that  $y$  sees every line segment in  $S(0)$  in its entirety except a set of lines whose union forms a set of measure zero. Hence any point in  $\text{intv } 0y$  has the same property. Since almost every point of  $K$  is in  $S(0)$ ,  $0y \subset P \ker K$ .

**COROLLARY.** *Let  $K$  be a compact set in  $E_n$ . Then  $I(K) = 1$  iff  $K = F \cup C$  where  $C$  is compact and convex and  $m(F) = 0$ .*

**COROLLARY.** (*Helly's Theorem for compact sets of Index 1.*) *Let  $\{K_\alpha\}$  be a collection of compact sets of Index 1 in  $E_n$  such that every  $n + 1$  intersect in a set of positive measure. Then  $\bigcap K_\alpha \neq \emptyset$ .*

**COROLLARY.** *If  $K \subset E_n$  is a compact set which is the closure of an open set and  $I(K) = 1$ , then  $K$  is convex.*

As a consequence of Theorem 5 we have the following

**COROLLARY.** *If a compact set  $K \subset E_n$  satisfies  $m(K) > 0$ ,  $I(K) = 1$  and  $v(x) > 0$  for every  $x$  in  $K$ , then  $K$  is convex.*

We are now in a position to classify those compact sets with nonempty pseudo kernels.

**THEOREM 7.** *Let  $K$  be a compact set in  $E_n$  with  $m(K) > 0$ . Suppose  $P \ker K \neq \emptyset$ . Then  $K = S \cup F$  where  $m(F) = 0$  and  $S$  is a compact starshaped set with convex kernel  $P \ker K$ . In addition,  $m(S(x)) = 0$  for every  $x \in F$ .*



*Proof.* Let  $\{a_n\}$  be a countable dense subset of  $P \ker K$ .  $m(\bigcup_{n=1}^{\infty} K/S(a_n)) = 0$  so that  $m(\bigcap_{n=1}^{\infty} S(a_n)) = m(K)$ . The compactness of  $K$  implies

$$\bigcap_{x \in P \ker K} S(x) = \bigcap_{n=1}^{\infty} S(a_n) \text{ so that } m(K) = m\left(\bigcap_{x \in P \ker K} S(x)\right).$$

Let  $y \in \bigcap_{x \in P \ker K} S(x)$  and  $x_0 \in P \ker K$  be arbitrary. If  $q \in \text{intv } x_0 y$ , we have  $\text{conv}(q \cup P \ker K) \subset \text{conv}(y \cup P \ker K) \subset K$  so that  $q \in \bigcap_{x \in P \ker K} S(x)$ . Hence  $\bigcap_{x \in P \ker K} S(x)$  is starshaped with respect to  $P \ker K$ , is compact and is of full measure. If  $y \in K / \bigcap_{x \in P \ker K} S(x)$ , Theorem 5 again implies  $v(y) = 0$ .

We note that any compact pseudo starshaped set has a visibility function identical with the visibility function of some compact starshaped set.

**THEOREM 8.** *Let  $\{K_n\}$  be a sequence of compact sets of positive measure with nonempty pseudo kernels  $\{P_n\}$  and suppose  $\{K_n\} \rightarrow K(\bar{d})$ . Then  $K$  has a pseudo kernel which contains a subsequential limit of  $\{P_n\}$ .*

*Proof.* Let  $K_n = S_n \cup F_n$  where  $S_n$  is compact and starshaped with respect to  $P_n$  and  $m(F_n) = 0$ . By the Blaschke convergence theorem for starshaped sets, passing to a subsequence we may assume  $\{S_n\} \rightarrow S$  and  $\{P_n\} \rightarrow P$  in the Hausdorff metric, where  $P \subset \text{conv } \ker S$ . We have  $m(S) \geq \limsup m(S_n) = \lim m(K_n) = m(K)$ . Since  $S \subset K$ , we have  $m(K/S) = 0$ .

Since the compact pseudo starshaped sets include all compact sets of measure zero, it follows from Theorem 8 that they form a closed metric subspace of the metric space of all compact sets under  $\bar{d}$ . It fails to be continuous on this subspace; to see this, we note that the unit disc in  $E_2$  is the limit of a sequence of radial Cantor sets of positive measure.

No classification theorem for open sets of Index 1 has yet been established. Clearly, if  $E$  is a bounded open set of Index 1, then  $\text{cl } E$  is convex which implies that a regular bounded open set is convex iff it has Index 1. It seems reasonable to conjecture that the Index of a pseudo kernel of a bounded open set would be 1. A property shared in common with convex open sets is the following.

**THEOREM 9.** *Let  $\{E_m\}$  be a countable collection of bounded open sets in  $E_n$  of Index 1 such that every  $n + 1$  contain a translate of some closed ball  $B$ . Then  $\bigcap_{m=1}^{\infty} E_m \neq \emptyset$ .*

*Proof.* The collection  $\{\text{cl}(E_m)\}$  satisfy the hypotheses of the following theorem of Klee [1]: If  $\{K_\alpha\}$  is a collection of compact convex sets in  $E_n$  such that the intersection of every subfamily of cardinality  $n + 1$  contains a translate of some convex compact set  $K$ , then  $\bigcap K_\alpha$  contains a translate of  $K$ . Therefore,  $\bigcap_{m=1}^{\infty} \text{cl}(E_m)$  contains  $B + p$  for some  $p \in E_n$ . Now  $\text{bd } E_m$  is nowhere dense in  $B + p$  for each  $m$ . Since  $B + p$  is a complete metric space,  $\bigcap_{m=1}^{\infty} E_m$  is nonempty.

EXAMPLE. For each  $(r_1, r_2) \in E_2$ , let

$$E_{r_1, r_2} = \{(x, y): 0 < x < 1, 0 < y < 1, (x, y) \neq (r_1, r_2)\}.$$

It is easy to see that the intersection of any three such sets contains a ball of radius  $1/12$ , but  $\bigcap_{(r_1, r_2) \in E_2} E_{r_1, r_2} = \emptyset$ .

#### REFERENCES

1. V. L. Klee, *The critical set of a convex body*, Amer. J. Math., **75** (1953), 178-188.
2. Walter Rudin, *Real and Complex Analysis*, McGraw-Hill, New York, 1966.
3. F. A. Valentine, *Convex Sets*, McGraw-Hill, New York, 1964.

Received September 14, 1971 and in revised form July 28, 1972.

CALIFORNIA STATE COLLEGE, LOS ANGELES



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

C. R. HOBBY  
University of Washington  
Seattle, Washington 98105

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Jimmy T. Arnold, <i>Power series rings over Prüfer domains</i> .....	1
Maynard G. Arsove, <i>On the behavior of Pincherle basis functions</i> .....	13
Jan William Auer, <i>Fiber integration in smooth bundles</i> .....	33
George Bachman, Edward Beckenstein and Lawrence Narici, <i>Function algebras over valued fields</i> .....	45
Gerald A. Beer, <i>The index of convexity and the visibility function</i> .....	59
James Robert Boone, <i>A note on mesocompact and sequentially mesocompact spaces</i> .....	69
Selwyn Ross Caradus, <i>Semiclosed operators</i> .....	75
John H. E. Cohn, <i>Two primary factor inequalities</i> .....	81
Mani Gagrat and Somashekhar Amrith Naimpally, <i>Proximity approach to semi-metric and developable spaces</i> .....	93
John Grant, <i>Automorphisms definable by formulas</i> .....	107
Walter Kurt Hayman, <i>Differential inequalities and local valency</i> .....	117
Wolfgang H. Heil, <i>Testing 3-manifolds for projective planes</i> .....	139
Melvin Hochster and Louis Jackson Ratliff, Jr., <i>Five theorems on Macaulay rings</i> .....	147
Thomas Benton Hoover, <i>Operator algebras with reducing invariant subspaces</i> .....	173
James Edgar Keesling, <i>Topological groups whose underlying spaces are separable Fréchet manifolds</i> .....	181
Frank Leroy Knowles, <i>Idempotents in the boundary of a Lie group</i> .....	191
George Edward Lang, <i>The evaluation map and EHP sequences</i> .....	201
Everette Lee May, Jr., <i>Localizing the spectrum</i> .....	211
Frank Belsley Miles, <i>Existence of special <math>K</math>-sets in certain locally compact abelian groups</i> .....	219
Susan Montgomery, <i>A generalization of a theorem of Jacobson. II</i> .....	233
T. S. Motzkin and J. L. Walsh, <i>Equilibrium of inverse-distance forces in three-dimensions</i> .....	241
Arunava Mukherjea and Nicolas A. Tserpes, <i>Invariant measures and the converse of Haar's theorem on semitopological semigroups</i> .....	251
James Waring Noonan, <i>On close-to-convex functions of order <math>\beta</math></i> .....	263
Donald Steven Passman, <i>The Jacobian of a growth transformation</i> .....	281
Dean Blackburn Priest, <i>A mean Stieltjes type integral</i> .....	291
Joe Bill Rhodes, <i>Decomposition of semilattices with applications to topological lattices</i> .....	299
Claus M. Ringel, <i>Socle conditions for <math>QF-1</math> rings</i> .....	309
Richard Rochberg, <i>Linear maps of the disk algebra</i> .....	337
Roy W. Ryden, <i>Groups of arithmetic functions under Dirichlet convolution</i> .....	355
Michael J. Sharpe, <i>A class of operators on excessive functions</i> .....	361
Erling Stormer, <i>Automorphisms and equivalence in von Neumann algebras</i> .....	371
Philip C. Tonne, <i>Matrix representations for linear transformations on series analytic in the unit disc</i> .....	385