

Pacific Journal of Mathematics

ON CLOSE-TO-CONVEX FUNCTIONS OF ORDER β

JAMES WARING NOONAN

ON CLOSE-TO-CONVEX FUNCTIONS OF ORDER β

JAMES W. NOONAN

For $\beta \geq 0$, denote by $K(\beta)$ the class of normalized functions f , regular and locally schlicht in the unit disc, which satisfy the condition that for each $r < 1$, the tangent to the curve $C(r) = \{f(re^{i\theta}) : 0 \leq \theta < 2\pi\}$ never turns back on itself as much as $\beta\pi$ radians. $K(\beta)$ is called the class of close-to-convex functions of order β . The purpose of this paper is to investigate the asymptotic behavior of the integral means and Taylor coefficients of $f \in K(\beta)$. It is shown that the function F_β , given by $F_\beta(z) = (1/(2(\beta+1)))\{((1+z)/(1-z))^{\beta+1} - 1\}$, is in some sense extremal for each of these problems. In addition, the class $B(\alpha)$ of Bazilevic functions of type $\alpha > 0$ is related to the class $K(1/\alpha)$. This leads to a simple geometric interpretation of the class $B(\alpha)$ as well as a geometric proof that $B(\alpha)$ contains only schlicht functions.

Let f be regular in $U = \{z : |z| < 1\}$ and be given by

$$(1.1) \quad f(z) = z + a_2 z^2 + a_3 z^3 + \dots$$

Following an argument due to Kaplan [9], we see that $f \in K(\beta)$ iff, for some normalized convex function φ and some constant c with $|c| = 1$, we have for all $z \in U$ that

$$(1.2) \quad \left| \arg \frac{cf'(z)}{\varphi'(z)} \right| \leq \beta\pi/2.$$

Equivalently,

$$(1.3) \quad cf'(z) = p(z)^\beta \varphi'(z),$$

where $p(z) = \sum_{n=0}^{\infty} p_n z^n$, $|p_0| = 1$, has positive real part in U .

It is geometrically clear that for $0 \leq \beta \leq 1$, $K(\beta)$ contains only schlicht functions. However, for any $\beta > 1$, Goodman [3] has shown that $K(\beta)$ contains functions of arbitrarily high valence. $K(0)$ is the class of convex functions, and $K(1)$ is the class of close-to-convex functions introduced by Kaplan [9]. For $0 \leq \alpha \leq 1$, Pommerenke [13, 14] has studied m -fold symmetric functions of class $K(\alpha)$. The following theorem shows that the study of these functions is closely related to the study of $K(\beta)$ for arbitrary $\beta \geq 0$.

THEOREM 1.1. *Let $\beta \geq 0$ and m be a positive integer. Then $f \in K(\beta)$ iff there exists an m -fold symmetric function $g \in K(\beta/m)$ such that $f'(z^m) = g'(z)^m$.*

Proof. Suppose $f \in K(\beta)$, and define g by $g'(z) = f'(z^m)^{1/m}$. From (1.3) it follows that

$$g'(z) = e^{-1/m} p(z^m)^{\beta/m} \psi'(z)$$

where the convex function ψ is defined by $\psi'(z) = \varphi'(z^m)^{1/m}$. Hence $g \in K(\beta/m)$, and g is clearly m -fold symmetric. To prove the converse implication, we merely reverse the above procedure.

Finally, for $k \geq 2$ denote by V_k the class of normalized functions with boundary rotation at most $k\pi$. From the proof of [2, Theorem 2.2], it follows that $V_k \subset K(k/2 - 1)$. However, $f \in V_k$ implies that f is at most $k/2$ valent [2], so $K(k/2 - 1)$ is in general a much larger class than V_k . The results in §2 and 3 of this paper are extensions to $K(\beta)$ of results of the author [10] for the class V_k . These results also generalize and improve some of the results of Pommerenke [13] for $K(\alpha)$, $0 \leq \alpha \leq 1$.

2. Behavior of the coefficients. We begin by studying $M(r, f') = \max_{|z|=r} |f'(z)|$.

THEOREM 2.1. *Let $f \in K(\beta)$. Then $((1-r)/(1+r))^{\beta+2} M(r, f')$ is a decreasing function of r , and hence $\omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} M(r, f')$ exists and is finite. If $\omega > 0$ and f is given by (1.3), then there exists θ_0 such that $\varphi'(z) = (1 - ze^{-i\theta_0})^{-2}$ and $\omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} |f'(re^{i\theta_0})|$.*

Proof. Since for each $\beta \geq 0$, $K(\beta)$ is a linear-invariant family of order $\beta + 1$ in the sense of Pommerenke [12] (See [4, Theorem 3] for a proof.), the first two statements of the theorem follow. Also, if φ' is not of the stated form, then $\varphi'(z) = O(1)(1-r)^{-\delta}$ for some $0 < \delta < 2$, and hence from (1.3) we see $\omega = 0$. Finally, if $\omega > 0$, then $\varphi'(z) = (1 - ze^{-i\theta_0})^{-2}$, and just as in the proof of [10, Theorem 3.1] we see that $\omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} |f'(re^{i\theta_0})|$.

We now begin to study the coefficient behavior. Our method is the major-minor arc technique used by Hayman [5], and the proofs are similar to the proofs of the corresponding results for the class V_k [10]. Hence we omit details wherever possible. We first require two lemmas.

LEMMA 2.1 *Let $f \in K(\beta)$ and $\omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} |f'(re^{i\theta_0})| > 0$. Then given $\delta > 0$, we may choose $C = C(\delta) > 0$ and $r_0 = r_0(\delta) < 1$ such that for $r_0 \leq r < 1$ we have*

$$\int_E |f'(re^{i\theta})| d\theta < \frac{\delta}{(1-r)^{\beta+1}}$$

where $E = \{\theta: C(\delta)(1-r) \leq |\theta - \theta_0| \leq \pi\}$.

Proof. Without loss of generality we may assume $\theta_0 = 0$, so from Theorem 2.1 and (1.3) we find, with $z = re^{i\theta}$,

$$|f'(z)| = |p(z)|^\beta |1-z|^{-2}.$$

Hence, with $C > 0$ and E as above, we find

$$\int_E |f'(z)| d\theta = \frac{O(1)}{(1-r)^\beta} \int_{C(1-r)}^\pi \theta^{-2} d\theta = O(1) \frac{1}{C} \frac{1}{(1-r)^{\beta+1}},$$

and the lemma now follows upon choosing C sufficiently large.

LEMMA 2.2. Let $f \in K(\beta)$, $\omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} |f'(re^{i\theta_0})| > 0$, $r_n = 1 - 1/n$, $\omega_n = (1-r_n)^{\beta+2} f'(r_n e^{i\theta_0})$, and

$$f'_n(z) = \frac{\omega_n}{(1 - ze^{-i\theta_0})^{\beta+2}}.$$

Let S be a fixed but arbitrary Stolz angle with vertex $e^{i\theta_0}$, and let $D_n = \{z \in S: |e^{i\theta_0} - z| < 2/n\}$. Then as $n \rightarrow \infty$, $f'_n \sim f'$ uniformly for $z \in D_n$.

Proof. Again assuming $\theta_0 = 0$, we have from (1.3) $cf'(z) = p(z)^\beta (1-z)^{-2}$, and so

$$f'_n(z) = \frac{[(1-r_n)p(r_n)]^\beta}{c(1-z)^{\beta+2}}.$$

Thus, to prove the lemma it suffices to show that as $n \rightarrow \infty$,

$$(2.1) \quad \frac{(1-r_n)p(r_n)}{(1-z)p(z)} \longrightarrow 1$$

uniformly for $z \in D_n$.

By a theorem of Hayman [6, Theorem 2], $\lim_{r \rightarrow 1} (1-r)p(r) = L$ exists, and it is clear that $(1-z)p(z)$ is bounded as $|z| \rightarrow 1$, providing $z \in S$. By a theorem of Lindelöf [8, p. 260], we have for $z \in S$ that $\lim_{z \rightarrow 1} (1-z)p(z) = L$ where the limit is approached uniformly as $|z| \rightarrow 1$. But $0 < \omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} |f'(r)| = \lim_{r \rightarrow 1} [(1-r)|p(r)|]^\beta$, so $L \neq 0$. Combining these remarks with the inequality

$$\begin{aligned} & \left| \frac{(1-z)p(z)}{(1-r_n)p(r_n)} - 1 \right| \\ & \leq \frac{1}{|(1-r_n)p(r_n)|} \{ |(1-z)p(z) - L| + |L - (1-r_n)p(r_n)| \}, \end{aligned}$$

we see that (2.1) holds, so the proof is complete.

We can now determine the asymptotic behavior of a_n as $n \rightarrow \infty$.

THEOREM 2.2. *Let $f \in K(\beta)$ be given by (1.1), and let $\omega = \lim_{r \rightarrow 1} (1 - r)^{\beta+2} M(r, f')$. Let Γ denote the gamma function. Then*

$$\lim_{n \rightarrow \infty} \frac{|a_n|}{n^\beta} = \frac{\omega}{\Gamma(\beta + 2)}.$$

Also, if $\omega = \lim_{r \rightarrow 1} (1 - r)^{\beta+2} |f'(re^{i\theta_0})| > 0$, then as $n \rightarrow \infty$

$$a_n \sim \frac{f'(r_n e^{i\theta_0}) e^{-i(n-1)\theta_0}}{n^2 \Gamma(\beta + 2)}$$

where $r_n = 1 - 1/n$.

Proof. Suppose first that $\omega > 0$, and define

$$f'_n(z) = \omega_n \sum_{m=0}^{\infty} d^m e^{-im\theta_0} z^m$$

as in Lemma 2.2. We note that

$$(2.2) \quad d_m = \frac{\Gamma(m + \beta + 2)}{\Gamma(m + 1) \Gamma(\beta + 2)},$$

so $d_m \sim m^{\beta+1}/\Gamma(\beta + 2)$ as $m \rightarrow \infty$. Computation shows that

$$(2.3) \quad na_n - \omega_n d_{n-1} e^{-i(n-1)\theta_0} = \frac{1}{2\pi r^{n-1}} \int_{-\pi}^{\pi} \{f'(re^{i\theta}) - f'_n(re^{i\theta})\} e^{-i(n-1)\theta} d\theta.$$

Given $\delta > 0$, we choose $C = C(\delta)$ and E as in Lemma 2.1, and we let $r_n = 1 - 1/n$. With n sufficiently large, Lemma 2.1 gives

$$\int_E |f'(r_n e^{i\theta})| d\theta < \delta n^{\beta+1},$$

and clearly this inequality is also true for f'_n . Hence, we see that

$$(2.4) \quad \left| \int_E \{f'(r_n e^{i\theta}) - f'_n(r_n e^{i\theta})\} e^{-i(n-1)\theta} d\theta \right| < 2\delta n^{\beta+1}$$

for n sufficiently large. We now choose a Stolz angle S , depending on δ , such that $\{r_n e^{i\theta} : \theta \in E'\} \subset S$ for large n , where $E' = [-\pi, \pi] \setminus E$. By Lemma 2.2, we have as $n \rightarrow \infty$ and with $\theta \in E'$,

$$\begin{aligned} f'(r_n e^{i\theta}) - f'_n(r_n e^{i\theta}) &= o(1) \{f'_n(r_n e^{i\theta})\} \\ &= o(1) n^{\beta+2}, \end{aligned}$$

where $o(1)$ is uniform for $\theta \in E'$, and hence as $n \rightarrow \infty$, we have

$$(2.5) \quad \left| \int_{E'} \{f'(r_n e^{i\theta}) - f'_n(r_n e^{i\theta})\} e^{-i(n-1)\theta} d\theta \right| \leq o(1)2C(\delta)(1 - r_n)n^{\beta+2} \\ = o(1)n^{\beta+1}.$$

Note that although $o(1)$ depends on δ , $o(1) \rightarrow 0$ as $n \rightarrow \infty$ once δ has been fixed.

Combining (2.3), (2.4), and (2.5), we find

$$|na_n - \omega_n d_{n-1} e^{-i(n-1)\theta_0}| < \{2\delta + o(1)\}n^{\beta+1}$$

for sufficiently large n . Since $\delta > 0$ is arbitrary and since $o(1) \rightarrow 0$ once δ has been fixed, we have

$$a_n = \omega_n \frac{d_{n-1}}{n} e^{-i(n-1)\theta_0} + o(1)n^\beta.$$

From (2.2) and the definition of ω_n we see that as $n \rightarrow \infty$,

$$a_n \sim \omega_n e^{-i(n-1)\theta_0} n^\beta / \Gamma(\beta + 2) \\ \sim \frac{f'(r_n e^{i\theta_0}) e^{-i(n-1)\theta_0}}{n^2 \Gamma(\beta + 2)}.$$

In particular,

$$\lim_{n \rightarrow \infty} \frac{|a_n|}{n^\beta} = \frac{\omega}{\Gamma(\beta + 2)}.$$

We now suppose $\omega = 0$. We shall subsequently prove (Theorem 3.1 with $\lambda = 1$) that if $\omega = 0$, then

$$\lim_{r \rightarrow 1} (1 - r)^{\beta+1} \int_0^{2\pi} |f'(re^{i\theta})| d\theta = 0.$$

Using a standard inequality relating coefficients and integral means [7, p. 11] we have $\lim_{n \rightarrow \infty} |a_n|/n^\beta = 0$. This completes the proof of the theorem. Note that if $\omega > 0$, then it follows easily from the theorem that $\lim_{n \rightarrow \infty} a_{n+1}/a_n = e^{-i\theta_0}$, and so the radius of maximal growth can be determined from the coefficients.

We now consider the problem of determining

$$\max \{|a_n| : f \in K(\beta)\}.$$

It is natural to conjecture that for each $n \geq 2$ this problem is solved by the function

$$F_\beta(z) = \frac{1}{2(\beta + 1)} \left\{ \left(\frac{1+z}{1-z} \right)^{\beta+1} - 1 \right\} = z + \sum_{j=2}^{\infty} A_j(\beta) z^j.$$

Toward this end we have the following theorem.

THEOREM 2.3. *Let $f \in K(\beta)$ be given by (1.1) and let F_β be as above.*

(i) *There exists an integer n_0 depending on f such that $|a_n| \leq A_n(\beta)$ for $n \geq n_0$.*

(ii) *If $n \leq \beta + 2$, then $|a_n| \leq A_n(\beta)$.*

(iii) *If β is an integer, then $|a_n| \leq A_n(\beta)$ for all n .*

Note that since $V_k \subset K(\beta)$ with $\beta = k/2 - 1$, we have from (ii) that $|a_n| \leq A_n(\beta)$ for $n \leq k/2 + 1$ and from (iii) that $|a_n| \leq A_n(\beta)$ for all n whenever k is an even integer.

Proof. We have from (1.3), with $|c| = 1$,

$$cf'(z) = p(z)^\beta \varphi'(z),$$

where p has positive real part and φ is convex. Suppose that $p(z) = \sum_{n=0}^{\infty} p_n z^n$, $|p_0| = 1$, and $p(z)^\beta = \sum_{n=0}^{\infty} q_n z^n$. Then it is easily verified by induction that for $m \geq 1$,

$$q_m = \frac{1}{m!} \sum_{j=1}^m \beta(\beta-1) \cdots (\beta-(j-1)) p_0^{\beta-j} D_j(p)$$

where $D_j(p)$ is a polynomial, with nonnegative coefficients, in the variables p_0, p_1, \dots, p_m .

Therefore, if β is an integer, $|q_m|$ is maximal for all $m \geq 1$ when $p_0 = 1$ and $p_j = 2$ for $j \geq 1$, which implies $p(z) = (1+z)/(1-z)$. Also, for any $\beta \geq 0$, we see as above that if $n \leq \beta + 2$, then $|q_m|$ is maximal for $1 \leq m \leq n-1$ when $p(z) = (1+z)/(1-z)$. In addition, if $\varphi'(z) = 1 + \sum_{j=2}^{\infty} u_j z^{j-1}$, it is well-known that $|u_j| \leq j$ for all j , with equality for $\varphi'(z) = (1-z)^{-2}$. But when $p(z) = (1+z)/(1-z)$ and $\varphi'(z) = (1-z)^{-2}$, we have $cf'(z) = F''(z)$. Hence, since

$$cna_n = \sum_{j=0}^{n-1} q_j u_{n-j}$$

where we define $u_1 = 1$, we see that (ii) and (iii) are proved.

We now prove (i). We first note that as $n \rightarrow \infty$,

$$(2.6) \quad A_n(\beta) \sim \frac{2^\beta n^\beta}{\Gamma(\beta+2)}.$$

Let $\omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} M(r, f')$. If $\omega = 0$, then Theorem 2.2 shows $a_n = o(1)n^\beta$, and so it is clear from (2.6) that (i) holds. We now suppose $\omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} |f'(re^{i\theta_0})| > 0$, and we recall that in this case $\omega = \lim_{r \rightarrow 1} [(1-r) |p(re^{i\theta_0})|]^\beta$. Hence, from [6, Theorem 2], it follows easily that $\omega \leq 2^\beta$ with equality only if

$$p(z) = \frac{1 + ze^{-i\theta_0}}{1 - ze^{-i\theta_0}}.$$

But $\omega > 0$ implies also that $\varphi'(z) = (1 - ze^{-i\theta_0})^{-2}$, and thus we have $\omega \leq 2^\beta$ with equality only if $cf'(z) = F'_\beta(e^{-i\theta_0}z)$, in which case $|a_n| = A_n(\beta)$ for all n , since $|c| = 1$. Thus we may suppose $\omega < 2^\beta$, and using Theorem 2.2 and (2.6) we see that (i) holds. This completes the proof of Theorem 2.3.

To conclude this section we examine the asymptotic behavior of the quantity $||a_{n+1}| - |a_n||$ for $f \in K(\beta)$.

THEOREM 2.4. *Let $f \in K(\beta)$ be given by (1.1). If $\omega > 0$, then*

$$\lim_{n \rightarrow \infty} \frac{||a_{n+1}| - |a_n||}{n^{\beta-1}} = \frac{\beta\omega}{\Gamma(\beta + 2)}.$$

The theorem is in general false when $\omega = 0$.

Proof. If $\beta = 0$ and $\omega > 0$, then from (1.3) it follows that $cf'(z) = (1 - ze^{-i\theta_0})^{-2}$, so $|a_n| = 1$ for all n , and the theorem is trivially true. Thus, we may assume without loss of generality that $\beta > 0$. The proof will be divided into three parts.

We first claim that given $\delta > 0$, there exists $C(\delta) > 0$ such that

$$(2.7) \quad \left| \frac{1}{2\pi} \int_E (1 - re^{i(\theta - \theta_0)}) f'(re^{i\theta}) d\theta \right| < \frac{\delta}{(1 - r)^\beta}$$

where θ_0 is as in Theorem 2.1 and $E = \{\theta: C(\delta)(1 - r) \leq |\theta - \theta_0| \leq \pi\}$. To prove (2.7), we note that $\omega > 0$ implies that

$$cf'(z) = p(z)^\beta(1 - z)^{-2},$$

where we have assumed without loss of generality that $\theta_0 = 0$. Also, for notational ease, we assume $c = 1$ and $p(0) = 1$, so

$$(1 - z)f'(z) = p(z)^\beta/(1 - z).$$

Choose $\lambda > 1$ such that $\lambda\beta > 1$, and let $1/\lambda + 1/\lambda' = 1$. If C is an arbitrary positive constant, we have from Hölder's inequality that

$$(2.8) \quad \int_E |(1 - z)f'(z)| d\theta \leq \left\{ \int_E |p(z)|^{\lambda\beta} d\theta \right\}^{1/\lambda} \left\{ \int_E |1 - z|^{-\lambda'} d\theta \right\}^{1/\lambda'}.$$

Since p is subordinate to $(1 + z)/(1 - z)$, and since $\lambda\beta > 1$,

$$(2.9) \quad \left\{ \int_0^{2\pi} |p(z)|^{\lambda\beta} d\theta \right\}^{1/\lambda} = O(1) \frac{1}{(1 - r)^{\beta-1/\lambda}}.$$

Also, as in the proof of Lemma 2.1, we have (since $\lambda' > 1$)

$$(2.10) \quad \int_E |1 - z|^{-\lambda'} d\theta = O(1) \frac{1}{C^{\lambda'-1}} \frac{1}{(1-r)^{\lambda'-1}}.$$

Hence, combining (2.8), (2.9), and (2.10), we find

$$\left| \int_E (1 - z) f'(z) d\theta \right| = O(1) \frac{1}{C^{1/\lambda}} \frac{1}{(1-r)^\beta},$$

which gives (2.7) if we choose C sufficiently large.

From this point on we proceed essentially as in the proof of [11, Theorem 2], and thus we merely sketch the proof. We define ω_n as in Lemma 2.2, $\lambda_n = \arg \omega_n$, and

$$f'_n(z) = \frac{\omega e^{i\lambda_n}}{(1 - ze^{-i\theta_0})^{\beta+2}} = \omega e^{i\lambda_n} \sum_{m=0}^{\infty} d_m e^{-im\theta_0} z^m.$$

Since $\omega_n = [(1 - r_n)p(r_n e^{i\theta_0})]^\beta$, $\lim_{n \rightarrow \infty} \lambda_n$ exists by [6, Theorem 2]. As in [11, Lemma 3] we find that as $n \rightarrow \infty$,

$$(2.11) \quad a_n - e^{-i\theta_0} a_{n-1} = -\frac{e^{-i\theta_0} a_{n-1}}{n} + \frac{\omega e^{i(\lambda_n - (n-1)\theta_0)}}{\Gamma(\beta+1)} n^{\beta-1} + o(1) n^{\beta-1},$$

and hence as $n \rightarrow \infty$,

$$(2.12) \quad \frac{a_n - e^{-i\theta_0} a_{n-1}}{n^{\beta-1}} = \frac{\omega e^{i(\lambda_n - (n-1)\theta_0)}}{\Gamma(\beta+1)} \left[1 - \frac{1}{\beta+1} (1 + o(1)) \right] + o(1),$$

where we have used (2.11) and Theorem 2.2. Theorem 2.2 also implies that as $n \rightarrow \infty$,

$$\arg e^{-i\theta_0} a_n = \arg \omega e^{i(\lambda_n - n\theta_0)} + o(1),$$

and since $\lim_{n \rightarrow \infty} \lambda_n$ exists we have as $n \rightarrow \infty$ that

$$(2.13) \quad \arg e^{-i\theta_0} a_{n-1} = \arg \omega e^{i(\lambda_n - (n-1)\theta_0)} + o(1).$$

Combining (2.12) with (2.13), we find

$$\frac{||a_n| - |a_{n-1}||}{n^{\beta-1}} = \frac{\beta\omega}{\Gamma(\beta+2)} + o(1)$$

as $n \rightarrow \infty$, which proves the theorem.

We now show that the theorem is false when $\omega = 0$. Let $\beta \geq 0$ be given, and define $f \in K(\beta)$ by

$$f'(z) = \frac{1}{(1 - z^2)^{\beta+1}}.$$

Clearly f is an odd function, and it is easily verified that $a_{2n+1} \sim n^{\beta-1}/2\Gamma(\beta+1)$ as $n \rightarrow \infty$, so

$$\lim_{n \rightarrow \infty} \frac{||a_{2n+1}| - |a_{2n}||}{n^{\beta-1}} = \lim_{n \rightarrow \infty} \frac{|a_{2n+1}|}{n^{\beta-1}} = \frac{1}{2\Gamma(\beta+1)}.$$

However, $\omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} M(r, f') = \lim_{r \rightarrow 1} (1-r)/(1+r)^{\beta+1} = 0$, so the theorem is false when $\omega = 0$. This is in sharp contrast to the corresponding result [11] for V_k , where the result is true for all $k > 2$ even if $\omega = 0$.

3. Behavior of the integral means. In this section we shall investigate the behavior of $I_\lambda(r, f')$ and $I_\lambda(r, f)$, where for $\lambda > 0$ we define

$$I_\lambda(r, g) = \frac{1}{2\pi} \int_0^{2\pi} |g(re^{i\theta})|^\lambda d\theta.$$

Our results again include as special cases previous results of the author [10] for the class V_k as well as generalizing results of Pommerenke [13] for the classes $K(\alpha)$, $0 \leq \alpha \leq 1$. Although the details of the proofs given here are slightly more involved than those for V_k , we refer to [10] whenever possible. We first need two lemmas, the first of which is proved in exactly the same way as [10, Lemma 4.1].

LEMMA 3.1. *Let $f \in K(\beta)$, $\omega = \lim_{r \rightarrow 1} (1-r)^{\beta+2} |f'(re^{i\theta_0})| > 0$. Let $C > 0$ and $\lambda > 0$ be fixed, and for $0 < R < 1$ define $E = \{\theta: C(1-R) \leq |\theta - \theta_0| \leq \pi\}$, $E' = [-\pi, \pi] \setminus E$. Define $\omega(R) = (1-R)^{\beta+2} |f'(Re^{i\theta_0})|$ and*

$$f'_R(z) = \frac{\omega(R)}{(1 - ze^{-i\theta_0})^{\beta+2}}.$$

Then as $R \rightarrow 1$,

$$\int_{E'} |f'_R(Re^{i\theta})|^\lambda d\theta \sim \int_{E'} |f'(Re^{i\theta})|^\lambda d\theta.$$

LEMMA 3.2. *Let $f \in K(\beta)$, $\omega > 0$, and f'_R be as above. If $\lambda(\beta+2) > 1$, then as $r \rightarrow 1$,*

$$I_\lambda(r, f') = I_\lambda(r, f'_r) + o(1)(1-r)^{1-\lambda(\beta+2)}.$$

Proof. By definition, with $z = re^{i\theta}$, we have

$$\begin{aligned} 2\pi |I_\lambda(r, f') - I_\lambda(r, f'_r)| &\leq \int_E |f'(z)|^\lambda d\theta + \int_E |f'_r(z)|^\lambda d\theta \\ &+ \int_{E'} \left\{ |f'(z)|^\lambda - |f'_r(r)|^\lambda \right\} d\theta, \end{aligned}$$

where E and E' are as in Lemma 3.1. If $\beta = 0$, then $\omega > 0$ implies

$f'(z) = (1 - z)^{-2}$, and so the lemma is trivial. With $\beta > 0$, let $\gamma = 1 + 2/\beta$ and $\gamma' = 1 + \beta/2$, so $1/\gamma + 1/\gamma' = 1$. Recalling that in (1.3) we have $\varphi'(z) = (1 - z)^{-2}$ since $\omega > 0$, we have from Hölder's inequality that

$$\int_E |f'(z)|^2 d\theta \leq \left\{ \int_E |p(z)|^{\lambda(\beta+2)} d\theta \right\}^{\beta/(\beta+2)} \left\{ \int_E |1 - z|^{-\lambda(\beta+2)} d\theta \right\}^{2/(\beta+2)}.$$

As in the proof of (2.9) and (2.10) it follows that

$$\int_E |p(z)|^{\lambda(\beta+2)} d\theta = O(1)(1 - r)^{1-\lambda(\beta+2)}.$$

Also, with $\delta > 0$, it follows that

$$\int_E |1 - z|^{-\lambda(\beta+2)} d\theta < \frac{\delta}{(1 - r)^{\lambda(\beta+2)-1}}$$

for $C(\delta)$ depending on δ and for $\lambda(\beta + 2) > 1$, and therefore

$$\int_E |f'(z)|^2 d\theta < \frac{\delta}{(1 - r)^{\lambda(\beta+2)-1}}$$

for r sufficiently close to 1. Clearly this inequality also holds for f'_r , and so using Lemma 3.1 we have for r sufficiently close to 1 that

$$\begin{aligned} 2\pi |I_\lambda(r, f') - I_\lambda(r, f'_r)| &< \frac{2\delta}{(1 - r)^{\lambda(\beta+2)-1}} + o(1) \int_{E'} |f'_r(z)|^2 d\theta \\ &< \frac{2\delta}{(1 - r)^{\lambda(\beta+2)-1}} + \frac{o(1)\omega(r)^\lambda}{(1 - r)^{\lambda(\beta+2)}} \int_0^{(1-r)C(\delta)} d\theta \\ &< \frac{2\delta}{(1 - r)^{\lambda(\beta+2)-1}} + \frac{o(1)\omega(r)^2 C(\delta)}{(1 - r)^{\lambda(\beta+2)-1}}. \end{aligned}$$

Since $\delta > 0$ was arbitrary and since $o(1)$ approaches zero once δ has been fixed, the lemma follows.

We can now determine the asymptotic behavior of $I_\lambda(r, f')$ when $\lambda(\beta + 2) > 1$. For notational convenience, define

$$G(\lambda, \beta) = \frac{\Gamma(\lambda(\beta + 2) - 1)}{2^{\lambda(\beta+2)-1} \Gamma^2\{(\lambda(\beta + 2))/2\}}.$$

THEOREM 3.1. *Let $f \in K(\beta)$ and $\lambda(\beta + 2) > 1$. Then*

$$\lim_{r \rightarrow 1} (1 - r)^{\lambda(\beta+2)-1} I_\lambda(r, f') = \omega^2 G(\lambda, \beta).$$

Proof. If $\omega > 0$, then the theorem is an immediate consequence of Lemma 3.2 and Pommerenke's result [13] that as $r \rightarrow 1$,

$$(3.1) \quad \frac{1}{2\pi} \int_0^{2\pi} |1 + re^{i\theta}|^{-m} d\theta \sim \frac{\Gamma(m-1)}{2^{m-1}\Gamma^2(m/2)} (1-r)^{1-m}$$

whenever $m > 1$. Hence, we now assume $\omega = 0$, and we divide the proof into two cases. We first assume that in (1.3) φ' is not of the form $(1 - ze^{-i\theta})^{-2}$. Then, as is well known, $M(r, \varphi') = O(1)(1-r)^{-\gamma}$ for some $0 < \gamma < 2$. Without loss of generality we assume $\gamma\lambda(\beta+2)/2 > 1$. As in the proof of Lemma 3.2, we find

$$\int_0^{2\pi} |f'(z)|^2 d\theta \leq \left\{ \int_0^{2\pi} |p(z)|^{\lambda(\beta+2)} d\theta \right\}^{\beta/(\beta+2)} \left\{ \int_0^{2\pi} |\varphi'(z)|^{(\lambda(\beta+2))/2} d\theta \right\}^{2/(\beta+2)}$$

and

$$\left\{ \int_0^{2\pi} |p(z)|^{\lambda(\beta+2)} d\theta \right\}^{\beta/(\beta+2)} = O(1)(1-r)^{\beta/(\beta+2)-\lambda\beta}.$$

Also, since φ is convex, $z\varphi'$ is starlike and schlicht, so from [7, Theorem 3.2] we have

$$\left\{ \int_0^{2\pi} |\varphi'(z)|^{(\lambda(\beta+2))/2} d\theta \right\}^{2/(\beta+2)} = O(1)(1-r)^{2/(\beta+2)-\gamma\lambda}.$$

Hence

$$\int_0^{2\pi} |f'(z)|^2 d\theta = O(1)(1-r)^{1-\lambda(\beta+\gamma)},$$

and since $\gamma < 2$ we have as $r \rightarrow 1$

$$(1-r)^{\lambda(\beta+2)-1} I_\lambda(r, f') \longrightarrow 0.$$

It remains only to consider the case $\omega = 0$ and $\varphi'(z) = (1 - ze^{-i\theta_0})^{-2}$ for some θ_0 . Assuming without loss of generality that $\theta_0 = 0$, we find from (1.3) and our hypothesis $\omega = 0$ that

$$0 = \lim_{r \rightarrow 1} (1-r)p(r).$$

As in Lemma 2.2, it now follows that for z in a Stolz angle with vertex at 1, we have $\lim_{|z| \rightarrow 1} (1-z)p(z) = 0$ where the limit is approached uniformly as $|z| \rightarrow 1$. Hence, since $(1-r)|p(z)| \leq |1-z||p(z)|$,

$$|p(z)| \leq \frac{h(r)}{1-r}$$

for z in the Stolz angle, where $h(r) \rightarrow 0$ as $r \rightarrow 1$. Thus, given $C > 0$,

$$(3.2) \quad \begin{aligned} \int_0^{C(1-r)} |f'(z)|^2 d\theta &= \int_0^{C(1-r)} |p(z)|^{2\beta} |1-z|^{-2\lambda} d\theta \\ &\leq \left\{ \int_0^{C(1-r)} |p(z)|^{\lambda(\beta+2)} d\theta \right\}^{\beta/(\beta+2)} \left\{ \int_0^{C(1-r)} |1-z|^{-\lambda(\beta+2)} d\theta \right\}^{2/(\beta+2)} \end{aligned}$$

$$\begin{aligned} &\leq \frac{(Ch(r))^{\beta\lambda}}{(1-r)^{\beta\lambda-\beta/(\beta+2)}} \cdot \frac{O(1)}{(1-r)^{2\lambda-2/(\beta+2)}} \\ &= \frac{o(1)}{(1-r)^{\lambda(\beta+2)-1}} \end{aligned}$$

where we have used (3.1). Exactly as in the proof of Lemma 3.2 we also have, given $\delta > 0$,

$$(3.3) \quad \int_{C(1-r)}^{\pi} |f'(z)|^2 d\theta < \frac{\delta}{(1-r)^{\lambda(\beta+2)-1}}$$

for an appropriate choice of $C = C(\delta)$, and hence from (3.2) and (3.3)

$$\lim_{r \rightarrow 1} (1-r)^{\lambda(\beta+2)-1} I_{\lambda}(r, f') = 0,$$

which completes the proof of Theorem 3.1.

To complete this section, we examine $I_{\lambda}(r, f)$.

THEOREM 3.2. *Let $f \in K(\beta)$ and let $G(\lambda, \beta)$ be as in Theorem 3.1.*

(i) *If $\lambda \geq 1$, then*

$$\liminf_{r \rightarrow 1} (1-r)^{\lambda(\beta+1)-1} I_{\lambda}(r, f) \geq \frac{\omega^{\lambda} G(\lambda, \beta)}{2^{\lambda(\beta+2)-1}}.$$

(ii) *If $\lambda \geq 1$ and $\lambda(\beta+1) > 1$, then*

$$\limsup_{r \rightarrow 1} (1-r)^{\lambda(\beta+1)-1} I_{\lambda}(r, f) \leq \frac{\omega^{\lambda} G(\lambda, \beta)}{(\beta+1 - (1/\lambda))^{\lambda}}.$$

Note that when $\omega = 0$, $\lim_{r \rightarrow 1} (1-r)^{\lambda(\beta+1)-1} I_{\lambda}(r, f) = 0$, and when $\omega > 0$ the growth of $I_{\lambda}(r, f)$ is regular in the sense that $\limsup_{r \rightarrow 1}$ and $\liminf_{r \rightarrow 1}$ are either both positive or both zero.

Proof. The proof of (i) is very similar to that of [10, Theorem 4.4], and so we omit the details. To prove (ii), we first note that

$$f(re^{i\theta}) = \int_0^r f'(te^{i\theta}) dt.$$

Since $\lambda \geq 1$, a generalization of Minkowski's inequality [15, p. 260] gives

$$I_{\lambda}(r, f)^{1/\lambda} \leq \int_0^r I_{\lambda}(t, f')^{1/\lambda} dt.$$

Since Theorem 3.1 gives us the asymptotic behavior of $I_{\lambda}(t, f')$ as $t \rightarrow 1$, a straightforward argument shows that whenever $\lambda(\beta+1) > 1$,

$$\limsup_{r \rightarrow 1} (1 - r)^{\lambda(\beta+1)-1} I_\lambda(r, f) \leq \frac{\omega^\lambda G(\lambda, \beta)}{(\beta + 1 - 1/\lambda)^\lambda}.$$

In conclusion, it should be noted that the basic result underlying the theorems of §§ 2 and 3 is the existence of $\omega = \lim_{r \rightarrow 1} (1 - r)^{\alpha+1} M(r, f')$, where $\alpha = \beta + 1$. Since this limit exists whenever f belongs to a linear-invariant family of order α , it is interesting to speculate as to whether the results of the previous sections remain true if we assume only that f belong to such a linear-invariant family. Nothing seems to be known concerning this question. The similarity between the results of the previous sections and results of Hayman [5] on mean p -valent functions should also be noted. In this direction, W. E. Kirwan has recently shown (unpublished) that given $f \in V_k$ with $2 \leq k \leq 4$, there exists a constant $d(f)$ such that $f - d(f)$ is circumferentially mean- $k/4$ valent.

4. Bazilevic functions and $K(\beta)$. For any $\alpha > 0$, define $B(\alpha)$ to be the class of functions g which are regular in U and which are given by

$$(4.1) \quad g(z) = \left\{ \alpha \int_0^z \xi^{\alpha-1} p(\xi) \left(\frac{h(\xi)}{\xi} \right)^\alpha d\xi \right\}^{1/\alpha},$$

where $p \in \mathcal{P}$, the class of functions P regular in U satisfying $\operatorname{Re} P(z) > 0$ and $P(0) = 1$, and where $h \in \mathcal{S}^*$, the class of normalized starlike functions. The powers appearing in (4.1) are meant as principal values. It is known [1] that $B(\alpha)$ contains only schlicht functions, and it is easy to verify that for various special choices of α , p , and h , the class $B(\alpha)$ reduces to the classes of convex, starlike, and close-to-convex functions. However, in general very little seems to be known about the geometry of $B(\alpha)$. In this section we shall relate $B(\alpha)$ to $K(1/\alpha)$. This relationship will allow us to give a simple geometric interpretation of $B(\alpha)$ as well as a simple geometric proof that $B(\alpha)$ contains only schlicht functions.

We first need a technical lemma.

LEMMA 4.1. *Let g be given by (4.1). Then g is locally schlicht and vanishes only at the origin.*

Proof. If $\alpha = 1$, then it is easily seen that g is close-to-convex, and hence the lemma is trivial. Thus we assume $\alpha \neq 1$. Let $z_0 \neq 0$ be given. We claim that $g(z_0) = 0$ iff $g'(z_0) = 0$. If $g(z_0) \neq 0$, then $(g(z)/z)^\alpha$ is regular in a neighborhood of z_0 , and from (4.1)

$$(4.2) \quad (g(z)/z)^{\alpha-1} g'(z) = p(z)(h(z)/z)^\alpha.$$

Since neither p nor h vanish at z_0 , it then follows that $g'(z_0) \neq 0$.

Suppose now that $g'(z_0) \neq 0$. We must show $g(z) \neq 0$. Since the zeros of g and g' are isolated, it is clear that we may choose (even if $g(z_0) = 0$) an arc γ ending at z_0 such that (4.2) holds for $z \in \gamma$, $z \neq z_0$, and such that $g'(z) \neq 0$ for $z \in \gamma$. Therefore, for $z \in \gamma$,

$$\lim_{z \rightarrow z_0} |g(z)/z|^{\alpha-1} = \left| \frac{p(z_0)}{g'(z_0)} \left(\frac{h(z_0)}{z_0} \right)^\alpha \right|,$$

and hence (since $\alpha \neq 1$) $g(z_0) \neq 0$, which establishes our claim.

To prove the lemma, it is now sufficient to show that g vanishes only at the origin. Suppose not; that is, suppose $g(z) = (z - z_0)^m q(z)$ where $m \geq 1$, $q(z_0) \neq 0$ and $z_0 \neq 0$. We choose an arc γ ending at z_0 such that for $z \in \gamma$ ($z \neq z_0$) we have $g(z) \neq 0$, $g'(z) \neq 0$, and such that (4.2) holds. Then with $z \in \gamma$,

$$(z - z_0)^{m\alpha-1} \left(\frac{q(z)}{z} \right)^{\alpha-1} [(z - z_0)q'(z) + mq(z)] = p(z) \left(\frac{h(z)}{z} \right)^\alpha.$$

We now allow $z \rightarrow z_0$, and we find that $m\alpha = 1$. We now define G for $z \in U$ by $G(z)^m = g(z^m)$. From (4.1) it follows that G is close-to-convex with respect to H , given by $H(z)^m = h(z^m)$ where h is as in (4.1). But $G(z_0^{1/m})^m = g(z_0) = 0$ and $z_0^{1/m} \neq 0$, which contradicts the fact that G is schlicht. This proves the lemma.

We now define $K_0(\beta)$ to be that subclass of $K(\beta)$ such that in (1.3) we have $c = 1$ and $p(0) = 1$. Therefore, $f \in K_0(\beta)$ iff

$$(4.3) \quad f'(z) = p(z)^\beta \frac{h(z)}{z}$$

where $p \in \mathcal{P}$ and $h \in \mathcal{S}^*$. We also assume $\beta > 0$.

THEOREM 4.1. *If $f \in K_0(\beta)$, then $g \in B(1/\beta)$ where*

$$g(z) = \left\{ \frac{1}{\beta} \int_0^z (\xi f'(\xi))^{1/\beta} \xi^{-1} d\xi \right\}^\beta.$$

Conversely, if $g \in B(\alpha)$, then $f \in K_0(1/\alpha)$ where

$$f(z) = \int_0^z \left(\frac{g(\xi)}{\xi} \right)^{1-1/\alpha} (g'(\xi))^{1/\alpha} d\xi.$$

Proof. Suppose first that $f \in K_0(\beta)$ and is given by (4.3). Then

$$f'(z)^{1/\beta} = p(z) \left(\frac{h(z)}{z} \right)^{1/\beta},$$

and from the definition of $B(1/\beta)$ it follows that g defined as in the

theorem belongs to $B(1/\beta)$.

Now we suppose $g \in B(\alpha)$, and we define f as in theorem. By Lemma 4.1 f is regular in U , and since $g \in B(\alpha)$ we have from the definition of f that

$$f'(z)^\alpha = p(z) \left(\frac{h(z)}{z} \right)^\alpha$$

where $p \in \mathcal{P}$ and $h \in \mathcal{S}^*$. Hence $f \in K_0(1/\alpha)$.

Note that although for $\beta > 1$ f may be of arbitrarily high valence, it is always true that the corresponding g is schlicht. Also note that since $V_k \subset K(k/2 - 1)$, we have a relation between V_k and $B(2/(k - 2))$.

We now investigate the geometry of $B(\alpha)$. We shall assume that g is regular and locally schlicht in U , is normalized as in (1.1), and vanishes only at the origin. Also, for $0 < r < 1$, we define the curve $C(r) = \{g(re^{i\theta})^\alpha : 0 \leq \theta < 2\pi\}$.

THEOREM 4.2. *With the above notation and hypothesis on g , we have that $g \in B(\alpha)$ iff for all $0 < r < 1$ the tangent to $C(r)$ never turns back on itself as much as π radians.*

Proof. If $g \in B(\alpha)$, then we see from Theorem 4.1 that $f \in K_0(1/\alpha)$ where

$$f'(z) = \left(\frac{g(z)}{z} \right)^{1-1/\alpha} (g'(z))^{1/\alpha}.$$

Denote by $T(f, re^{i\theta})$ the tangent to the curve $f(|z| = r)$ at $f(re^{i\theta})$. Then with $z = re^{i\theta}$,

$$\arg T(f, re^{i\theta}) = (1 - 1/\alpha) \arg g(z) + (1/\alpha) \arg zg'(z) + \pi/2,$$

from which it follows by a standard argument that

$$\frac{\partial}{\partial \theta} \arg T(f, re^{i\theta}) = (1 - 1/\alpha) \operatorname{Re} \frac{zg'(z)}{g(z)} + \frac{1}{\alpha} \operatorname{Re} \left\{ 1 + \frac{zg''(z)}{g'(z)} \right\}.$$

Since $f \in K_0(1/\alpha)$,

$$\int_{\theta_1}^{\theta_2} \frac{\partial}{\partial \theta} \arg T(f, re^{i\theta}) d\theta > -\pi/\alpha$$

for any $\theta_1 < \theta_2 < \theta_1 + 2\pi$, and so

$$(4.4) \quad (\alpha - 1) \int_{\theta_1}^{\theta_2} \operatorname{Re} \frac{zg'(z)}{g(z)} d\theta + \int_{\theta_1}^{\theta_2} \operatorname{Re} \left(1 + \frac{zg''(z)}{g'(z)} \right) d\theta > -\pi.$$

Noting that locally we have $(g^\alpha(z))' = \alpha g(z)^{\alpha-1} g'(z)$, we see by a standard

argument that (4.4) is equivalent to the fact that the tangent to $C(r)$ never turns back on itself by as much as π radians.

To prove the converse, we have from Lemma 4.1 that for $z \neq 0$, $(g(z))^\alpha$ is locally regular, so we may assume that (4.4) holds. If f is defined by

$$f(z) = \int_0^z \left(\frac{g(\xi)}{\xi} \right)^{1-1/\alpha} (g'(\xi))^{1/\alpha} d\xi,$$

then f is regular in U and from (4.4) we have

$$(4.5) \quad \int_{\theta_1}^{\theta_2} \frac{\partial}{\partial \theta} \arg T(f, re^{i\theta}) d\theta > -\pi/\alpha$$

for any $\theta_1 < \theta_2 < \theta_1 + 2\pi$. Since f' never vanishes, an argument due to Kaplan [9] shows that (4.5) implies $f \in K_0(1/\alpha)$, and thus

$$f'(z) = p(z)^{1/\alpha} \frac{h(z)}{z}$$

where $p \in \mathcal{P}$ and $h \in \mathcal{S}^*$. We now see from the definition of f that

$$g(z) = \left\{ \alpha \int_0^z \xi^{\alpha-1} p(\xi) \left(\frac{h(\xi)}{\xi} \right)^\alpha d\xi \right\}^{1/\alpha},$$


and so $g \in B(\alpha)$. This proves Theorem 4.2.

In conclusion, we prove geometrically that $B(\alpha)$ contains only schlicht functions.

COROLLARY 4.3. *$B(\alpha)$ contains only schlicht functions.*

Proof. Suppose $g \in B(\alpha)$ and g is not schlicht. For each $0 < r < 1$, let $C(r) = \{g(re^{i\theta}) : 0 \leq \theta \leq 2\pi\}$, and let $R = \inf\{r : C(r) \text{ is not a simple curve}\}$. Since $g'(0) = 1$, it is clear that $R > 0$. Also, $R < 1$, since it follows from the argument principle that there exists $r < 1$ such that g is not schlicht on $|z| = r$.

Consider now the curve $C(R)$. Clearly $C(R)$ is nonsimple, and g is schlicht in $\{z : |z| < R\}$. Hence we may choose $w, z_1 = Re^{i\theta_1}$, and $z_2 = Re^{i\theta_2}$ (with $\theta_1 < \theta_2$) such that $g(z_1) = g(z_2) = w$, and such that the curve $C(R)$ is simple for $\theta \in (\theta_1, \theta_2)$.



By Lemma 4.1 g is locally schlicht and vanishes only at the origin, so from Theorem 4.2, with $z = Re^{i\theta}$,

$$(\alpha - 1) \int_{\theta_1}^{\theta_2} d \arg g + \int_{\theta_1}^{\theta_2} d \arg zg'(z) > -\pi .$$

However, by the choice of θ_1 and θ_2 we have $\int_{\theta_1}^{\theta_2} d \arg g = 0$, and so

$$(4.6) \quad \int_{\theta_1}^{\theta_2} d \arg zg' > -\pi .$$

But it is clear geometrically that between θ_1 and θ_2 the argument of the tangent vector to $C(R)$ turns back on itself by π radians, which contradicts (4.6). Therefore g must be schlicht.

Acknowledgement. After completing this paper, the author became aware of the paper [4] by Professor A. W. Goodman. I wish to thank Professor Goodman for providing me with a copy of his manuscript. Aside from the geometrical interpretation of the class $K(\beta)$, the only results appearing both here and in [4] are parts (ii) and (iii) of Theorem 2.3. (See Theorems 8 and 9 of [4].).

REFERENCES

1. I. E. Bazilevič, *On a case of integrability in quadratures of the Loewner-Kufarev equation*, Mat. Sborn., **37** (1955), 471-476. (Russian)
2. D. A. Brannan, *On functions of bounded boundary rotation I*, Proc. Edinburgh Mat. Soc., **16** (1968-69), 339-347.
3. A. W. Goodman, *A note on the Noshiro-Warschawski theorem*, (to appear).
4. ———, *On close-to-convex functions of higher order*, (to appear).
5. W. K. Hayman, *The asymptotic behavior of p -valent functions*, Proc. London Math. Soc., **5** (1955), 257-284.
6. ———, *On functions with positive real part*, J. London Math. Soc., **36** (1961), 35-48.
7. ———, *Multivalent Functions*, (Cambridge, 1958).
8. E. Hille, *Analytic Function Theory, Vol. II*, (Blaisdell, 1962).
9. W. Kaplan, *Close-to-convex schlicht functions*, Michigan Math. J., **1** (1952), 169-185.
10. J. W. Noonan, *Asymptotic behavior of functions with bounded boundary rotation*, Trans. Amer. Math. Soc., **164** (1972), 397-410.
11. ———, and D. K. Thomas, *On successive coefficients of functions of bounded boundary rotation*, J. London Math. Soc., **5** (1972), 656-662.
12. Ch. Pommerenke, *Linear-invariante Familien analytischer Funktionen I*, Math. Annalen, **155** (1964), 108-154.
13. ———, *On the coefficients of close-to-convex functions*, Michigan Math. J., **9** (1962), 259-269.
14. ———, *On close-to-convex analytic functions*, Trans. Amer. Math. Soc., **114** (1964), 176-186.

15. A. Zygmund, *Trigonometric Series Vol. I, II*, 2nd edition (Cambridge, 1968).

Received June 30, 1971. NRC-NRL Post-doctoral Research Associate.

E. O. HULBURT CENTER FOR SPACE RESEARCH
U. S. NAVAL RESEARCH LABORATORY
WASHINGTON, D. C. 20390

Current address: College of the Holy Cross Worcester, Massachusetts 01610

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Pacific Journal of Mathematics

Vol. 44, No. 1

May, 1973

Jimmy T. Arnold, <i>Power series rings over Prüfer domains</i>	1
Maynard G. Arsove, <i>On the behavior of Pincherle basis functions</i>	13
Jan William Auer, <i>Fiber integration in smooth bundles</i>	33
George Bachman, Edward Beckenstein and Lawrence Narici, <i>Function algebras over valued fields</i>	45
Gerald A. Beer, <i>The index of convexity and the visibility function</i>	59
James Robert Boone, <i>A note on mesocompact and sequentially mesocompact spaces</i>	69
Selwyn Ross Caradus, <i>Semiclosed operators</i>	75
John H. E. Cohn, <i>Two primary factor inequalities</i>	81
Mani Gagrat and Somashekhar Amrith Naimpally, <i>Proximity approach to semi-metric and developable spaces</i>	93
John Grant, <i>Automorphisms definable by formulas</i>	107
Walter Kurt Hayman, <i>Differential inequalities and local valency</i>	117
Wolfgang H. Heil, <i>Testing 3-manifolds for projective planes</i>	139
Melvin Hochster and Louis Jackson Ratliff, Jr., <i>Five theorems on Macaulay rings</i>	147
Thomas Benton Hoover, <i>Operator algebras with reducing invariant subspaces</i> ...	173
James Edgar Keesling, <i>Topological groups whose underlying spaces are separable Fréchet manifolds</i>	181
Frank Leroy Knowles, <i>Idempotents in the boundary of a Lie group</i>	191
George Edward Lang, <i>The evaluation map and EHP sequences</i>	201
Everette Lee May, Jr, <i>Localizing the spectrum</i>	211
Frank Belsley Miles, <i>Existence of special K-sets in certain locally compact abelian groups</i>	219
Susan Montgomery, <i>A generalization of a theorem of Jacobson. II</i>	233
T. S. Motzkin and J. L. Walsh, <i>Equilibrium of inverse-distance forces in three-dimensions</i>	241
Arunava Mukherjea and Nicolas A. Tserpes, <i>Invariant measures and the converse of Haar's theorem on semitopological semigroups</i>	251
James Waring Noonan, <i>On close-to-convex functions of order β</i>	263
Donald Steven Passman, <i>The Jacobian of a growth transformation</i>	281
Dean Blackburn Priest, <i>A mean Stieltjes type integral</i>	291
Joe Bill Rhodes, <i>Decomposition of semilattices with applications to topological lattices</i>	299
Claus M. Ringel, <i>Socle conditions for QF - 1 rings</i>	309
Richard Rochberg, <i>Linear maps of the disk algebra</i>	337
Roy W. Ryden, <i>Groups of arithmetic functions under Dirichlet convolution</i>	355
Michael J. Sharpe, <i>A class of operators on excessive functions</i>	361
Erling Stormer, <i>Automorphisms and equivalence in von Neumann algebras</i>	371
Philip C. Tonne, <i>Matrix representations for linear transformations on series analytic in the unit disc</i>	385