

Pacific Journal of Mathematics

**CLOSED RANGE THEOREMS FOR CONVEX SETS AND
LINEAR LIFTINGS**

TSUYOSHI ANDÔ

CLOSED RANGE THEOREMS FOR CONVEX SETS AND LINEAR LIFTINGS

T. ANDO

Let M be a closed subspace of a Banach space E such that its annihilator M^\perp is the range of a projection P . Given a closed convex subset S containing 0 , the first problem of this paper is to find a condition for $\tau(S)$ to be closed where τ is the canonical map from E to E/M . Closure is guaranteed if S is splittable in the sense that the polar S° coincides with the norm-closed convex hull of $P(S^\circ) \cup Q(S^\circ)$, where $Q = 1 - P$. The second problem is to give a condition for existence of a linear map φ , called a linear lifting, from E/M to E such that $\tau \circ \varphi = 1$ and $\varphi \circ \tau(S) \subseteq S$. A linear lifting exists if and only if M is the kernel of a projection making S invariant. Of special interest is the case where S is a ball or a cone. When the unit ball is splittable, existence of a linear lifting of norm one is guaranteed under suitable conditions on E/M , which are satisfied by separable L_p and $C(X)$ on compact metrizable X . If further E is an ordered Banach space, and if both P and Q are positive, M is shown to be the kernel of a positive projection of norm one.

Though the closed range theorem (Theorem 1) yields immediately an abstract version of the Rudin-Carleson-Bishop theorem on norm-preserving extensions of functions defined on a peak set, in §2 further modification (Theorem 2) is shown to include Gamelin's extension [5] of the Rudin-Carleson-Bishop theorem in abstract form. Recently a different approach to generalization of the Gamelin theorem was made by Alfsen and Hirsberg [1]. In §3 it is indicated how the closed range theorem is applied to give unified proofs for results of Davies [4] and Perdrizet [9] on closedness of a cone in a quotient space and on order-preserving extensions. In §4 an idea of Pełczyński-Michael [8] is further developed for the closed range theorem to produce existence of linear liftings under suitable conditions. The Pełczyński-Michael theorems are generalized in abstract form (Theorems 5 and 6).

1. Preliminary. Let E be a real or complex Banach space with unit ball U . E^* and E^{**} are its dual and second dual respectively, and E is always imbedded canonically into E^{**} . x, y, z, \dots are vectors in E or E^{**} while f, g, h, \dots are functionals in E^* . For $x \in E^{**}$ and $f \in E^*$ $f(x)$ is used instead of $x(f)$. The weak topology $\sigma(E^*, E)$ on E^* is called the *weak** topology while $\sigma(E^{**}, E^*)$ on E^{**} is the *weak*** topology. For a subset S of E its norm-closure and its *weak***

closure (in E^{**}) are denoted by \bar{S} and S^\sim respectively.

The *polar* S° is defined as the set of all f such that $\operatorname{Re} f(x) \leq 1$ or $f(x) \leq 1$ on S according as the scalar field is complex or real. When S is a subspace its polar coincides with its *annihilator* S^\perp consisting of all f vanishing on it. The following basic facts are used frequently in this paper. Proofs are found, for instance, in [10]. Let S_1 and S_2 be closed convex subsets of E containing 0. $(S_1 \cap S_2)^\circ$ coincides with the weak* closure of $\operatorname{conv}(S_1^\circ \cup S_2^\circ)$ where $\operatorname{conv}(\cdot)$ denotes the convex hull. The weak** closure S_1^\sim coincides with the polar of S_1° in duality $\langle E^{**}, E^* \rangle$ and $S_1 = E \cap S_1^\sim$. Thus coincidence $(S_1 \cap S_2)^\sim = S_1^\sim \cap S_2^\sim$ occurs if and only if the weak* and the norm closure of $\operatorname{conv}(S_1^\circ \cup S_2^\circ)$ coincide. In some case $\operatorname{conv}(S_1^\circ \cup S_2^\circ)$ becomes itself weak* closed. Here the Krein-Smulian theorem is quite useful: $\operatorname{conv}(S_1^\circ \cup S_2^\circ)$ is weak* closed if (and only if) $\gamma U^\circ \cap \operatorname{conv}(S_1^\circ \cup S_2^\circ)$ is weak* closed for every $0 \leq \gamma < \infty$. If S_1 contains 0 in its interior then S_1° is weak* compact and the norm closure of $\operatorname{conv}(S_1^\circ \cup S_2^\circ)$ is weak* closed. If S_1 is a subspace or a cone, the weak* closure of $\operatorname{conv}(S_1^\circ \cup S_2^\circ)$ is just that of $S_1^\perp + S_2^\circ$. In case both S_1 and S_2 are subspaces, $S_1^\perp + S_2^\perp$ is weak* closed if and only if $S_1 + S_2$ is norm-closed.

Suppose now that E is a real Banach space provided with a closed proper cone E_+ . E_+ gives rise to natural ordering in E under which it becomes the set of all *positive* vectors: $x \leq y$ means $y - x \in E_+$. In this respect E_+ is called the *positive cone*. The dual positive cone E_+^* is defined as the set of f nonnegative on E_+ , or equivalently $E_+^* = -E_+^\circ$. E is called an *ordered Banach space* if $E = E_+ - E_+$ and if there is $\gamma < \infty$ with $(U - E_+) \cap (U + E_+) \subseteq \gamma U$. The latter condition is equivalent to that every subset of the form $\{x; y_1 \leq x \leq y_2\}$ is norm-bounded. For notational convenience the relation $x \leq y + \varepsilon$ in an ordered Banach space means that there is $z \geq 0$ such that $\|z\| < \varepsilon$ and $x \leq y + z$. An ordered Banach space or its norm is called *regular* if $\|x\| = \inf\{\|y\|; -y \leq x \leq y\}$ for every x . A regular norm is monotone on the positive cone in the sense that $0 \leq x \leq y$ implies $\|x\| \leq \|y\|$. An ordered Banach space admits an equivalent regular norm. In fact, the functional $\|x\|_0 = \inf\{\|y\|; -y \leq x \leq y\}$ gives a regular norm.

It is known (cf. [2] and [4]) that E is regular if and only if E^* is regular. An ordered Banach space is said to have the *Riesz interpolation property* if for $y_i \geq x_j$ ($i, j = 1, 2, \dots, n$) there is z such that $x_i \leq z \leq y_i$ ($i = 1, 2, \dots, n$). A regular ordered Banach space is called a *Banach lattice* if it is lattice under the ordering. A Banach lattice has the Riesz interpolation property. It is known (cf. [2] and [4]) that E has the Riesz interpolation property if and only if E^* is a lattice. A continuous linear operator between ordered Banach spaces is called *positive* if it transforms a positive cone into another.

2. Closed range theorems. E is a real or complex Banach space with unit ball U and M is a closed subspace. The canonical map from E to the quotient space E/M is denoted by τ .

Throughout this section it is assumed:

There is a continuous projection P from E^ to M^\perp , and Q stands for $1 - P$.*

Remark that the adjoint Q^* projects E^{**} onto M^\sim but M is not necessarily range of any projection. S, S_1 and S_2 will denote closed convex subsets of E containing 0. S is said to be *splittable*, or more precisely, *P-splittable* if its polar S^0 coincides with the norm-closure of $\text{conv}(P(S^0) \cup Q(S^0))$.

LEMMA 1. *The following conditions are equivalent.*

- (a) S is splittable.
- (b) $S^\sim = \{x \in E^{**}; P^*x \in S^\sim \text{ and } Q^*x \in S^\sim\}$.
- (c) $\theta(f) = \theta(Pf) + \theta(Qf)$ ($f \in E^*$)

where $\theta(f)$ is defined by $\theta(f) = \sup \{\text{Re } f(x); x \in S\}$.

Proof. Since the polar of $P(S^0)$ (resp. of $Q(S^0)$) in E^{**} coincides with the set $\{x \in E^{**}; P^*x$ (resp. $Q^*x \in S^\sim\}$ equivalence of (a) and (b) is clear (cf. §1).

(b) \Rightarrow (c). Obviously $\theta(f)$ can be defined by

$$\theta(f) = \sup \{\text{Re } f(x); x \in S^\sim\}.$$

Take x and y in S . Then by (b) $P^*x + Q^*y$ belongs to S^\sim so that

$$\text{Re } Pf(x) + \text{Re } Qf(y) = \text{Re } f(P^*x + Q^*y) \leq \theta(f),$$

leading to $\theta(Pf) + \theta(Qf) \leq \theta(f)$. The reverse inequality is obvious.

(c) \Rightarrow (b). Since the functional θ is nonnegative because of $S \ni 0$, (c) implies $P(S^0) \cup Q(S^0) \subseteq S^0$. Therefore S^\sim is contained in the set $\{x \in E^{**}; P^*x \in S^\sim \text{ and } Q^*x \in S^\sim\}$. Take x with $P^*x, Q^*x \in S^\sim$. Then by (c)

$$\text{Re } f(x) = \text{Re } Pf(P^*x) + \text{Re } Qf(Q^*x) \leq \theta(Pf) + \theta(Qf) = \theta(f).$$

Thus x belongs to the polar of S^0 in E^{**} .

COROLLARY 1. *The unit ball U is splittable if and only if*

$$\|f\| = \|Pf\| + \|Qf\| \quad (f \in E^*).$$

COROLLARY 2. *If both S_1 and S_2 are splittable and $(S_1 \cap S_2)^\sim = S_1^\sim \cap S_2^\sim$, then $S_1 \cap S_2$ is splittable.*

COROLLARY 3. *A closed subspace (resp. cone) is splittable if and only if its polar is invariant under P (resp. under P and Q).*

Proof. Let N be a closed cone. $P(N^0) \subseteq N^0$ and $Q(N^0) \subseteq N^0$ implies $N^0 = P(N^0) + Q(N^0) = \text{conv}(P(N^0) \cup Q(N^0))$. If N is further a subspace, $Q(N^0) \subseteq N^0$ follows already from $P(N^0) \subseteq N^0$.

LEMMA 2. *If S_1 and S_2 are splittable, for any $\varepsilon > 0$ and $\rho > 0$ the following inclusion relation holds:*

$$\frac{\{S_1 \cap (S_2 + \varepsilon U) + \rho U\} \cap \overline{(S_1 + M)} \cap \overline{(S_2 + M)}}{\subseteq \overline{S_1 \cap (S_2 + \alpha \varepsilon U) + \alpha \rho U \cap M}}$$

where $\alpha = \|Q\|$.

Proof. Take any x in the set on the left hand side. Then it follows by splittability that

$$Q^*x \in S_1^\sim \cap (S_2^\sim + \alpha \varepsilon U^\sim \cap M^\sim) + \alpha \rho U^\sim \cap M^\sim.$$

There is $y \in \alpha \rho U^\sim \cap M^\sim$ such that

$$Q^*(x - y) = Q^*x - y \in S_1^\sim \cap (S_2^\sim + \alpha \varepsilon U^\sim \cap M^\sim)$$

and

$$P^*(x - y) = P^*x \in P^*(\overline{S_1 + M}) \cap P^*(\overline{S_2 + M}) \subseteq S_1^\sim \cap S_2^\sim.$$

Then by Lemma 1 $x - y \in S_1^\sim$ and there is $z \in \alpha \varepsilon U^\sim \cap M^\sim$ such that $x - y - z \in S_2^\sim$. Finally in view of arguments of §1 x belongs to

$$\begin{aligned} & E \cap \{S_1^\sim \cap (S_2^\sim + \alpha \varepsilon U^\sim) + \alpha \rho U^\sim \cap M^\sim\} \\ & \subseteq E \cap \{S_1 \cap (S_2 + \alpha \varepsilon U) + \alpha \rho U \cap M\}^\sim \\ & = \overline{S_1 \cap (S_2 + \alpha \varepsilon U) + \alpha \rho U \cap M}. \end{aligned}$$

By definition of the quotient topology $\tau(x)$ belongs to the closure $\overline{\tau(S)}$ if and only if x is contained in $\overline{S + M}$. In particular, $\tau(S)$ is closed if and only if $S + M$ is closed.

LEMMA 3. *Suppose that S_1 and S_2 are splittable. If $\tau(x)$ belongs to $\overline{\tau(S_1)} \cap \overline{\tau(S_2)}$ and $\|x - S_1 \cap S_2\| < \gamma$ there is $y \in S_1$ such that $\tau(x) = \tau(y)$ and $\|x - y\| < \gamma \|Q\|$. In case $\|Q\| = 1$ for any $\varepsilon > 0$ y can be chosen in $S_1 \cap (S_2 + \varepsilon U)$.*

Proof. Let $\alpha = \|Q\|$ and take ε' with $0 < \varepsilon' < \varepsilon$. By hypothesis x is contained in

$$\{S_1 \cap (S_2 + \varepsilon' U) + \gamma' U\} \cap \overline{(S_1 + M)} \cap \overline{(S_2 + M)}$$

for some $\gamma > \gamma' > 0$. Choose $\varepsilon_n > 0$ such that $\sum_{n=1}^{\infty} \varepsilon_n < \gamma - \gamma'$. By Lemma 2 there is $x_0 \in M$ such that $\|x_0\| \leq \alpha\gamma'$ and $\|x + x_0 - S_1 \cap (S_2 + \alpha\varepsilon'U)\| < \varepsilon_1$. Then

$$x + x_0 \in \overline{S_i + M} + M = \overline{S_i + M}$$

and

$$x + x_0 \in S_1 \cap (S_2 + \alpha\varepsilon'U) + \varepsilon_1 U.$$

Now inductive procedure based on Lemma 2 makes it possible to find a sequence $\{x_n\}$ in M such that $\|x_n\| \leq \alpha\varepsilon_n$ and $\|x + \sum_{i=0}^{n-1} x_i - S_1 \cap (S_2 + \alpha^n\varepsilon'U)\| < \varepsilon_n$. Since $\sum_{n=0}^{\infty} \|x_n\| < \infty$, $y = x + \sum_{n=0}^{\infty} x_n$ is well defined. Obviously y belongs to S_1 , and in case $\alpha = 1$, to $S_1 \cap (\overline{S_2 + \varepsilon'U}) \subseteq S_1 \cap (S_2 + \varepsilon U)$. Finally $\tau(x) = \tau(y)$ and $\|x - y\| \leq \alpha \sum_{n=1}^{\infty} \varepsilon_n + \alpha\gamma' < \alpha\gamma$.

Now the main result of this paper is near at hand.

THEOREM 1. *Suppose that the annihilator of a closed subspace M is the range of a projection P and that S, S_1 and S_2 are closed convex subsets containing 0. Then (a) the image of S under the canonical map τ from E to E/M is closed whenever S is splittable. (b) If both S_1 and S_2 are splittable and $(S_1 \cap S_2)^\sim = S_1^\sim \cap S_2^\sim$ then $\tau(S_1) \cap \tau(S_2) = \tau(S_1 \cap S_2)$. (c) If both S_1 and S_2 are splittable and $\|1 - P\| \leq 1$, then the inclusion $\tau(S_1) \cap \tau(S_2) \subseteq \tau(S_1 \cap (S_2 + \varepsilon U))$ holds for any $\varepsilon > 0$.*

Proof. (a) follows from Lemma 3 with $S_1 = S_2 = S$. Also (c) is a direct consequence of Lemma 3. (b) $S_1 \cap S_2$ is splittable by Corollary 2 and $S_1 \cap S_2 + M$ is closed by (a). Now since by hypothesis

$$\begin{aligned} P^*\{(S_1 + M) \cap (S_2 + M)\} &\subseteq P^*(S_1) \cap P^*(S_2) \subseteq S_1^\sim \cap S_2^\sim \\ &= (S_1 \cap S_2)^\sim, \end{aligned}$$

it follows that

$$(S_1 + M) \cap (S_2 + M) \subseteq E \cap (S_1 \cap S_2 + M)^\sim = S_1 \cap S_2 + M,$$

showing $\tau(S_1) \cap \tau(S_2) \subseteq \tau(S_1 \cap S_2)$. The reverse inclusion is obvious. This completes the proof.

It follows immediately from Theorem 1 that if the unit ball U is splittable and if N is a closed splittable subspace then $\tau(N \cap U)$ is closed and coincides with $\tau(N) \cap \tau(U)$. Let us show that the same conclusion holds for a non-splittable subspace under suitable conditions.

Since by Corollary 3 splittability of N is characterized by $P(N^\perp) \subseteq N^\perp \cap M^\perp$, it follows by Corollary 1 that under the splittability of U , N is splittable if and only if

$$\|Qf\| = \|f - M^\perp \cap N^\perp\| \quad (f \in N^\perp).$$

On the other hand, Corollary 1 implies $\|Qf\| = \|f - M^\perp\|$. Thus if the unit ball is splittable, splittability of N is characterized by

$$\|f - M^\perp \cap N^\perp\| \leq \|f - M^\perp\| \quad (f \in N^\perp).$$

LEMMA 4. *Let N_1 and N_2 be closed subspace. Then for $\rho > 0$ the following assertions are equivalent.*

- (a) $\|x - N_1 \cap N_2\| \leq \rho \|x - N_1\| \quad (x \in N_2)$
- (b) $\|f - N_1^\perp \cap N_2^\perp\| \leq \rho \|f - N_2^\perp\| \quad (f \in N_1^\perp)$

Proof. (a) means that

$$N_2 \cap (\overline{N_1 + U}) \subseteq \overline{N_1 \cap N_2 + \rho U}$$

which implies by polar formation

$$(N_1^\perp + N_2^\perp) \cap U^0 \subseteq N_2^\perp + N_1^\perp \cap \rho U^0.$$

The last relation can be converted to

$$N_1 \cap (N_2^\perp + U^0) \subseteq N_1^\perp \cap N_2^\perp + \rho U^0$$

which is nothing but (b). The reverse process can be pursued because (b) implies that $(N_1^\perp + N_2^\perp) \cap U^0$ is weak* compact, hence by the Krein-Smulian theorem that $N_1^\perp + N_2^\perp$ is weak* closed.

COROLLARY 4. *Suppose that the unit ball is splittable. Then the following assertions for a closed subspace N are equivalent.*

- (a) N is splittable.
- (b) $\|x - N \cap M\| \leq \|x - N\| \quad (x \in M)$
- (c) $\|f - N^\perp \cap M^\perp\| \leq \|f - M^\perp\| \quad (f \in N^\perp)$.

THEOREM 2. *Let M and N be closed subspaces, and suppose that the annihilator M^\perp is the range of a projection P such that*

$$(1) \quad \|f\| = \|Pf\| + \|f - Pf\| \quad (f \in E^*).$$

If for some $1 \leq \rho < 2$

$$(2) \quad \|x - N \cap M\| \leq \rho \|x - N\| \quad (x \in M),$$

then the images of the unit ball U , N and $N \cap U$ under the canonical map τ from E to E/M are closed and

$$\tau(N \cap U) = \tau(N) \cap \tau(U).$$

Proof. Closedness of $\tau(U)$ follows from (1) by Corollary 1 and

Theorem 1. Let $Q = 1 - P$. Then in view of (1) relation (2) is converted by Lemma 4 to

$$(3) \quad \|Pf - M^\perp \cap N^\perp\| \leq \gamma \|Qf\| \quad (f \in N^\perp)$$

where $\gamma = \rho - 1 < 1$. Then for any $f \in N^\perp$ and $g \in M^\perp$.

$$\begin{aligned} \|g - M^\perp \cap N^\perp\| &\leq \|Pf - M^\perp \cap N^\perp\| + \|g - Pf\| \\ &\leq \gamma \|Qf\| + \|g - Pf\| \leq \|g - f\|, \end{aligned}$$

showing

$$\|g - M^\perp \cap N^\perp\| \leq \|g - N^\perp\| \quad (g \in M^\perp),$$

which is converted by Lemma 4 to

$$(4) \quad \|x - M \cap N\| \leq \|x - M\| \quad (x \in N).$$

This last relation means that the canonical map from the Banach space $N/M \cap N$ onto $\tau(N)$ has bounded inverse. Therefore $\tau(N)$ is closed. Further (4) implies

$$N \cap (U + M) \subseteq \overline{U + N \cap M}.$$

Let us prove that really

$$N \cap (U + M) \subseteq U + N \cap M$$

holds, which is equivalent to the required relation:

$$\tau(N) \cap \tau(U) \subseteq \tau(N \cap U).$$

Suppose for contradiction that there exists x in $N \cap (U + M)$ with $(U - x) \cap N \cap M = \emptyset$. Since $N \cap M$ is a subspace, it follows that

$$\text{conv}((U - x) \cup \{0\}) \cap N \cap M = \{0\}.$$

Since $\text{conv}((U - x) \cup \{0\})$ is closed, the last relation implies by polar formation that $(U - x)^0 + N^\perp + M^\perp$ is weak* dense in E^* . Weak* closedness of $(U - x)^0 + N^\perp + M^\perp$, if proved, leads to

$$\text{conv}((U^\sim - x) \cup \{0\}) \cap N^\sim \cap M^\sim = \{0\}$$

hence to a contradiction:

$$x \notin E \cap (U^\sim + N^\sim \cap M^\sim) = \overline{U + N \cap M}.$$

Now let us prove the above weak* closedness. To this end, in view of the Krein-Smulian theorem, it suffices to prove that for any $n > 0$

$$\{(U - x)^0 + N^\perp + M^\perp\} \cap nU^0 \subseteq \delta U^0 \cap (U - x)^0 + \delta U^0 \cap N^\perp + M^\perp$$

where δ is a constant depending on n . Remark that $(U - x)^0$ consists of all f with $\|f\| \leq \operatorname{Re} f(x) + 1$. Since $x \in U + M$ implies $\|P^*x\| \leq 1$, it follows from (1) that

$$\begin{aligned} & 1 + \operatorname{Re} f(x) - \|f\| \\ & \leq 1 + \operatorname{Re} Qf(x) - \|Qf\| \\ & \quad - \{\|Pf\| \|P^*x\| - |Pf(P^*x)|\} \\ & \leq 1 + \operatorname{Re} Qf(x) - \|Qf\|. \end{aligned}$$

This indicates that Q makes $(U - x)^0$ invariant. Now take $f \in (U - x)^0$, $g \in N^\perp$ and $h \in M^\perp$ with $\|f + g + h\| \leq n$. Then by (1) $\|Qf + Qg\| \leq n$. Since $x \in N \cap (U + M)$ and $g \in N^\perp$,

$$\begin{aligned} \operatorname{Re} Qf(x) & \leq n\|x\| - \operatorname{Re} Qg(x) = n\|x\| + \operatorname{Re} Pg(x) \\ & \leq n\|x\| + \|Pg - N^\perp \cap M^\perp\| \|x - M\| \\ & \leq n\|x\| + \|Pg - N^\perp \cap M^\perp\|. \end{aligned}$$

Since Qf belongs to $(U - x)^0$ as f , it follows that

$$\begin{aligned} \|Qg\| & \leq n + \|Qf\| \leq n + \operatorname{Re} Qf(x) + 1 \\ & \leq n(\|x\| + 2) + \|Pg - N^\perp \cap M^\perp\|. \end{aligned}$$

Then (3) applied to g yields

$$\|Pg - N^\perp \cap M^\perp\| \leq \frac{\gamma(n\|x\| + 2)}{1 - \gamma} \equiv \delta_1$$

and consequently

$$\|Qf + g - N^\perp \cap M^\perp\| \leq n + \delta_1 \equiv \delta_2.$$

Since $x \in N$, $g \in N^\perp$ and $Qf \in (U - x)^0$,

$$\|Qf\| \leq \operatorname{Re} Qf(x) + 1 \leq \delta_2\|x\| + 1 \equiv \delta_3$$

and

$$\|g - N^\perp \cap M^\perp\| \leq \delta_2 + \delta_3 \equiv \delta.$$

This implies that

$$\begin{aligned} f + g + h & = Qf + g + (Pf + h) \\ & \equiv \delta U^0 \cap (U - x)^0 + \delta U^0 \cap N^\perp + M^\perp. \end{aligned}$$

This completes the proof.

Consider the sup-norm Banach space $C(X)$ of continuous functions on a compact Hausdorff space X . By the Riesz theorem its dual is realized by the space of regular Borel measures on X with total-variation norm. Given a closed subset Y of X , let M be the subspace

of functions in $C(X)$ vanishing on Y . Then M^\perp is the set of measures with support in Y and becomes the range of a projection $P: Pm = \chi m$ for each measure m where χ is the characteristic function of Y . Obviously (1) is satisfied. Now let N be a closed subspace of $C(X)$. As shown in [5] (3) is equivalent to the property that for any $x \in N$ with $|x(t)| < 1$ ($t \in Y$) and any closed subset $Z \subset X$ with $Y \cap Z = \emptyset$ there is $y \in N$ such that $x(t) = y(t)$ ($t \in Y$), $|y(s)| < \gamma$ ($s \in Z$) and $\|y\| < \max(1, \gamma)$. Remark that $\|x - M\|$ coincides with the norm of the restriction $x|_Y$ of x to Y and that $x(t) = y(t)$ ($t \in Y$) is equivalent to $x - y \in M$. Thus Theorem 2 shows that if (3) with $\gamma < 1$, or equivalently (2) with $\rho < 2$, is satisfied then for any $x \in N$ there is $y \in N$ such that $x|_Y = y|_Y$ and $\|y\| = \|x|_Y\|$. The case $\gamma = 0$ is the generalized Carleson-Rudin theorem (cf. [6] Chap. II). As Gamelin [5] shows, Theorem 2 can further yield the following: suppose that (3) with $\gamma < 1$, or equivalently (2) with $\rho < 2$, is satisfied and that $p \in C(X)$ satisfies $p(t) = 1$ ($t \in Y$) and $p(s) > \gamma$ ($s \in X$). Then if $x \in N$ satisfies $|x(t)| \leq p(t)$ ($t \in Y$) there is $y \in N$ such that $x(t) = y(t)$ ($t \in Y$) and $|y(s)| \leq p(s)$ ($s \in X$). The case $\gamma = 0$ is the Bishop theorem (cf. [6] Chap. II). Generalization of the Gamelin theorem to other direction is treated by Alfsen and Hirsberg [1].

3. Ordered Banach spaces. Let E be an ordered Banach space with positive cone E_+ . A closed subspace M is called an *ideal* if $(M - E_+) \cap (M + E_+) \subseteq M$. An ideal M is *hypostrict* if its annihilator M^\perp is the range of a projection P such that $f \geq Pf \geq 0$ for every $f \geq 0$. The requirement means that both P and $Q = 1 - P$ are positive. Perdrizet [9] shows that a closed subspace M is a hypostrict ideal if and only if the following two conditions are satisfied: (1) Given $x_1, x_2 \in M$ and $y \in E$ with $x_1, x_2 \leq y$, for any $\varepsilon > 0$ there is $z \in M$ such that $x_1, x_2 \leq z \leq y + \varepsilon$, and (2) given $x \in M$ and $y_1, y_2 \in E_+$ with $x \leq y_1 + y_2$ there are $x_1, x_2 \in M$ such that $x = x_1 + x_2$ and $x_i \leq y_i + \varepsilon$ $i = 1, 2$. Under the Riesz interpolation property an ideal M is hypostrict if and only if it is positively generated in the sense: $M = M \cap E_+$.

When M is an ideal, the Banach space E/M is preordered by the cone $\tau(E_+)$ where τ is the canonical map from E to E/M . The following theorem was first proved by Davies [4] under the Riesz interpolation property and then by Perdrizet [9] in general case. Let us give a proof based on Theorem 1.

THEOREM 3. *Let E be an ordered Banach space with positive cone E_+ . If M is a hypostrict ideal then E/M is an ordered Banach space with $\tau(E_+)$ as its positive cone. If E is regular in addition, so is E/M .*

Proof. Since hypostriictness means that $E_+^* = -E_+^0$ is invariant under both P and Q , E_+ is splittable by Corollary 3. Then $\tau(E_+)$ is closed by Theorem 1. M^\perp is isometric to the dual of E/M , and the dual positive cone is identified with $M^\perp \cap E_+^*$. Suppose that E is regular. Then E^* is regular. Since P is positive and is of norm one in this case, M^\perp is a regular ordered Banach space with $M^\perp \cap E_+^*$ as its positive cone. Therefore E/M is regular as stated in §1. This completes the proof because every ordered Banach space admits an equivalent regular norm.

COROLLARY 5. *Suppose that the positive cone E_+ has nonempty interior and that M is a hypostriict ideal. If a closed subspace N is splittable and if it contains an interior point of E_+ then $\tau(N \cap E_+)$ is closed and $\tau(N \cap E_+) = \tau(N) \cap \tau(E_+)$.*

Proof. Since E_+ is splittable, in view of Corollary 2 and Theorem 1 it suffices to prove that $(N \cap E_+)^0 = N^\perp + E_+^0$. Remark that f belongs to $(N \cap E_+)^0$ if and only if the restriction of $-f$ to N is positive. However it is known (cf. [10] Chap. V §5) that when N contains an interior point of E_+ every continuous positive linear functional on N admits a continuous positive linear extension to E , in other words, $-(N \cap E_+)^0 = -(N^\perp + E_+^0)$.

Since E/M is ordered by the cone $\tau(E_+)$, for any y, z with $\tau(z) \leq \tau(y)$ there is y' such that $z \leq y'$ and $\tau(y) = \tau(y')$. The next task is to treat the case $\tau(z) \leq \tau(y) \leq \tau(x)$ and $z \leq x$ and to find a condition of existence y'' such that $z \leq y'' \leq x$ and $\tau(y) = \tau(y'')$.

LEMMA 5. *Let S_1 and S_2 be closed convex subsets containing 0. If for any $0 < \lambda < 1$, $f \in S_1^0$ and $g \in S_2^0$ there are $f' \in S_1^0$ and $g' \in S_2^0$ such that*

$$\lambda f + (1 - \lambda)g = \lambda f' + (1 - \lambda)g'$$

and

$$\lambda \|f'\|, (1 - \lambda)\|g'\| \leq \|\lambda f + (1 - \lambda)g\|$$

*then $(S_1 \cap S_2)^\sim$ coincides with $S_1^\sim \cap S_2^\sim$ where $(\cdot)^\sim$ denotes the weak** closure.*

Proof. In view of the Krein-Smulian theorem it suffices to prove that for any $\gamma > 0$ the weak* closure of $\text{conv}(S_1^0 \cup S_2^0) \cap \gamma U^0$ is contained in the norm closure of $\text{conv}(S_1^0 \cup S_2^0)$. Suppose that $0 < \lambda_\alpha < 1$, $f_\alpha \in S_1^0$, $g_\alpha \in S_2^0$ and $\|\lambda_\alpha f_\alpha + (1 - \lambda_\alpha)g_\alpha\| \leq \gamma$ and that the net $\{\lambda_\alpha f_\alpha +$

$(1 - \lambda_\alpha)g_\alpha\}$ weak* converges to h and the net $\{\lambda_\alpha\}$ converges to λ . By hypothesis $\{\lambda_\alpha f_\alpha\}$ and $\{(1 - \lambda_\alpha)g_\alpha\}$ can be assumed to be bounded, hence to weak* converge to f' and g' respectively. If $0 < \lambda < 1$, $\{f_\alpha\}$ and $\{g_\alpha\}$ can be assumed to weak* converge to $f'' \in S_1^0$ and $g'' \in S_2^0$ respectively. Then $h = \lambda f'' + (1 - \lambda)g''$ belongs to $\text{conv}(S_1^0 \cup S_2^0)$. In case $\lambda = 0$, $h = f' + g''$ and nf' belongs to S_1^0 for any $n > 0$. Therefore h , as the norm limit of $1/n(nf') + (1 - 1/n)g''$, belongs to the norm closure of $\text{conv}(S_1^0 \cup S_2^0)$. The case $\lambda = 1$ is treated similarly.

COROLLARY 6. $(\bigcap_{i=1}^n (x_i - E_+))^\sim = \bigcap_{i=1}^n (x_i - E_+)^\sim$ whenever $x_i \geq 0$ $i = 1, 2, \dots, n$.

Proof. E , hence E^* , can be assumed to be regular. $(x_i - E_+)^0$ consists of all $0 \leq f$ with $f(x_i) \leq 1$. Suppose that $\lambda_i \geq 0$, $\sum_{i=1}^n \lambda_i = 1$ and $f_i \in (x_i - E_+)^0$. Since the norm is monotone on the dual positive cone by regularity, it follows that $\lambda_j \|f_j\| \leq \|\sum_{i=1}^n \lambda_i f_i\|$ $j = 1, 2, \dots, n$. Now inductive application of Lemma 5 yields the assertion.

The following theorem was proved by Perdrizet [9]. Let us give a proof based on Theorem 1.

THEOREM 4. *Let E be an ordered Banach space with positive cone E_+ . Suppose that M is a hypostrict ideal and E/M is ordered by the cone $\tau(E_+)$ where τ is the canonical map from E to E/M . If $z_i \leq 0 \leq x_i$ and $\tau(z_i) \leq \tau(y) \leq \tau(x_i)$ $i = 1, 2, \dots, n$, then for any $\varepsilon > 0$ there is y' such that $z_i \leq y' \leq x_i + \varepsilon$ $i = 1, 2, \dots, n$ and $\tau(y) = \tau(y')$. Further ε can be made 0 if every x_i is an interior point of E_+ or if E has the Riesz interpolation property.*

Proof. E is assumed to be regular, hence Q is of norm one. $z_i + E_+$ is a closed convex set containing 0. It is splittable because both P^* and Q^* are positive and z_i is negative. Similarly $x_i - E_+$ is splittable. Let $S_1 = \bigcap_{i=1}^n (z_i + E_+)$ and $S_2 = \bigcap_{i=1}^n (x_i - E_+)$. Then by Corollaries 2, 6, and Theorem 1 both S_1 and S_2 are splittable and

$$\begin{aligned} & \bigcap_{i=1}^n \tau(z_i + E_+) \cap \bigcap_{i=1}^n \tau(x_i - E_+) \\ &= \tau(S_1) \cap \tau(S_2) \subseteq \tau(S_1 \cap (S_2 + \varepsilon U)) \end{aligned}$$

which is just the first assertion.

If every x_i is an interior point of E_+ , S_2 contains 0 in its interior and by Corollary 2 and Theorem 1

$$\tau(S_1) \cap \tau(S_2) = \tau(S_1 \cap S_2) .$$

Suppose finally that E has the Riesz interpolation property. Since E^{**} becomes a lattice as stated in §1, S_1^\sim consists of all $w \in E^{**}$ with $\bigvee_{i=1}^n z_i \leq w$, where $\bigvee_{i=1}^n z_i$ denotes the supremum of z_1, \dots, z_n in E^{**} . Then S_1^0 consists of all $0 \leq f$ with $f(\bigvee_{i=1}^n z_i) \leq 1$. Similarly S_2^0 consists of all $0 \leq g$ with $g(\bigwedge_{i=1}^n x_i) \leq 1$ where $\bigwedge_{i=1}^n x_i$ denotes the infimum of x_1, \dots, x_n , in E^{**} . Take $0 < \lambda < 1$, $f \in S_1^0$ and $g \in S_2^0$ and let $h = \lambda f + (1 - \lambda)g$. Since E^* is a Banach lattice as stated in §1, and since both $-f$ and g are positive, it follows that $0 \geq h \wedge 0 \geq \lambda f$ and $0 \leq h \vee 0 \leq (1 - \lambda)g$. Let $f' = (1/\lambda)(h \wedge 0)$ and $g' = (1)/(1 - \lambda)h \vee 0$. Then it follows from the above characterization of S_i^0 that $f' \in S_1^0$, $g' \in S_2^0$ and $h = \lambda f' + (1 - \lambda)g'$. Now since $\|h \wedge 0\|, \|h \vee 0\| \leq \|h\|$, Lemma 5 yields $(S_1 \cap S_2)^\sim = S_1^\sim \cap S_2^\sim$ and the assertion follows from Theorem 1.

4. Linear lifting. Let E be a Banach space with unit ball U and M a closed subspace. The canonical map from E to E/M is denoted by τ . A continuous linear map φ from E/M to E is called a *linear lifting* if $\tau \circ \varphi = 1$. If φ is a linear lifting, $\varphi \circ \tau$ is a projection with M as its kernel. Conversely, a linear lifting exists if M is the kernel of a continuous projection.

In this section it is assumed:

There is a projection P from E^ to M^\perp such that*

$$\|f\| = \|Pf\| + \|f - Pf\| \quad (f \in E^*)$$

and Q stands for $1 - P$.

Let F be a finite dimensional Banach space with unit ball V . Consider the dual system $\langle F^* \otimes E, F \otimes E^* \rangle$ of tensor products. When $F^* \otimes E$ is provided with the Minkowski functional of $(V \otimes U^0)^0$ as norm, it is called the *inductive* tensor product of F^* and E and is denoted by $F^* \check{\otimes} E$. When $F \otimes E^*$ is provided with the Minkowski functional of $\text{conv}(V \otimes U^0)$ as norm, it is called the *projective* tensor product of F and E^* and is denoted by $F \hat{\otimes} E^*$. Let $B = B(F, E)$ denote the Banach space of all continuous linear maps from F to E , provided with operator-norm. Since F is finite dimensional, B is isometric to the inductive tensor product $F^* \check{\otimes} E$ under the canonical correspondence. The following lemma, whose proof is found in [10] Chap. IV §9, is basic in the subsequent development.

LEMMA 6. *The dual of $B(F, E)$ is isometric to the projective tensor product $F \hat{\otimes} E^*$ while the second dual is isometric to the inductive tensor product $F^* \check{\otimes} E^{**}$, hence to $B(F, E^{**})$.*

In view of Lemma 6 B^{**} is always identified with $B(F, E^{**})$.

In this case the imbedding of B to B^{**} is just the natural imbedding of $B(F, E)$ to $B(F, E^{**})$. In accordance with the terminology in §1 the weak^{**} closure of a subset G of B is formed in $B(F, E^{**})$ and is denoted by G^\sim . When K and S are a subset of F and a closed convex subset of E containing 0 respectively, $G(K, S)$ and $\mathcal{G}(K, S^\sim)$ denote the set of all $\varphi \in B$ with $\varphi(K) \subseteq S$ and the set of all $\psi \in B^{**}$ with $\psi(K) \subseteq S^\sim$. Obviously $G(K, S)$ is a closed convex subset of B containing 0 and its weak^{**} closure is contained in $\mathcal{G}(K, S^\sim)$.

COROLLARY 6. (a) $G(V, U)^\sim = \mathcal{G}(V, U^\sim)$. (b) $\{G(H, 0) \cap G(F, N)\}^\sim = \mathcal{G}(H, 0) \cap \mathcal{G}(F, N^\sim)$ if H and N are closed subspaces of F and E respectively. (c) $G(K, S)^\sim = \mathcal{G}(K, S^\sim)$ if K is a cone generated by a linearly independent basis $\{x_1, \dots, x_n\}$ of F and S is a cone.

Proof. (a) is an immediate consequence of Lemma 6. (b) $G(F, N)^\sim = \mathcal{G}(F, N^\sim)$ follows from Lemma 6 applied to N instead of E . Since F is finite dimensional, H is the kernel of a projection σ . Then

$$\begin{aligned} \mathcal{G}(H, 0) \cap \mathcal{G}(F, N^\sim) &= \mathcal{G}(F, N^\sim) \circ \sigma = G(F, N)^\sim \circ \sigma \\ &= \{G(F, N) \circ \sigma\}^\sim = \{G(H, 0) \cap G(F, N)\}^\sim, \end{aligned}$$

while the reverse inclusion is obvious. (c) Take any φ in $\mathcal{G}(K, S^\sim)$ and let $y_i = \varphi(x_i)$ $i = 1, 2, \dots, n$. Since each y_i belongs to S^\sim , there are nets $\{y_{i,\alpha}\}$ in S , weak^{**} converging to y_i $i = 1, 2, \dots, n$. Consider a net $\{\varphi_\alpha\}$ in B defined by $\varphi_\alpha(x_i) = y_{i,\alpha}$ $i = 1, 2, \dots, n$. By hypothesis it is contained in $G(K, S)$ and weak^{**} converges to φ . Thus $\mathcal{G}(K, S^\sim)$ is contained in $G(K, S)^\sim$ with the reverse inclusion is obvious.

Since B^* is identified with the projective tensor product $F \hat{\otimes} E^*$ by Lemma 6, the operators $1 \otimes P$ and $1 \otimes Q$ are considered to define projections in B^* . The adjoints of $1 \otimes P$ and $1 \otimes Q$ are realized in $B(F, E^{**})$ according to the following formula:

$$(5) \quad (1 \otimes P)^* \varphi = P^* \circ \varphi \text{ and } (1 \otimes Q)^* \varphi = Q^* \circ \varphi \quad (\varphi \in B(F, E^{**})) .$$

LEMMA 7. *The annihilator $G(F, M)^\perp$ is the range of $1 \otimes P$.*

Proof. Since Q^* is a projection onto M^\sim , by (5) $(1 \otimes Q)^*$ projects B^{**} onto $\mathcal{G}(F, M^\sim)$, which coincides with $G(F, M)^\sim$ by Corollary 6. Then $1 \otimes P$ is obviously a projection from B^* to $G(F, M)^\perp$.

On the basis of Lemma 7, a sentence “ $G(K, S)$ is *splittable*” will always mean that $G(K, S)$ is $1 \otimes P$ -splittable.

COROLLARY 7. *If S is splittable and $G(K, S)^\sim = \mathcal{G}(K, S^\sim)$ then*

$G(K, S)$ is splittable.

Proof. This follows from (5) by Lemma 1.

The following lemma can be considered a development of a basic device in Michael and Pełczyński [8], treating linear lifting in a special case. The crucial requirement for P plays a decisive role in the proof.

LEMMA 8. Suppose that S is splittable and $G(K, S)^\sim = \mathcal{G}(K, S^\sim)$ for a subset K of F . If ψ belongs to

$$G(\pi(K), S) \cap G(\pi(V), U) \cap G(K, S + M) \cap G(V, U + M)$$

where π is a projection of F to a subspace H , then for any $\varepsilon > 0$ there is φ in $G(K, S) \cap G(V, U)$ such that

$$\tau \circ \varphi = \tau \circ \psi \quad \text{and} \quad \|(\varphi - \psi)|H\| < \varepsilon.$$

Proof. Remark first of all that the requirement for P means by Corollary 1 that the unit ball U is splittable.

Let $\psi_1 = \psi - Q^* \circ \psi \circ (1 - \pi)$. Since

$$Q^* \circ \psi_1(K) \subseteq Q^* \circ \psi \circ \pi(K) \subseteq Q^*(S) \subseteq S^\sim$$

and

$$P^* \circ \psi_1(K) \subseteq P^* \circ \psi(K) \subseteq P^*(S + M) \subseteq S^\sim$$

by splittability of S , ψ_1 belongs to $\mathcal{G}(K, S^\sim)$ by Lemma 1, hence to $G(K, S)^\sim$ by hypothesis. Since U is splittable and $G(V, U)^\sim = \mathcal{G}(V, U^\sim)$ by Corollary 6, the same argument shows that ψ_1 belongs also to $G(V, U)^\sim$. Moreover it belongs to $\{G(K, S) \cap G(V, U)\}^\sim$ because $G(V, U)$ is the unit ball of B . On the other hand, $Q^* \circ \psi \circ (1 - \pi)$ belongs to $\mathcal{G}(H, 0) \cap \mathcal{G}(F, M^\sim)$, hence to $\{G(H, 0) \cap G(F, M)\}^\sim$ by Corollary 6. Thus ψ belongs to

$$\{G(K, S) \cap G(V, U) + G(H, 0) \cap G(F, M)\}^\sim.$$

It follows that ψ must be contained in the norm closure of

$$G(K, S) \cap G(V, U) + G(H, 0) \cap G(F, M).$$

Therefore there is $\psi_2 \in B$ such that $\psi - \psi_2 \in G(H, 0) \cap G(F, M)$ and

$$\|\psi_2 - G(K, S) \cap G(V, U)\| < \varepsilon.$$

Since $G(K, S) \cap G(V, U)$ is splittable by hypothesis and Corollary 7, Lemma 3 guarantees that there is $\varphi \in G(K, S) \cap G(V, U)$ such that $\varphi - \psi_2 \in G(F, M)$ and $\|\varphi - \psi_2\| < \varepsilon$. Now $\psi_2 - \psi \in G(H, 0) \cap G(F, M)$

implies that $\tau \circ \varphi = \tau \circ \psi$ and

$$\|(\varphi - \psi)|H\| = \|(\varphi - \psi_2)|H\| \leq \|\varphi - \psi_2\| < \varepsilon.$$

Let S be a closed splittable subset of E and L a subset of $\tau(S)$. Suppose that there is a sequence of projections $\{\pi_n\}$ in E/M such that (1) the range F_n of π_n is of finite dimension, (2) $\|\pi_n\| \leq 1$, (3) $\pi_n \cdot \pi_m = \pi_n$ for $n \leq m$, (4) $\pi_n(L) \subseteq L$ and (5) π_n converges strongly to the identity as $n \rightarrow \infty$.

Let \mathcal{G}_n denote the set of all $\varphi \in B(F_n, E)$ with $\varphi \circ \pi_n(L) \subseteq S$ while G_n is the set of all $\psi \in B(F_n, E^{**})$ with $\psi \circ \pi_n(L) \subseteq S^\sim$. As before, the second dual of $B(F_n, E)$ is identified with $B(F_n, E^{**})$.

LEMMA 9. *If the weak** closure of G_n coincides with \mathcal{G}_n , $n = 1, 2, \dots$, then there is a linear lifting φ from E/M to E such that $\varphi(L) \subseteq S$ and $\|\varphi\| \leq 1$.*

Proof. Let $\pi_0 = 0$ and $\varphi_0 = 0$. Assume that linear maps $\varphi_j \in B(F_j, E)$ $j = 0, 1, \dots, n$ have been found in such a way that $\tau \circ \varphi_j = 1$ on F_j , $\|\varphi_j\| \leq 1$, $\varphi_j \circ \pi_j(L) \subseteq S$ and $\|(\varphi_{j-1} - \varphi_j)|F_{j-1}\| < 1/2^{j-1}$ $j = 0, 1, \dots, n$. Since F_{n+1} is finite dimensional by hypothesis, there is $\psi \in B(F_{n+1}, E)$ such that $\tau \circ \psi = 1$ on F_{n+1} . Consider the map $\psi' = \varphi_n \circ \pi_n + \psi \circ (1 - \pi_n)$ from F_{n+1} to E . Then by assumption

$$\psi' \circ \pi_n(\pi_{n+1}(L)) = \varphi_n \circ \pi_n(L) \subseteq S$$

and in view of $\|\pi_n\| \leq 1$

$$\psi' \circ \pi_n(V_{n+1}) = \varphi_n(V_n) \subseteq U$$

where V_i denotes the unit ball of F_i . Since $V_{n+1} \subseteq \tau(U)$ by Theorem 1 and $\pi_{n+1}(L) \subseteq \tau(S)$,

$$\psi'(V_{n+1}) \subseteq U + M \text{ and } \psi'(\pi_{n+1}(L)) \subseteq S + M.$$

Since the weak** closure of G_n coincides with \mathcal{G}_n by hypothesis, Lemma 8, applied to F_{n+1} , $\pi_{n+1}(L)$ and π_n instead of F , K and π , yields that there is $\varphi_{n+1} \in G_{n+1}$ such that $\|\varphi_{n+1}\| \leq 1$, $\tau \circ \varphi_{n+1} = 1$ on F_{n+1} and $\|(\varphi_{n+1} - \varphi_n)|F_n\| < 1/2^n$, completing induction. Now the sequence $\{\varphi_n \circ \pi_n\}$ is uniformly bounded and

$$\sum_{k=n}^{\infty} \left\| (\varphi_{k+1} - \varphi_k) \Big| F_n \right\| \leq \sum_{j=n}^{\infty} 1/2^j < \infty$$

guaranteeing convergence of $\varphi_k(x)$ for every $x \in F_n$ as $k \rightarrow \infty$. Then $\{\varphi_n \circ \pi_n\}$ converges strongly to some map φ from E/M to E . Obviously φ is a required linear lifting.

It is better to introduce some terminology before stating the main

result on linear lifting. A Banach space E is called a π -space if there is a sequence $\{F_n\}$ of finite dimensional subspaces such that $F_1 \subseteq F_2 \subseteq \dots$ with $\overline{\bigcup_{n=1}^{\infty} F_n} = E$ and each F_n is the range of a projection of norm one. Here projections π_n can be assumed to have the property that $\pi_n \pi_m = \pi_n$ for $n \leq m$ and that π_n converges strongly to the identity as $n \rightarrow \infty$. An ordered Banach space is called a Π -space if, in addition, projections can be chosen positive and if each F_n has the positive cone generated by a linearly independent basis.

THEOREM 5. *Suppose that the annihilator of a closed subspace M is the range of a projection P such that*

$$\|f\| = \|Pf\| + \|f - Pf\| \quad (f \in E^*).$$

If the quotient space E/M becomes a π -space then there is a linear lifting of norm one, or equivalently, M is the kernel of a projection of norm one.

Proof. Since the unit ball U is splittable by Corollary 1, all requirements in Lemma 9 are fulfilled with $S = U$ and $L = \tau(U)$ by Corollary 7.

COROLLARY 8. *Let N and M be closed subspaces and suppose that M^\perp is the range of a projection P such that $P(N^\perp) \subseteq N^\perp$ and*

$$\|f\| = \|Pf\| + \|f - Pf\| \quad (f \in E^*).$$

If the quotient space $N/N \cap M$ is a π -space, there is a linear lifting of norm one from $N/N \cap M$ to N .

Proof. In view of Theorem 5 it suffices to prove that the annihilator of $N \cap M$ in N^* is the range of a projection \mathcal{P} such that

$$\|g\| = \|\mathcal{P}g\| + \|g - \mathcal{P}g\| \quad (g \in N^*).$$

When N^* is identified with E^*/N^\perp , the annihilator of $N \cap M$ becomes the image of $(N \cap M)^\perp$ under the canonical map from E^* to E^*/N^\perp . Since hypothesis implies splittability of N by Corollary 3, $N + M$ is closed by Theorem 1 so that $N^\perp + M^\perp$ is weak* closed and coincides with $(N \cap M)^\perp$. Therefore the annihilator of $N \cap M$ in N^* becomes the image of M^\perp in E^*/N^\perp . Since N^\perp is invariant under P , there arises a natural projection \mathcal{P} from N^* to the annihilator $N \cap M$. Since by hypothesis

$$\|f - N^\perp\| \geq \|Pf - N^\perp\| + \|f - Pf - N^\perp\|,$$

\mathcal{P} is easily seen to have the required property.

When E is the space of continuous functions on a compact set and M consists of functions vanishing on a fixed closed subset, Corollary 8 was proved by Michael and Pełczyński [8].

THEOREM 6. *Let M be a closed subspace of an ordered Banach space E . Suppose that M is the range of a projection P such that $f \geq Pf \geq 0$ ($f \geq 0$) and*

$$\|f\| = \|Pf\| + \|f - Pf\| \quad (f \in E^*).$$

If E/M is a Π -space under the canonical ordering, there is a positive linear lifting of norm one, or equivalently, M is the kernel of a positive projection of norm one.

Proof. Since the positive cone E_+ is splittable by Corollary 3, all requirements in Lemma 9 are fulfilled with $S = E_+$ and $L = \tau(E_+)$ by definition of a Π -space and Corollary 6.

To be a π -space or a Π -space is not so severe restriction. Let us prove:

Separable complex (resp. real) L_p ($1 \leq p < \infty$) and complex (resp. real) $C(X)$ on compact metrizable X are π -spaces (resp. Π -spaces).

In fact, it suffices for the first part to treat a L_p space on a finite measure space (\mathcal{B}, μ) . Since the Borel field is separable with respect to μ , there is an increasing sequence $\{\mathcal{B}_n\}$ of finite Borel subfields such that $\bigcup_{n=1}^{\infty} L_p(\mathcal{B}_n)$ is dense in L_p where $L_p(\mathcal{B}_n)$ is the subspace of \mathcal{B}_n -measurable functions. Each $L_p(\mathcal{B}_n)$ is finite dimensional and the conditional expectation relative to \mathcal{B}_n becomes a (positive) projection of norm one from L_p to $L_p(\mathcal{B}_n)$ (cf. [3]). The assertion for $C(X)$ is proved in [7] by using peaked partition.

REFERENCES

1. E. M. Alfsen and B. Hirsberg, *On dominated extensions in linear subspaces of $C_c(X)$* , Pacific J. Math., **36** (1971), 567-584.
2. T. Ando, *On fundamental properties of a Banach space with a cone*, Pacific J. Math., **12** (1962), 1163-1169.
3. ———, *Contractive projections in L_p spaces*, Pacific J. Math., **17** (1966), 391-405.
4. E. B. Davies, *The structure and ideal theory of the predual of a Banach lattice*, Trans. Amer. Math. Soc., **131** (1968), 544-555.
5. T. Gamelin, *Restrictions of subspaces of $C(X)$* , Trans. Amer. Math. Soc., **112** (1964), 278-286.
6. ———, *Uniform algebra*, Prentice-Hall, Englewood Cliffs, N. J. 1969.
7. E. Michael and A. Pełczyński, *Peaked partition subspaces of $C(X)$* , Illinois J. Math., **11** (1967), 555-562.

8. E. Michael and A. Pełczynski, *A linear extension theorem*, Illinois J. Math., **11** (1967), 563-579.
9. F. Perdrizet, *Espaces de Banach ordonnés et idéaux*, J. Math. Pures Appl., **49** (1970), 61-98.
10. H. H. Schaefer, *Topological Vector Spaces*, MacMillan N. Y., 1966.

Received October 13, 1971.

RESEARCH INSTITUTE OF APPLIED ELECTRICITY
HOKKAIDO UNIVERSITY, SAPPORO, JAPAN

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$48.00 a year (6 Vols., 12 issues). Special rate: \$24.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics

Vol. 44, No. 2

June, 1973

Tsuyoshi Andô, <i>Closed range theorems for convex sets and linear liftings</i>	393
Richard David Bourgin, <i>Conically bounded sets in Banach spaces</i>	411
Robert Jay Buck, <i>Hausdorff dimensions for compact sets in R^n</i>	421
Henry Cheng, <i>A constructive Riemann mapping theorem</i>	435
David Fleming Dawson, <i>Summability of subsequences and stretchings of sequences</i>	455
William Thomas Eaton, <i>A two sided approximation theorem for 2-spheres</i>	461
Jay Paul Fillmore and John Herman Scheuneman, <i>Fundamental groups of compact complete locally affine complex surfaces</i>	487
Avner Friedman, <i>Bounded entire solutions of elliptic equations</i>	497
Ronald Francis Gariepy, <i>Multiplicity and the area of an $(n - 1)$ continuous mapping</i>	509
Andrew M. W. Glass, <i>Archimedean extensions of directed interpolation groups</i>	515
Morisuke Hasumi, <i>Extreme points and unicity of extremum problems in H^1 on polydiscs</i>	523
Trevor Ongley Hawkes, <i>On the Fitting length of a soluble linear group</i>	537
Garry Arthur Helzer, <i>Semi-primary split rings</i>	541
Melvin Hochster, <i>Expanded radical ideals and semiregular ideals</i>	553
Keizō Kikuchi, <i>Starlike and convex mappings in several complex variables</i>	569
Charles Philip Lanski, <i>On the relationship of a ring and the subring generated by its symmetric elements</i>	581
Jimmie Don Lawson, <i>Intrinsic topologies in topological lattices and semilattices</i>	593
Roy Bruce Levow, <i>Counterexamples to conjectures of Ryser and de Oliveira</i>	603
Arthur Larry Lieberman, <i>Some representations of the automorphism group of an infinite continuous homogeneous measure algebra</i>	607
William George McArthur, <i>G_δ-diagonals and metrization theorems</i>	613
James Murdoch McPherson, <i>Wild arcs in three-space. II. An invariant of non-oriented local type</i>	619
H. Millington and Maurice Sion, <i>Inverse systems of group-valued measures</i>	637
William James Rae Mitchell, <i>Simple periodic rings</i>	651
C. Edward Moore, <i>Concrete semispaces and lexicographic separation of convex sets</i>	659
Jingyal Pak, <i>Actions of torus T^n on $(n + 1)$-manifolds M^{n+1}</i>	671
Merrell Lee Patrick, <i>Extensions of inequalities of the Laguerre and Turán type</i>	675
Harold L. Peterson, Jr., <i>Discontinuous characters and subgroups of finite index</i>	683
S. P. Philipp, <i>Abel summability of conjugate integrals</i>	693
R. B. Quintana and Charles R. B. Wright, <i>On groups of exponent four satisfying an Engel condition</i>	701
Marlon C. Rayburn, <i>On Hausdorff compactifications</i>	707
Martin G. Ribe, <i>Necessary convexity conditions for the Hahn-Banach theorem in metrizable spaces</i>	715
Ryōtarō Satō, <i>On decomposition of transformations in infinite measure spaces</i>	733
Peter Drummond Taylor, <i>Subgradients of a convex function obtained from a directional derivative</i>	739
James William Thomas, <i>A bifurcation theorem for k-set contractions</i>	749
Clifford Edward Weil, <i>A topological lemma and applications to real functions</i>	757
Stephen Andrew Williams, <i>A nonlinear elliptic boundary value problem</i>	767
Pak-Ken Wong, <i>$*$-actions in A^*-algebras</i>	775