

Pacific Journal of Mathematics

**SIMULTANEOUS APPROXIMATION AND INTERPOLATION IN
 L_1 AND $C(T)$**

JOSEPH MICHAEL LAMBERT

SIMULTANEOUS APPROXIMATION AND INTERPOLATION IN L_1 AND $C(T)$

JOSEPH M. LAMBERT

Given a dense subspace of M of a Banach space X , an element x in X and a finite collection of linear functions in X^* , the problem of simultaneous approximation and interpolation is to interpolate x at the given functionals in X^* by an element m of M , with the restriction that the norms of x and m be equal and their difference in norm be arbitrarily small. A solution is given for the space L_1 with dense subspace, the simple functions in L_1 , and any collection of functions in L_∞ . In addition the problem is studied in the space $C(T)$, with any dense subalgebra and any finite collection of linear functionals in $C(T)^*$.

In [1] the concept of simultaneous approximation and interpolation which preserves the norm, (SAIN), was introduced.

DEFINITION [1]. Let X be a normed linear space, M a dense subset of X , L a finite dimensional subspace of X^* . The triple (X, M, L) has property (SAIN) if for every x in X and $\varepsilon > 0$ there exists y in M such that $\|x - y\| < \varepsilon$, $\|x\| = \|y\|$ and $\lambda(x) = \lambda(y)$ for all λ in L .

Other papers concerned with this topic are [4], [5], and [6].

In [5] it was shown that if L is any finite dimensional subspace of l_∞ and if M is the subspace of l_1 consisting of the elements having only finitely many nonzero components, then (l_1, M, L) had property (SAIN). In this paper, we let M be the subspace of simple functions in L_1 . We show (L_1, M, T) has property (SAIN) for any finite dimensional subspace T in L_∞ .

In [1], the space $C(T)$ is studied, where T is a compact Hausdorff space. One finds there

THEOREM 4.1. *Let A be a dense subalgebra of $C(T)$ and t_1, \dots, t_n in T . Then $(C(T), A, \{\delta_{t_1}, \dots, \delta_{t_n}\})$ has property (SAIN). (δ_t is the linear functional on $C(T)$ given by point evaluation at t .)*

When arbitrary linear functionals in $C(T)$ are used, examples in [1] show that $(C(T), A, \{\nu\})$ may or may not have property (SAIN) depending on ν .

In this paper we wish to find sufficient conditions on f in $C(T)$ and M dense in $C(T)$ such that given $\{\nu_1, \dots, \nu_n\}$ in $C(T)^*$ and $\varepsilon > 0$ there exists m in M such that $\|f - m\| < \varepsilon$, $\|f\| = \|m\|$ and

$$\int f d\nu_i = \int m d\nu_i, i = 1, \dots, n.$$

In particular one finds that if f attains its norm at most a finite number of times, then any dense subalgebra of $C(T)$ will satisfy these conditions.

In this paper the following notation and terminology is used. X is to denote a real normed linear space. X^* is to denote the continuous dual of X , $U(X)$ and $S(X)$, the closed unit ball and its boundary in X . A set E contained in a set F is F -extremal if whenever $tx + (1-t)y$ is in E , with $0 < t < 1$ and x, y in F then x, y are in E . A hyperplane H supports a set K , if it bounds K and intersects K . The real valued function $\text{sgn}(\cdot): \text{Reals} \rightarrow \{-1, 0, 1\}$ is defined via $\text{sgn}(0) = 0$ and $\text{sgn}(x) = x/|x|, x \neq 0$. Then convex hull of a set A is to be denoted by $\text{co}(A)$. All other notation will correspond to that of [3].

1. Minimal closed $U(X)$ extremal subsets.

DEFINITION 1.1. $F(x)$ is to denote the minimal closed $U(X)$ -extremal set containing x . $Q(x)$ is the intersection of all $U(X)$ supporting hyperplanes at x .

THEOREM 1.1. *Let X be a normed linear space, M a dense subspace of X and $L = \text{span}\{\varphi_1, \dots, \varphi_k\}$ a finite dimensional subspace of X^* , and x in $S(X)$. If $F(x) \cap M$ is dense in $F(x)$ then given $\varepsilon > 0$ there exists m in $S(X)$ such that $\varphi_i(x) = \varphi_i(m), i = 1, \dots, k$ and $\|x - m\| < \varepsilon$.*

Proof. Define the continuous function $\varphi: F(x) \rightarrow R^k$ via $\varphi(x) = (\varphi_1(x), \dots, \varphi_k(x))$. Assume that $F(x) \subset \varphi_i^{-1}(\varphi_i(x))$ for $i = 0, 1, \dots, k$ and that this is the largest set of linearly independent elements of L for which this is true. If no such set exists, $k = 0$. In R^{n-k} we assert the existence of $m_\alpha \in F(x) \cap M$ with $\|x - m_\alpha\| < \varepsilon$ such that

$$(\varphi_{k+1}(x), \dots, \varphi_n(x)) \in \text{co}(\varphi_{k+1}(m_\alpha), \dots, \varphi_n(m_\alpha) \mid \alpha \in A),$$

A an arbitrary index set. If not, then in R^{n-k} there exists a linear functional τ , a linear combination of the $\varphi_i, i > k$ such that without loss of generality $\tau(m) \leq \tau(x)$ for all $m \in F(x) \cap M$ such that $\|x - m\| < \varepsilon$. But this implies $\tau(m) \leq \tau(x)$ for all $m \in F(x) \cap M$, since if there exists $m_0 \in F(x) \cap M$ with $\|x - m_0\| > \varepsilon$ then the set

$$\{y \in F(x) \mid \tau(y) > \tau(x), \|y - x\| < \varepsilon\}$$

is $F(x)$ relatively open and nonempty (choose a suitable combination

of x and m_0) and hence contains m in $F(x) \cap M$ contradicting $\tau(m) \leq \tau(x)$ with $\|m - x\| < \varepsilon$. Since $F(x) \cap M$ is dense in $F(x)$ this implies $\tau(y) \leq \tau(x) \forall y \in F(x)$. Let $K = \{y \in F(x) \mid \tau(y) = \tau(x)\}$. K is convex closed and $F(x)$ -extremal since $tz + (1 - t)y$ in K implies $t\tau(z) + (1 - t)\tau(y) = \tau(x)$ with $\tau(z) \leq \tau(x)$, $\tau(y) \leq \tau(x)$. Hence $\tau(z) = \tau(y) = \tau(x)$ and $z, y \in K$. Hence K is closed $U(X)$ -extremal and $K = F(x)$. Thus $F(x) \subset \tau^{-1}(\tau(x))$. Since τ is linearly independent of $\varphi_i, i = 1, \dots, k$, this contradicts the maximal choice of φ_i at the start of the proof. Therefore

$$(\varphi_{k+1}(x), \dots, \varphi_n(x)) \in \text{co}(\varphi_{k+1}(m_\alpha), \dots, \varphi_n(m_\alpha) \mid \alpha \in A)$$

with $\|x - m_\alpha\| < \varepsilon$. This yields the result by the convexity of M and $\varphi(M)$.

In a recent paper of Deutsch and Lindahl [2], they showed that in certain spaces that the set $Q(x)$, the intersection of all $U(X)$ supporting hyperplanes at x , is equal to the closure of the minimal extremal subset containing x . Thus $Q(x)$ is equal to the minimal closed extremal subset containing x . This occurs, in particular [2, Theorem 4.2], if (T, Σ, ν) is a σ -finite measure space, in $L_1(T, \Sigma, \nu)$. Also, this occurs [2, Theorem 3.3] in the space $C_0(T)$, the space of continuous functions vanishing at infinity, T locally compact.

THEOREM 2.1. *Let (T, Σ, ν) be a σ -finite measure space with $L_1^*(T, \Sigma, \nu) = L_\infty(T, \Sigma, \nu)$. Let M be the dense subspace of L_1 consisting of the simple functions. Then (L_1, M, H) has property (SAIN) for any finite dimensional subspace H contained in L_∞ .*

Proof. Given x in $S(L_1)$. By [2, Theorem 4.2], $Q(x) = \{z \in S(L_1) \mid \int z \text{sgn}(x) = 1\}$ and $Q(x) = F(x)$. M is dense in $Q(x)$ and by Theorem 1.1 the result follows.

THEOREM 2.2. *Let T be a compact Hausdorff space, $C(T)$ the space of real valued continuous functions on T . Let f in $S(C(T))$ be such that $Q(f) = \bigcap_{i=1}^r \varphi_i^{-1}(\|f\|)$ with φ_i in $\text{rca}(T)$. If*

$$(C(T), M, \{\varphi_i \mid i = 1, \dots, n\})$$

has property SAIN then given any finite collection μ_i in $\text{rca}(T)$, $\varepsilon > 0$ there exists m in M such that $\|f - m\| < \varepsilon, \|f\| = \|m\|$ and $\int f d\mu_i = \int m d\mu_i$.

Proof. By [2, Theorem 3.3] $Q(f) = \{x \in C(T) \mid x(t) = f(t) \text{ for } t \in T \text{ such that } |f(t)| = \|f\|\}$ and $Q(f) = F(f)$. $(C(T), M, \{\varphi_i \mid i = 1, \dots, n\})$ having property (SAIN) implies $F(f) \cap M$ is dense in $F(f)$. By Theorem 1.1 the result follows.

COROLLARY 2.1. *Let f be in $S(C(T))$ $\varepsilon > 0$. If $|f|$ attains its norm finitely often then given $\mu_i, i = 1, \dots, n$ in $rca(T)$ there exists p in A (any dense subalgebra of $C(T)$) such that $\|p - f\| < \varepsilon \|p\| = \|f\|$ and $\int p d\mu_i = \int f d\mu_i$.*

Proof. By [1, Theorem 4.1] quoted in the introduction of this article ($(C(T), A, \{\delta_i | f(t) | = 1\})$ has property (SAIN). But $Q(f) = \cap \{\delta_i^{-1}(f(t) | |f(t) | = 1\}$. Hence apply Theorem 2.2.

ACKNOWLEDGMENT. The author is indebted to the referee for his criticism of the initially submitted article and for his suggestions for revision.

REFERENCES

1. F. Deutsch and P. Morris, *On simultaneous approximation and interpolation which preserves the norm*, J. Approximation Theory, **2** (1969), 355-373.
2. F. Deutsch and R. Lindahl, *Minimal extremal subsets of the unit sphere*, Math. Annalen (to appear).
3. N. Dunford and J. T. Schwartz, *Linear Operators, Part I, General Theory*, Interscience, New York, 1958.
4. R. Holmes and J. Lambert, *A geometrical approach to property (SAIN)*, (to appear)
5. J. Lambert, *Simultaneous approximation and interpolation in l_1* , Proc. Amer. Math. Soc. **32** (1972), 150-152.
6. H. McLaughlin and P. Zaretski, *Simultaneous approximation and interpolation with norm preservation*, J. Approximation Theory, **4** (1970), 54-58.

Received November 21, 1971 and in revised form April 17, 1972.

THE PENNSYLVANIA STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

William George Bade, <i>Complementation problems for the Baire classes</i>	1
Ian Douglas Brown, <i>Representation of finitely generated nilpotent groups</i>	13
Hans-Heinrich Brungs, <i>Left Euclidean rings</i>	27
Victor P. Camillo and John Cozzens, <i>A theorem on Noetherian hereditary rings</i>	35
James Cecil Cantrell, <i>Codimension one embeddings of manifolds with locally flat triangulations</i>	43
L. Carlitz, <i>Enumeration of up-down permutations by number of rises</i>	49
Thomas Ashland Chapman, <i>Surgery and handle straightening in Hilbert cube manifolds</i>	59
Roger Cook, <i>On the fractional parts of a set of points. II</i>	81
Samuel Harry Cox, Jr., <i>Commutative endomorphism rings</i>	87
Michael A. Engber, <i>A criterion for divisoriality</i>	93
Carl Clifton Faith, <i>When are proper cyclics injective</i>	97
David Finkel, <i>Local control and factorization of the focal subgroup</i>	113
Theodore William Gamelin and John Brady Garnett, <i>Bounded approximation by rational functions</i>	129
Kazimierz Goebel, <i>On the minimal displacement of points under Lipschitzian mappings</i>	151
Frederick Paul Greenleaf and Martin Allen Moskowitz, <i>Cyclic vectors for representations associated with positive definite measures: nonseparable groups</i>	165
Thomas Guy Hallam and Nelson Onuchic, <i>Asymptotic relations between perturbed linear systems of ordinary differential equations</i>	187
David Kent Harrison and Hoyt D. Warner, <i>Infinite primes of fields and completions</i>	201
James Michael Hornell, <i>Divisorial complete intersections</i>	217
Jan W. Jaworowski, <i>Equivariant extensions of maps</i>	229
John Jobe, <i>Dendrites, dimension, and the inverse arc function</i>	245
Gerald William Johnson and David Lee Skoug, <i>Feynman integrals of non-factorable finite-dimensional functionals</i>	257
Dong S. Kim, <i>A boundary for the algebras of bounded holomorphic functions</i>	269
Abel Klein, <i>Renormalized products of the generalized free field and its derivatives</i> ...	275
Joseph Michael Lambert, <i>Simultaneous approximation and interpolation in L_1 and $C(T)$</i>	293
Kelly Denis McKennon, <i>Multipliers of type (p, p) and multipliers of the group L_p-algebras</i>	297
William Charles Nemitz and Thomas Paul Whaley, <i>Varieties of implicative semi-lattices. II</i>	303
Donald Steven Passman, <i>Some isolated subsets of infinite solvable groups</i>	313
Norma Mary Piacun and Li Pi Su, <i>Wallman compactifications on E-completely regular spaces</i>	321
Jack Ray Porter and Charles I. Votaw, <i>$S(\alpha)$ spaces and regular Hausdorff extensions</i>	327
Gary Sampson, <i>Two-sided L_p estimates of convolution transforms</i>	347
Ralph Edwin Showalter, <i>Equations with operators forming a right angle</i>	357
Raymond Earl Smithson, <i>Fixed points in partially ordered sets</i>	363
Victor Snaith and John James Ucci, <i>Three remarks on symmetric products and symmetric maps</i>	369
Thomas Rolf Turner, <i>Double commutants of weighted shifts</i>	379
George Kenneth Williams, <i>Mappings and decompositions</i>	387