

Pacific Journal of Mathematics

**UNIFORM INTEGRABILITY OF DERIVATIVES ON
 σ -LATTICES**

ALLAN FRANCIS ABRAHAMSE

UNIFORM INTEGRABILITY OF DERIVATIVES ON σ -LATTICES

ALLAN F. ABRAHAMSE

This paper contains a new derivation of the Radon-Nikodym derivative on a σ -lattice. Absolute continuity is defined in this setting, and the definition is justified by obtaining an extension of a standard result on the uniform integrability of derivatives. An application to mean convergence of martingales and a version of Jensen's inequality are given.

Let (Ω, \mathcal{F}, P) be a probability space, φ a finite signed measure on \mathcal{F} , and \mathcal{M} a σ -lattice of elements of \mathcal{F} . By this, we mean \mathcal{M} contains Ω and the empty set \emptyset , and is closed under countable unions and countable intersections. We denote by \mathcal{M}^c the collection of complements of elements of \mathcal{M} . An extended real valued function X on Ω is said to be measurable with respect to \mathcal{M} (we will write $X \in \mathcal{M}$) if sets of the form $\{X \geq a\} = \{\omega \mid X(\omega) \geq a\}$ are in \mathcal{M} for every real a .

In [3] it is shown that there exists an a.s. unique (a.s. with respect to P) function $X \in \mathcal{M}$ such that for every real a ,

$$\begin{aligned} \varphi(A \cap \{X \geq a\}) &\geq aP(A \cap \{X \geq a\}) & A \in \mathcal{M}^c \\ \varphi(A \cap \{X \leq a\}) &\leq aP(A \cap \{X \leq a\}) & A \in \mathcal{M} . \end{aligned}$$

X is called the derivative of φ with respect to P on \mathcal{M} . We denote this by $X = D(\varphi, \mathcal{M})$.

In this paper we present a new proof of the existence of this derivative X , based on the observation that the set $\{X \geq a\}$ maximizes $\varphi(A) - aP(A)$ as A runs over \mathcal{M} . We introduce what appears to be suitable definition of absolute continuity in the present setting. Our main result is that the collection of derivatives $\{D(\varphi, \mathcal{M}_0)\}$, for $\mathcal{M}_0 \subseteq \mathcal{M}$, is uniformly integrable if and only if φ is absolutely continuous with respect to P on \mathcal{M} . As an application of this, we can strengthen a result in [2] concerning mean convergence of martingales over σ -lattices.

Finally, we prove a version of Jensen's inequality for the present setting. This enables one to obtain very easily the results in [2].

The σ -lattice \mathcal{M} is a σ -field if and only if $\mathcal{M} = \mathcal{M}^c$. It will be noted that in all that follows, if \mathcal{M} is a σ -field the standard theory is obtained. We have here direct generalizations of the classical theory. What is remarkable is that little extra work seems to be entailed.

2. The derivative. We begin by defining:

$$(1) \quad \begin{aligned} \varphi^+ &= \sup \{ \varphi(A) \mid A \in \mathcal{M} \} \\ \varphi^- &= \inf \{ \varphi(A) \mid A \in \mathcal{M}^c \}. \end{aligned}$$

Then the following holds for $A_1 \in \mathcal{M}^c, A_2 \in \mathcal{M}$:

$$(2) \quad \varphi(A_1) - \varphi^- \geq \varphi(A_1 \cap A_2) \geq \varphi(A_2) - \varphi^+.$$

To establish the right hand inequality, we note that $A_1^c \cap A_2 \in \mathcal{M}$, hence $\varphi(A_1^c \cap A_2) \leq \varphi^+$. Then $\varphi(A_1 \cap A_2) = \varphi(A_2) - \varphi(A_1^c \cap A_2) \geq \varphi(A_2) - \varphi^+$. The left hand inequality follows similarly.

THEOREM 1. (a) *There is a $A \in \mathcal{M}$ such that $\varphi(A) = \varphi^+$.* (b) *There is a $A \in \mathcal{M}^c$ such that $\varphi(A) = \varphi^-$.*

Proof. For each $n \geq 1$, choose $A_n \in \mathcal{M}$ such that $\varphi(A_n) \geq \varphi^+ - 2^{-n}$. Then

$$\begin{aligned} \varphi\left(\bigcup_{m=n}^{\infty} A_m\right) &= \varphi(A_n) + \sum_{m=n+1}^{\infty} \varphi(A_m - (A_n \cup \dots \cup A_{m-1})) \\ &\geq \varphi(A_n) + \sum_{m=n+1}^{\infty} (\varphi(A_m) - \varphi^+) \\ &\geq \varphi^+ - \sum_{m=n}^{\infty} 2^{-m} = \varphi^+ - 2^{-n+1}. \end{aligned}$$

Let $A = \limsup A_n$, and by continuity of φ , $\varphi(A) = \varphi^+$.

The proof of (b) is similar.

COROLLARY 1. $\varphi(\Omega) = \varphi^+ + \varphi^-$.

Proof. Select $A \in \mathcal{M}$ so that $\varphi(A) = \varphi^+$. Then $\varphi(\Omega) \geq \varphi^+ + \varphi^-$. The reverse inequality is obtained in an obvious way.

COROLLARY 2. *For $A \in \mathcal{M}$, $\varphi(A) = \varphi^+$ if and only if $\varphi(A^c) = \varphi^-$.*

Proof. If $A \in \mathcal{M}$ and $\varphi(A) = \varphi^+$, then $\varphi(A^c) = \varphi(\Omega) - \varphi(A) = \varphi(\Omega) - \varphi^+ = \varphi^-$.

THEOREM 2. *If $A \in \mathcal{M}$, the following are equivalent:*

- (a) $\varphi(A) = \varphi^+$
- (b) $\varphi(A_0 \cap A) \geq 0$ for all $A_0 \in \mathcal{M}^c$, and $\varphi(A_0 \cap A^c) \leq 0$ for all $A_0 \in \mathcal{M}$.

Proof. Assume (a). Then $\varphi(A^c) = \varphi^-$. The first inequality of (b) follows from the right hand inequality of (2), and the second from the left hand.

Conversely, assume (b) is valid for $A \in \mathcal{M}$. Select $A_1 \in \mathcal{M}$ so that $\varphi(A_1) = \varphi^+$. Then $\varphi(A^c \cap A_1) \geq 0$ from (2) and $\varphi(A^c \cap A_1) \leq 0$ from (b), hence $\varphi(A^c \cap A_1) = 0$. Similarly, $\varphi(A \cap A_1^c) = 0$, and so $\varphi(A) = \varphi(A_1) = \varphi^+$. This completes our proof.

DEFINITION. A function $X \in \mathcal{M}$ is a derivative of φ with respect to P if for every real a ,

$$\begin{aligned} \varphi(A \cap \{X \geq a\}) &\geq aP(A \cap \{X \geq a\}) & A \in \mathcal{M}^c \\ \varphi(A \cap \{X \leq a\}) &\leq aP(A \cap \{X \leq a\}) & A \in \mathcal{M}. \end{aligned}$$

For any real a , let $\varphi_a = \varphi - aP$, and define φ_a^+ and φ_a^- by (1), using φ_a . Since for each fixed $A \in \mathcal{F}$, $\varphi_a(A)$ is nonincreasing, so are φ_a^+ and φ_a^- .

COROLLARY. $X \in \mathcal{M}$ is a derivative of φ with respect to P on \mathcal{M} if and only if $\varphi_a(\{X \geq a\}) = \varphi_a^+$ for every real a .

The reader is referred to [4], p. 108, for a classical version of the following.

THEOREM 3. A derivative of φ with respect to P on \mathcal{M} exists, and is a.s. unique.

Proof. Define a collection K of functions by setting

$$K = \{X \in \mathcal{M} \mid \varphi_a(A \cap \{X \geq a\}) \geq 0 \text{ all } A \in \mathcal{M}^c, \text{ all real } a\}.$$

For any real a , we can choose $A_a \in \mathcal{M}$ so that $\varphi(A_a) = \varphi_a^+$. Let

$$(3) \quad \begin{aligned} X_a(\omega) &= a & \omega \in A_a \\ &= -\infty & \omega \in A_a^c. \end{aligned}$$

Then we claim $X_a \in K$. For suppose $b > a$. Then $\{X_a \geq b\} = \emptyset$, so $\varphi_a(A \cap \{X_a \geq b\}) = 0$. On the other hand, if $b \leq a$, then $\{X_a \geq b\} = A_a$, and $\varphi_a(A \cap \{X_a \geq b\}) \geq \varphi_a(A \cap \{X_a \geq b\}) \geq 0$ by Theorem 2.

We now note that if $X \in K$, $Y \in \mathcal{M}$ and $X \geq Y$, then for any $A \in \mathcal{M}^c$,

$$\varphi_a(A \cap \{X \geq a\}) = \varphi_a(A \cap \{Y \geq a\}) + \varphi_a(A \cap \{a > Y\} \cap \{X \geq a\}).$$

Since $A \cap \{a > Y\} \in \mathcal{M}^c$, the right-most term is nonnegative, hence

$$(4) \quad \varphi_a(A \cap \{X \geq a\}) \geq \varphi_a(A \cap \{Y \geq a\}).$$

Now suppose X_1, X_2, \dots are elements of K . Let

$$Z = \sup(X_1, X_2, \dots).$$

Certainly $Z \in \mathcal{M}$. We define

$$\begin{aligned} A_1 &= \Omega \\ A_k &= \{X_1 \leq a, \dots, X_{k-1} \leq a\} \quad k > 1. \end{aligned}$$

Then $A_k \in \mathcal{M}^c$ for each $k \geq 1$, and

$$\{Z > a\} = \bigcup_{k \geq 1} A_k \cap \{X_k > a\}.$$

The terms on the right are pair-wise disjoint, so for any $A \in \mathcal{M}^c$,

$$\varphi_a(A \cap \{Z > a\}) = \sum_{k \geq 1} \varphi_a(A \cap A_k \cap \{X_k > a\}) \geq 0.$$

That $\varphi_a(A \cap \{Z \geq a\}) \geq 0$ follows by continuity, whence $Z \in K$.

Let Q be a countable dense subset of the reals containing the discontinuities of φ_a^+ . Define

$$X = \sup \{X_q \mid q \in Q\}$$

where X_q is defined by (3). From the preceding argument $X \in K$, and from (4),

$$\varphi_q^+ \geq \varphi_q(X \geq q) \geq \varphi_q(X_q \geq q) = \varphi_q^+$$

for all $q \in Q$. It follows that $\varphi_a(X \geq a) = \varphi_a^+$, hence X is a derivative. Uniqueness a.s. is implied by the following lemma.

LEMMA 1. *Suppose X, Y are measurable with respect to \mathcal{M} , and for all real a ,*

$$\begin{aligned} \varphi_a(A \cap \{Y \geq a\}) &\geq 0 & A \in \mathcal{M}^c \\ \varphi_a(A \cap \{X \leq a\}) &\leq 0 & A \in \mathcal{M}. \end{aligned}$$

Then $Y \leq X$ a.s.

Proof. Choose $a < b$. Then

$$0 \leq \varphi_b(\{X \leq a < b \leq Y\}) \leq \varphi_a(\{X \leq a < b \leq Y\}) \leq 0.$$

It follows that

$$bP(\{X \leq a < b \leq Y\}) = aP(\{X \leq a < b \leq Y\})$$

and so $P(\{X \leq a < b \leq Y\}) = 0$, from which our result clearly follows.

We now give some elementary properties of the derivative.

LEMMA 2. *Let φ^1, φ^2 be finite signed measures on \mathcal{F} , with $\varphi^1 \leq \varphi^2$. Then $D(\varphi^1, \mathcal{M}) \leq D(\varphi^2, \mathcal{M})$ a.s.*

Proof. Let $X^k = D(\varphi^k, \mathcal{M})$, $k = 1, 2$. Then

$$\varphi_a^1(\Lambda \cap \{X^2 \leq a\}) \leq \varphi_a^2(\Lambda \cap \{X^2 \leq a\}) \leq 0$$

for $\Lambda \in \mathcal{M}$, any real a . Since $\varphi_a^1(\Lambda \cap \{X^1 \geq a\}) \geq 0$ for $\Lambda \in \mathcal{M}^c$, any real a , the result follows from Lemma 1.

LEMMA 3.

$$\begin{aligned} D(a\varphi + bP, \mathcal{M}) &= aD(\varphi, \mathcal{M}) + b & a \geq 0 \\ &= aD(\varphi, \mathcal{M}^c) + b & a < 0. \end{aligned}$$

Suppose $a < 0$, and let $\Psi = a\varphi + bP$, and $X = D(\varphi, \mathcal{M})$. Since $aX + b \in \mathcal{M}^c$, we must show that $\Psi_t^+ = \Psi_t(aX + b \geq t)$. But

$$\begin{aligned} \Psi_t^+ &= a\varphi_{(t-b)/a}^- \\ &= a\varphi_{(t-b)/a}(X \leq (t-b)/a) \\ &= a\varphi_{(t-b)/a}(aX + b \geq t) \\ &= \Psi_t(aX + b \geq t). \end{aligned}$$

This proves our assertion. When $a = 0$ or $a > 0$, similar proofs work.

Finally, we note that $|D(\varphi, \mathcal{M})|$ is a.s. finite. For

$$aP(X \geq a) \leq \varphi(X \geq a) \leq \varphi^+$$

hence

$$P(X \geq a) \leq \frac{\varphi^+}{a}$$

and so

$$P(X = \infty) = \lim_{a \rightarrow \infty} P(X \geq a) = 0.$$

Similarly we can prove $P(X = -\infty) = 0$.

3. Absolute continuity and uniform integrability. When $X \in \mathcal{F}$, $\Lambda \in \mathcal{F}$, we will write

$$E(X; \Lambda) = \int_{\Lambda} X dP$$

if the right hand side is well defined. The following result is in [3], but since the proof is not long, we will include it here.

THEOREM 4. If $X = D(\varphi, \mathcal{M})$, then for a, b real,

$$(5) \quad \begin{aligned} \varphi(\Lambda \cap \{a \leq X \leq b\}) &\geq E(X; \Lambda \cap \{a \leq X \leq b\}) & \Lambda \in \mathcal{M}^c \\ &\leq E(X; \Lambda \cap \{a \leq X \leq b\}) & \Lambda \in \mathcal{M}. \end{aligned}$$

Proof. For any $n \geq 1$, choose a_0, a_1, \dots, a_n , with $a_0 = a - 1/n$, $a_{k+1} - a_k = (b - a_0)/n$, $a_n = b$. For $A \in \mathcal{M}^c$,

$$\begin{aligned} \varphi(A \cap \{a - \frac{1}{n} < X \leq b\}) &= \sum_{k=0}^{n-1} \varphi(A \cap \{a_k < X \leq a_{k+1}\}) \\ &\geq \sum_{k=0}^{n-1} a_k P(A \cap \{a_k < X \leq a_{k+1}\}). \end{aligned}$$

As n increases, the right and left hand terms of this inequality approach the corresponding terms of (5). For $A \in \mathcal{M}$, a similar proof works.

COROLLARY. $D(\varphi, \mathcal{M})$ is integrable.

Proof. Since $\Omega \in \mathcal{M} \cap \mathcal{M}^c$, both inequalities in Theorem 4 hold, and we obtain

$$\varphi(\{a \leq X \leq b\}) = E(X; \{a \leq X \leq b\}).$$

Our conclusion follows from the fact that $P(|X| < \infty) = 1$ and φ is finite.

DEFINITION. φ is absolutely continuous with respect to P on \mathcal{M} if the following conditions hold: (a) If $A \in \mathcal{M}$ and $P(A) = 0$, then $\varphi(A) \leq 0$. (b) If $A \in \mathcal{M}^c$ and $P(A) = 0$, then $\varphi(A) \geq 0$.

THEOREM 5. *The following are equivalent:*

- (a) φ is absolutely continuous with respect to P on \mathcal{M} .
- (b) $\varphi(D(\varphi, \mathcal{M}) = \infty) = \varphi(D(\varphi, \mathcal{M}^c) = -\infty) = 0$
- (c) $\lim_{a \rightarrow \infty} \varphi_a^+ = \lim_{b \rightarrow -\infty} \varphi_b^- = 0$
- (d) There is an $X \in \mathcal{M}$ such that

$$\begin{aligned} \varphi(A) &\geq E(X; A) && A \in \mathcal{M}^c \\ &\leq E(X, A) && A \in \mathcal{M}. \end{aligned}$$

Proof. Let $X = D(\varphi, \mathcal{M})$. Then since

$$aP(a \leq X) \leq \varphi(a \leq X < \infty)$$

we always have $\lim_{a \rightarrow \infty} aP(a \leq X) = 0$. Now we note

$$\begin{aligned} 0 \leq \varphi_a^+ &= \varphi_a(X \geq a) \\ &= \varphi(X \geq a) - aP(X \geq a). \end{aligned}$$

Hence

$$\lim_{a \rightarrow \infty} \varphi_a^+ = \varphi(X = \infty).$$

Similarly,

$$\lim_{b \rightarrow -\infty} \varphi_b^- = \varphi(X = -\infty).$$

Hence, (b) and (c) are equivalent.

Now assume (a). Since $P(X = \infty) = 0$, it follows that $\varphi(X = \infty) \leq 0$. But since $\varphi(X \geq n) \geq nP(X \geq n) \geq 0$, $\varphi(X = \infty) \geq 0$ always holds. Hence $\varphi(X = \infty) = 0$. Similarly we can prove $\varphi(X = -\infty) = 0$, hence (b) holds.

Assume (c). Select $A \in \mathcal{M}$, and suppose $P(A) = 0$. Then for all a ,

$$\varphi(A) \leq \varphi_a^+ + aP(A) = \varphi_a^+$$

hence $\varphi(A) \leq 0$. In a similar way we can prove $\varphi(A) \geq 0$ if $A \in \mathcal{M}^c$ and $P(A) = 0$, so (c) implies (a).

Clearly (d) implies (a). Conversely, if (b) holds, then (d) follows from Theorem 4 by letting $a \rightarrow -\infty$, $b \rightarrow \infty$. This completes the proof.

EXAMPLE. Let (Ω, \mathcal{F}, P) denote the unit interval under Lebesgue measure. Let \mathcal{M} denote the class of all intervals of Ω , each of which contains the number 1. Define $\varphi(\{1\}) = -1$, $\varphi(\{0\}) = 1$, and $\varphi(A) = 0$ if $\{0, 1\} \cap A = \emptyset$. Then φ is absolutely continuous with respect to P on \mathcal{M} , but is singular with respect to P on \mathcal{F} . It is therefore not the case that there exists a function $X \in \mathcal{F}$ such that $\varphi(A) = E(X; A)$ for all $A \in \mathcal{F}$. It is not hard to verify that $D(\varphi, \mathcal{M}) = 0$ a.s.

In what follows, let \mathcal{M} be a fixed σ -lattice, and let \mathcal{D} denote the class of derivatives $X_0 = D(\varphi, \mathcal{M}_0)$, where $\mathcal{M}_0 \subseteq \mathcal{M}$. For each $X_0 \in \mathcal{D}$, define $\varphi_a^\pm(X_0)$ by (1) using \mathcal{M}_0 . Then

$$\begin{aligned} \varphi_a^+(X_0) &= \varphi_a(X_0 \geq a) \\ \varphi_a^-(X_0) &= \varphi_a(X_0 \leq a) \end{aligned}$$

and from the definition of φ^+ , it is clear that $\varphi_a^+(X_0) \leq \varphi_a^+(X)$ and $\varphi_a^-(X_0) \geq \varphi_a^-(X)$, where $X = D(\varphi, \mathcal{M})$. We now prove our main result:

THEOREM 6. \mathcal{D} is a uniformly integrable class if and only if φ is absolutely continuous with respect to P on \mathcal{M} .

Proof. For $X_0 \in \mathcal{D}$, for $a > 0$,

$$E(|X_0|; |X_0| \geq a) = E(X_0; X_0 \geq a) - E(X_0; X_0 \leq -a).$$

Since $\varphi(X_0 = \infty) = \lim_{n \rightarrow \infty} \varphi(X_0 \geq n) \geq 0$, we obtain from Theorem 4,

$$E(X_0; X_0 \geq a) \leq \varphi(\infty > X_0 \geq a) \leq \varphi(X_0 \geq a).$$

Now,

$$\begin{aligned} \varphi(X_0 \geq a) &= \varphi(X_0 \geq a, X \geq b) + \varphi(X_0 \geq a, X < b) \\ &\leq \varphi(X_0 \geq a, X \geq b) - bP(X_0 \geq a, X \geq b) \\ &\quad + bP(X_0 \geq a, X \geq b) + bP(X_0 \geq a, \\ &\quad X < b) \leq \varphi_b^+(X) + bP(X_0 \geq a). \end{aligned}$$

Now,

$$aP(X_0 \geq a) \leq \varphi(X_0 \geq a) \leq \varphi_b^+(X_0) \leq \varphi_b^+(X)$$

hence,

$$(6) \quad E(X_0, \{X_0 \geq a\}) \leq \varphi_b^+(X) + \frac{b}{a} \varphi_b^+(X).$$

If φ is absolutely continuous with respect to P on \mathcal{M} , then $\lim_{b \rightarrow \infty} \varphi_b^+(X) = 0$, and we can get a uniform bound on $E(X_0, \{X_0 \geq a\})$ for large enough a . In a similar fashion, we can prove $-E(X_0; \{X_0 \leq -a\})$ is uniformly bounded if φ is absolutely continuous with respect to P on \mathcal{M} , proving sufficiency of this condition.

Now if \mathcal{D} is not uniformly integrable, there is an $\varepsilon > 0$ such that for any $a > 0$, there is an $X_0 \in \mathcal{D}$ such that $E(|X_0|, |X_0| > a) > \varepsilon$. Without loss of generality, we can assume this statement holds for the left hand side of (6), hence $\lim_{b \rightarrow \infty} \varphi_b^+(X) > 0$, so φ is not absolutely continuous with respect to D on \mathcal{M} . This completes our proof.

Application. If $\mathcal{M}_1 \subseteq \mathcal{M}_2 \subseteq \dots$ are σ -lattices, it is shown in [1] that $D(\varphi, \mathcal{M}_n) \rightarrow D(\varphi, \mathcal{M})$ a.s., where \mathcal{M} is the minimal σ -lattice containing each \mathcal{M}_n , $n \geq 1$. It follows from Theorem 6 that if φ is absolutely continuous with respect to D on \mathcal{M} , this convergence holds in the mean as well. This strengthens a result in [2], where mean convergence of such a sequence is shown to hold when φ is absolutely continuous with respect to P on \mathcal{F} . However, our example above shows this is not the most general situation.

4. Jensen's inequality. Suppose φ is a absolutely continuous with respect to P on \mathcal{F} , so that there is a $Y \in \mathcal{F}$ such that $\varphi(A) = E(Y; A)$, $A \in \mathcal{F}$. Let $\mathcal{M} \subseteq \mathcal{F}$ be a σ -lattice. We define

$$E(Y | \mathcal{M}) = D(\varphi, \mathcal{M}).$$

and call $E(Y | \mathcal{M})$ the conditional expectation of Y given \mathcal{M} .

THEOREM 7. *Let F be a nondecreasing, convex function. If $F(Y)$ is integrable, then*

$$F(E(Y | \mathcal{M})) \leq E(F(Y) | \mathcal{M}) .$$

If G is a nonincreasing convex function, and $G(Y)$ is integrable, then

$$G(E(Y | \mathcal{M})) \leq E(G(Y) | \mathcal{M}^c) .$$

Proof. We prove the second assertion—the first being similar. Let $\{a_n x + b_n; n \geq 1\}$ be an affine lower envelope for G , with $a_n \leq 0$ all n . From Lemma 3,

$$a_n E(Y | \mathcal{M}) + b_n = E(a_n Y + b_n | \mathcal{M}^c)$$

and since $a_n Y + b_n \leq G(Y)$, it follows from Lemma 2 that

$$a_n E(Y | \mathcal{M}) + b_n \leq E(G(Y) | \mathcal{M}^c) .$$

Our result follows by taking the supremum of the left hand side.

Suppose H is a nonnegative convex function, with $H(0) = 0$. We say $X \in \mathcal{L}(H)$ if $X \in \mathcal{F}$ and $E(H(X)) < \infty$. If we apply Theorem 7 to the functions

$$\begin{aligned} F(x) &= H(x) & x \geq 0 \\ &= 0 & x < 0 \\ G(x) &= H(x) - F(x) \end{aligned}$$

then the first assertion of the following theorem clearly holds. The second assertion is implied by Theorem 6.

THEOREM 8. If $X \in \mathcal{L}(H)$, then so is $E(X | \mathcal{M})$. If $X \in \mathcal{L}(H)$, then the collection $\{H(E(X | \mathcal{M}_0))\}$, as \mathcal{M}_0 runs over the σ -lattices in \mathcal{F} , is uniformly integrable.

Integrability of conditional expectation $E(X | \mathcal{M}_0)$, and mean convergence of martingales of form $E(X | \mathcal{M}_n)$, where $\mathcal{M}_1 \subseteq \mathcal{M}_2 \subseteq \dots$, can be obtained Theorems 7 and 8, thus obtaining results in [2].

The author expresses his gratitude to the referee for correcting several blunders, and suggesting a simpler proof of Theorem 1.

REFERENCES

1. H. D. Brunk and S. Johansen, *A generalized Radon-Nikodym derivative*, Pacific J. Math., **34** (1970), 585-617.
2. R. B. Darst and G. A. DeBoth, *Norm convergence of Radon-Nikodym derivatives given σ -lattices*, To appear, Pacific J. Math.
3. S. Johansen, *The descriptive approach to the derivative of a set function with respect to a σ -lattice*, Pacific J. Math., **21** (1967), 49-58.

4. J. Neveu, *Mathematical Foundations of the Calculus of Probability*, Holden-Day, Inc., San Francisco, 1965.

Received January 10, 1972 and in revised form April 27, 1972. Supported by NSF Grant GP-24239.

UNIVERSITY OF SOUTHERN CALIFORNIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 46, No. 1

November, 1973

Allan Francis Abrahamse, <i>Uniform integrability of derivatives on σ-lattices</i>	1
Ronald Alter and K. K. Kubota, <i>The diophantine equation $x^2 + D = p^n$</i>	11
Grahame Bennett, <i>Some inclusion theorems for sequence spaces</i>	17
William Cutler, <i>On extending isotopies</i>	31
Robert Jay Daverman, <i>Factored codimension one cells in Euclidean n-space</i>	37
Patrick Barry Eberlein and Barrett O'Neill, <i>Visibility manifolds</i>	45
M. Edelstein, <i>Concerning dentability</i>	111
Edward Graham Evans, Jr., <i>Krull-Schmidt and cancellation over local rings</i>	115
C. D. Feustel, <i>A generalization of Kneser's conjecture</i>	123
Avner Friedman, <i>Uniqueness for the Cauchy problem for degenerate parabolic equations</i>	131
David Golber, <i>The cohomological description of a torus action</i>	149
Alain Gouillet de Rugy, <i>Un théorème du genre "Andô-Edwards" pour les Fréchet ordonnés normaux</i>	155
Louise Hay, <i>The class of recursively enumerable subsets of a recursively enumerable set</i>	167
John Paul Helm, Albert Ronald da Silva Meyer and Paul Ruel Young, <i>On orders of translations and enumerations</i>	185
Julien O. Hennefeld, <i>A decomposition for $B(X)^*$ and unique Hahn-Banach extensions</i>	197
Gordon G. Johnson, <i>Moment sequences in Hilbert space</i>	201
Thomas Rollin Kramer, <i>A note on countably subparacompact spaces</i>	209
Yves A. Lequain, <i>Differential simplicity and extensions of a derivation</i>	215
Peter Lorimer, <i>A property of the groups $\text{Aut PU}(3, q^2)$</i>	225
Yasou Matsugu, <i>The Levi problem for a product manifold</i>	231
John M.F. O'Connell, <i>Real parts of uniform algebras</i>	235
William Lindall Paschke, <i>A factorable Banach algebra without bounded approximate unit</i>	249
Ronald Joel Rudman, <i>On the fundamental unit of a purely cubic field</i>	253
Tsuan Wu Ting, <i>Torsional rigidities in the elastic-plastic torsion of simply connected cylindrical bars</i>	257
Philip C. Tonne, <i>Matrix representations for linear transformations on analytic sequences</i>	269
Jung-Hsien Tsai, <i>On E-compact spaces and generalizations of perfect mappings</i>	275
Alfons Van Daele, <i>The upper envelope of invariant functionals majorized by an invariant weight</i>	283
Giulio Varsi, <i>The multidimensional content of the frustum of the simplex</i>	303