

# Pacific Journal of Mathematics

**ON REALIZING HNN GROUPS IN 3-MANIFOLDS**

C. D. FEUSTEL AND ROBERT JOHN GREGORAC

## ON REALIZING HNN GROUPS IN 3-MANIFOLDS

C. D. FEUSTEL AND R. J. GREGORAC

**In this paper we suppose that the fundamental group of a 3-manifold  $M$  has a presentation as an HNN group. We then show that under suitable conditions we can realize this presentation by embedding a closed, connected incompressible surface in  $M$ .**

In [2], [3], and [4] we show that if  $\pi_1(M^3)$  is constructed in certain ways, one can realize this construction by a surface embedded in  $M^3$ . In this paper we show that one can realize the HNN construction when certain relationships between  $\pi_1(M^3)$  and  $M^3$  are present. The results in this paper are related to Theorem 2.4 in [10].

In this paper all spaces will be simplicial complexes, all maps will be piecewise linear, and all 3-manifolds will be 3-manifolds with boundary. However the boundary may be vacuous. Let  $X$  be a connected subspace of a space  $Y$ . As usual we shall denote the boundary, closure, and interior of  $X$  in  $Y$  by  $\text{bd}(X)$ ,  $\text{cl}(X)$ , and  $\text{int}(X)$  respectively. The natural inclusion map from  $X$  into  $Y$  will be denoted by  $\rho$  and the induced homomorphism from  $\pi_1(X)$  into  $\pi_1(Y)$  by  $\rho_*$ . Let  $S$  be a closed connected surface other than the 2-sphere of projective plane embedded in a space  $Y$ . Then  $S$  is *incompressible* in  $Y$  if  $\rho_*: \pi_1(S) \rightarrow \pi_1(Y)$  is one-to-one. If  $S$  is a closed surface embedded in  $Y$ , then  $S$  is *incompressible* in  $Y$  if each component of  $S$  is incompressible in  $Y$ . Irreducible and  $P^2$ -irreducible are defined as in [7]. We denote the unit interval  $[0, 1]$  by  $I$  throughout.

**DEFINITION 1.** Let  $K$  be a group and  $A$  a subgroup of  $K$ . Let  $S$  be a closed connected surface other than the projective plane or 2-sphere. Let  $A_j \cong \pi_1(S)$  and  $A_j \subset A$  for  $j = 1, 2$ . Let  $k$  be an element of  $K$  not in  $A$  such that  $A_1 = k^{-1}A_2k$ . Then if  $A$  and  $k$  generate  $K$  and all relations of  $K$  are consequences of the relations of  $A$  together with the relations  $k$  induces between the elements of  $A_1$  and  $A_2$ , we shall say that  $K$  is an *extension of  $A$  by  $k$  across  $A_1$  and  $A_2$* . The reader will note that the class of groups defined above is a subclass of the Higmann, Neumann, Neumann (H.N.N.) groups [8].

Let  $M$  be a 3-manifold,  $x$  a point in  $M$ , and  $S$  an incompressible surface in  $M$  such that  $M - S$  is connected. Then it is a consequence of Van Kampen's Theorem that  $\pi_1(M, x)$  is an extension of  $\pi_1(M - S, x)$  by some element of  $\pi_1(M, x)$  across appropriate subgroups of  $\pi_1(M, x)$ . One might then wonder "If  $\pi_1(M, x)$  were such an extension, could we embed in  $M$  an incompressible surface which realizes this exten-

sion." We will show below that this can, in fact, be done. Let  $M$  be a compact 3-manifold and  $x$  a point of  $M$ . We suppose that  $\pi_1(M, x)$  is an extension of  $A$  by  $k$  across  $A_1$  and  $A_2$  as given in Definition 1 above. We can represent this extension by an ordered sequence  $\langle \pi_1(M, x), A, A_1, A_2, k \rangle$ . If for each component  $F$  of the boundary of  $M$  some conjugate  $\rho_*\pi_1(F)$  is contained in  $A$ , we shall say that the extension preserves the peripheral structure of  $M$ . Suppose a second representation of  $\pi_1(M, x)$  is given by  $\langle \pi_1(M, x), B, B_1, B_2, \hat{k} \rangle$  and this extension of  $B$  is induced by an incompressible, closed, two-sided surface  $S$  embedded in  $M$  and a loop  $l$  meeting  $S$  in the single point  $x$ , i.e.,  $B$  is generated by the elements of  $\pi_1(M, x)$  having representative loops which do not cross  $S$ ,  $\hat{k} = [l]$ ,  $B_1 = \rho_*\pi_1(S, x)$  and  $B_2 = [l]B_1[l]^{-1}$ . We shall say that  $S$  realizes the extension of  $B$  if there is an isomorphism

$$\Phi: \pi_1(M, x) \longrightarrow \pi_1(M, x)$$

such that

- (1)  $\Phi(A) = B$
- (2)  $\Phi(A_j) = B_j \quad j = 1, 2$
- (3)  $\Phi(k) = \hat{k}$ .

**THEOREM 1.** *Let  $M$  be a compact 3-manifold such that  $\pi_2(M) = 0$ . Let  $S$  be a closed connected surface other than the 2-sphere or projective plane. Suppose  $\pi_1(M, x)$  has a representation given by*

$$\langle \pi_1(M, x), A, A_1, A_2, k \rangle$$

where  $A_i \cong \pi_1(S)$  and the extension above preserves the peripheral structure of  $M$ . Then there is an embedding of  $S$  in  $M$  which realizes the given extension.

The proof of Theorem 1 above is similar in many respects to the proof of Theorem 1 in [3]. One first constructs a complex  $X$  having the same fundamental group as  $M$ . One then finds a map  $f: M \rightarrow X$  inducing an isomorphism from  $\pi_1(M)$  to  $\pi_1(X)$ . The complex  $X$  is constructed to contain an embedded surface  $S$  realizing the given extension. One shows that there is a map  $g$  homotopic to  $f$  such that  $g^{-1}(S)$  is an incompressible, connected, closed surface in  $M$  and that  $g^{-1}(S)$  realizes the given extension.

The following three lemmas appear in [4]. We omit the proofs which are not difficult.

**LEMMA 1.** *Let  $M$  be a compact, connected 3-manifold such that  $\pi_2(M) = 0$ . Let  $X$  be a connected complex and  $S$  a closed incompressi-*

ble surface embedded in  $X$  and having a neighborhood homeomorphic to  $S \times I$ . We suppose that no component of  $S$  is a 2-sphere or projective plane. Let  $X_k, k = 1, \dots, n$  be the components of  $X - S$ . We suppose that  $\pi_i(X) = \pi_i(X_k) = 0$  for  $i \geq 2$  and  $k = 1, \dots, n$ . Let  $f: M \rightarrow X$  be a map such that  $f_*: \pi_1(M) \rightarrow \pi_1(X)$  is one-to-one  $f \text{ bd}(M)$  does not meet  $S$ . Then there is a homotopy, constant on  $\text{bd}(M)$ , of  $f$  to a map  $g$  such that  $g^{-1}(S)$  is an incompressible surface in  $M$ .

LEMMA 2. Let  $S_1$  and  $S_2$  be disjoint, incompressible, connected, two-sided surfaces which are embedded in a  $P^2$ -irreducible 3-manifold  $M$ . Then if  $S_1$  is homotopic to  $S_2$  in  $M$ ,  $S_1 \cup S_2$  bounds an  $S_1 \times I$  embedded in  $M$ .

LEMMA 3. Let  $M_1$  be a compact, connected, 3-manifold,  $X$  a connected complex, and  $F$  and  $S$  incompressible connected surfaces in  $M_1$  and  $X$  respectively. We suppose that  $S$  is neither a 2-sphere or projective plane and  $\pi_i(X) = 0$  for  $i \geq 2$ .

Let  $f: (M_1, F) \rightarrow (X, S)$  be a map of pairs such that for some  $x \in F$

$$f_*\pi_1(M_1, x) \subset \pi_1(S, f(x)) .$$

Then  $f$  is homotopic under a deformation, constant on  $F$ , to a map into  $S$ .

*Proof of Theorem 1.* It is a consequence of Remark 1 in [9] that we may assume that  $M$  is irreducible.

Let  $(M_A, \hat{x}, p)$  be the covering space of  $(M, x)$  associated with  $A \subset \pi_1(M, x)$ . Let  $f_1, f_2: (S, y) \rightarrow (M, x)$  be maps such that  $f_{j*}(\pi_1(S, y)) = A_j$ , for  $j = 1, 2$ . Since  $f_{j*}(\pi_1(S, y)) \subset p_*\pi_1(M_A, \hat{x})$ , there is a map  $\hat{f}_j: (S, y) \rightarrow (M_A, \hat{x})$  such that  $p\hat{f}_j = f_j$  for  $j = 1, 2$ . Let  $X$  be the union of  $M_A$  and  $S \times I$  with identifications  $\hat{f}_1(s) = (s, 0)$  and  $\hat{f}_2(s) = (s, 1)$ . We note that the arc  $\{y\} \times [0, 1] \subset S \times I$  becomes a simple loop  $\hat{l}$  after the identification above since  $\hat{f}_1(y) = \hat{f}_2(y) = \hat{x}$ . Let  $\Phi: A \cup \{k\} \rightarrow \pi_1(X, \hat{x})$  be a function defined by  $\Phi(k) = [l]$  and  $\Phi(a) = P_*^{-1}(a)$  for  $a \in A$ . Then  $\Phi$  can be extended to an isomorphism of  $\pi_1(M, x)$  onto  $\pi_1(X, \hat{x})$  since  $X$  has been constructed so that  $\pi_1(X, \hat{x})$  will have a presentation identical to the given presentation of  $\pi_1(M, x)$ .

It can be shown as in the proof of the theorem in [2] that  $\pi_i(X) = \pi_i(X - S) = 0$  for  $i \geq 2$ .

We denote  $S \times \{1/2\} \subset X$  by  $S$ .

Let the boundary of  $M$  be expressed as  $\bigcup_{m=1}^n F_m$  where  $F_m$  is a closed connected 2-manifold. Then some conjugate of  $\rho_*\pi_1(F_m)$  is contained in  $A$  for  $m = 1, \dots, n$ . Thus we can find a collection  $\{\alpha_m | m = 1, \dots, n\}$  of simple arcs embedded in  $M$  such that intersec-

tion of each pair of these arcs is  $x$ ,  $\alpha_m$  meets  $F_m$  in a single point, and there is a map  $\hat{\rho}: \mathbf{U}_{m=1}^n (F_m \cup \alpha_m) \rightarrow M_A$  such that  $p\hat{\rho} = \rho$ . Note that for each loop  $l_0$  in  $\mathbf{U}_{m=1}^n (F_m \cup \alpha_m)$  based at  $x$ ,  $[\hat{\rho}l_0] = \Phi[l_0]$ . Since  $\hat{\rho}_*\rho_* = \Phi\rho_*: \pi_1(\mathbf{U}_{m=1}^n (F_m \cup \alpha_m), x) \rightarrow \pi_1(X, \hat{x})$ , we can extend  $\hat{\rho}$  to a map  $f: M \rightarrow X$  such that  $\Phi = f_*: \pi_1(M, x) \rightarrow \pi_1(X, \hat{x})$  by using standard techniques from obstruction theory. (See [2] or [3] for the details of this construction.) It is a consequence of Lemma 1 that there is a map  $g_1$  homotopic to  $f$  such that  $g_1^{-1}(S)$  is an incompressible surface in  $M$  and  $g_1 = f$  on the boundary of  $M$ .

Since  $g_1^{-1}(S)$  and  $S$  are incompressible in  $M$  and  $X$  respectively, if  $S_0$  is any component of  $g_1^{-1}(S)$ , the homomorphism  $(g_1|S_0)_*: \pi_1(S_0) \rightarrow \pi_1(S)$  is one-to-one. Thus by Theorem 1 in [6]  $g_1|S_0$  is homotopic to a covering map. Thus after a deformation, constant outside of a small neighborhood of  $S_0$ , we may assume that  $g_1|S_0$  is a local homeomorphism. Thus we may assume that  $g_1$  is a local homeomorphism on  $g_1^{-1}(S)$ .

Let  $z$  be a point on  $S_0$ . Suppose that the isomorphism  $\Phi_0 = g_{1*}: \pi_1(M, z) \rightarrow \pi_1(X, g_1(z))$  does not carry  $\pi_1(S_0, z)$  onto  $\pi_1(S, g_1(z))$ . It is a consequence of the result in [1] that  $M$  is  $P^2$ -irreducible. Since  $\Phi_0^{-1}\pi_1(S, g_1(z))$  would properly contain  $\pi_1(S_0, z)$ , we would have by Theorem 6 in [7] that  $S_0$  bounds a twisted line bundle  $N \subset M$ . One can easily show using the techniques of [7], as has been done in [5], that  $\rho_*\pi_1(N, z)$  may be taken to be  $\Phi_0^{-1}(\rho_*\pi_1(S, g_1(z)))$ . It follows from Lemma 3 that there is a deformation of  $g_1$  to a map  $g_2$  which pushes  $g_1(N)$  first onto  $S$  and then to one side of  $S$  so that  $g_2^{-1}(S) = g_1^{-1}(S) - S_0$ . Thus we can assume that  $(g_1|S_0)_*: \pi_1(S_0) \rightarrow \pi_1(S)$  is an epimorphism for each component  $S_0$  of  $g_1^{-1}(S)$ .

Since  $\pi_1(M) \not\subset A$ ,  $g_1^{-1}(S)$  is not empty.

Let  $S_0$  and  $S_1$  be components of  $g_1^{-1}(S)$ . We claim that  $S_0 \cup S_1$  bounds a copy of  $S_0 \times [0, 1]$  embedded in  $M$ . Since  $M$  is  $P^2$ -irreducible, this will follow from Lemma 2 after we show that  $S_0$  and  $S_1$  are homotopic. Let  $z_0$  be a point on  $S_0$ . Since  $g_1|S_0$  and  $g_1|S_1$  are assumed to be homeomorphisms, there is a unique point  $z_1$  on  $S_1$  such that  $g_1(z_0) = g_1(z_1)$ . Let  $\alpha$  be an arc running from  $z_0$  to  $z_1$ . Since  $g_{1*}$  is an isomorphism, we can find a loop  $l_1$  based at  $z_0$  such that the loops  $g_1(l_1)$  and  $g_1(\alpha)$  represent the same element in  $\pi_1(X, g_1(z_0))$ . Thus we may assume that  $[g_1(\alpha)] = 1 \in \pi_1(X)$ . Let  $\lambda_0$  be a loop on  $S_0$  based at  $z_0$  and  $\lambda_1$  a loop on  $S_1$  such that  $g_1(\lambda_0) = g_1(\lambda_1)$ . Since the loop  $g_1(\lambda_0)g_1(\alpha)(g_1(\lambda_1))^{-1}(g_1(\alpha))^{-1}$  is nullhomotopic and  $\pi_2(X) = 0$ , we can show as in the proof of Theorem 1 in [3] that  $S_0$  and  $S_1$  are homotopic. Our claim follows.

We wish to show that we may assume  $g_1^{-1}(S)$  contains exactly one component.

Suppose there is more than one component in  $g_1^{-1}(S)$  and that the

number of components of  $g_1^{-1}(S)$  cannot be decreased by a small deformation of  $g_1$ . Let  $l: S^1 \rightarrow M$  be a loop in  $M$  such that  $g_{1*}[l] = [\hat{l}]$ . We may assume that

(i)  $g_1(l)$  meets  $S$  since the intersection number of  $[\hat{l}]$  and  $S$  is one. Thus we can take our basepoint to lie on one of the surfaces in  $g_1^{-1}(S)$ .

(ii)  $l$  crosses  $g_1^{-1}(S)$  at each point in  $l \cap g_1^{-1}(S)$  and thus  $(g_1 l)^{-1}(S)$  is a finite set whose cardinality cannot be reduced.

(iii)  $g_1(l \cap g_1^{-1}(S))$  is a single point.

Let  $D$  be a disk and  $\beta_1$  and  $\beta_2$  arcs in the boundary of  $D$  such that  $\beta_1 \cap \beta_2 = \text{bd}(\beta_1)$ . Then we can define a map  $\gamma: D \rightarrow X$  such that  $\gamma(\beta_1)$  is the loop  $g_1 l(S^1)$  and  $\gamma(\beta_2)$  is the loop  $\hat{l}$ .

We wish to show that  $g_1^{-1}(S)$  may be taken to be homeomorphic to  $S$  (connected). Assume that  $g_1^{-1}(S)$  is not connected; then it has been shown that each pair of distinct surfaces in  $g_1^{-1}(S)$  bounds a copy of  $S \times I$  embedded in  $M$ . If this is the case, it is clear that  $l^{-1}g_1^{-1}(S)$  contains more than one point. Let  $\delta_1, \dots, \delta_v$  be the closures of the components of  $S^1 - l^{-1}g_1^{-1}(S)$ . After a general position argument we may assume  $\gamma^{-1}(S)$  contains an arc  $\beta_3$  which cuts off an arc  $\beta_4 \subset \beta_1$  and that  $g_1 l(\delta_1) = \gamma(\beta_4)$ . Now  $l$  carries  $\text{bd}(\delta_1)$  to one or two components of  $g_1^{-1}(S)$ .

If  $l(\text{bd}(\delta_1))$  is a single point, the loop  $l(\delta_1)$  is homotopic to a loop  $l_1 \subset g_1^{-1}(S)$  such that  $g_1(l_1) = \gamma(\beta_3)$  since the restriction of  $g_1$  to each component of  $g_1^{-1}(S)$  is a homeomorphism and  $g_{1*}$  is an isomorphism. It would follow that the number of points in  $l^{-1}g_1^{-1}(S)$  could have been reduced by a different choice of  $l$ . Thus we conclude that  $l$  carries the points of  $\text{bd}(\delta_1)$  to distinct components of  $g_1^{-1}(S)$ .

Let  $N$  be closure of the component of  $M - g_1^{-1}(S)$  which meets  $l(\delta_1)$ . Let  $S_0$  be a component of  $\text{bd}(N)$ . Since  $g_1|_{S_0}$  is a homeomorphism and the loop  $g_1 l(\delta_1)$  is homotopic to a loop in  $S$ , we may assume that the loop  $g_1 l(\delta_1)$  is homotopic to a point. (One alters the image of  $l$  in a neighborhood of  $S_0$ .)

Since the loop  $g_1 l(\delta_1)$  is nullhomotopic in  $X$ , it can be shown that the map  $g_1|_N$  is homotopic mod  $\text{bd}(N)$  to a map into  $S$ ; full details of a similar argument appear in [3]. It follows after an argument by induction that there exists a map  $g: M \rightarrow X$  homotopic to  $g_1$  mod  $\text{bd}(M)$  such that  $g^{-1}(S)$  contains exactly one component  $S_0$  and  $g|_{S_0}$  is a homeomorphism. After an argument similar to the one given above, we can find a loop  $l$  meeting  $S_0$  in a single point and based at  $x \in M$  such that  $g_*[l] = [\hat{l}]$ .

We observe that  $S_0$  and  $l$  induce an expression of  $\pi_1(M, x)$  as an extension of a subgroup  $B$  of  $\pi_1(M, x)$ . Let  $B_1$  and  $B_2$  be the associated subgroups of  $\pi_1(M, x)$ . Then we see that our map  $g$  induces

an isomorphism  $g_*: \pi_1(M, x) \rightarrow \pi_1(M, x)$  such that

- (1)  $g_*(B) \subset A$
- (2)  $g_*(B_i) = A_i$
- (3)  $g_*B_2 = A_2$ .

Thus Theorem 1 is an immediate consequence of the remark preceding Lemma 2 on page 238 in [8] which shows that  $g_*$  sends  $B$  onto  $A$ .

REMARK 1. The remark mentioned above allows us to strengthen the statement of the theorem in [2] so that the splitting and the cutting are both actually realized.

REMARK 2. We can also realize geometrically more general presentations of  $\pi_1(M)$  as an HNN group. In particular one might have that  $\pi_1(M)$  has a presentation as in the first definition in §4 in [8] where each of the subgroups  $L_i$  of  $K$  is isomorphic to the fundamental group of a closed connected surface other than  $S^2$  or the projective plane and there are only finitely many of the  $t_i$ . The proof of this result varies only slightly from the one given above.

REMARK 3. Theorem 1 in this paper together with Theorem 1 in [3] or [4] give us a sort of converse to Van Kampen's theorem as applied to a closed, connected, incompressible surface, other than  $S^2$  or the projective plane, embedded in the interior of a compact 3-manifold.

REMARK 4. This paper is in some sense a generalization of Stallings's work in [11].

#### REFERENCES

1. C. D. Feustel,  $M^3$  admitting a certain embedding of  $P^2$  is a pseudo  $P^3$ , Proc. Amer. Math. Soc., **26** (1970), 215-216.
2. ———, On replacing proper Dehn maps with proper embeddings, Trans. Amer. Math. Soc., **166** (1972), 261-267.
3. ———, A splitting theorem for closed orientable 3-manifolds, Topology, **11** (1972), 151-158.
4. ———, A generalization of Kneser's conjecture, Pacific J. Math., to appear.
5. ———, On  $S$ -maximal subgroups of  $\pi_1(M^2)$ , Canad. J. Math., **24** (1972), 439-449.
6. W. Heil, On  $P^2$ -irreducible 3-manifolds, Bull. Amer. Math. Soc., **75** (1969), 772-775.
7. W. Jaco, Finitely presented subgroups of three-manifold groups, Inventiones Math., **13** (1971), 335-346.
8. A. Karrass and D. Solitar, The subgroups of a free product of two groups with an amalgamated subgroups, Trans. Amer. Math. Soc., **150** (1970), 227-255.
9. J. Milnor, A unique decomposition theorem for 3-manifolds, Amer. J. Math., **84** (1962), 1-7.

10. G. P. Scott, *On sufficiently large 3-manifolds*, Quart. J. Math. Oxford, **23** (1972), 159-172.
11. J. Stallings, *On Fiberings Certain 3-Manifolds*, Topology of 3-manifolds, Prentice-Hall, Englewood Cliffs, N. J., 1962, 95-100.

Received April 28, 1972 and in revised form October 3, 1972.

VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY  
AND  
IOWA STATE UNIVERSITY





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

D. GILBARG AND J. MILGRAM

Stanford University  
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington  
Seattle, Washington 98105

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$48.00 a year (6 Vols., 12 issues). Special rate: \$24.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Christopher Allday, <i>Rational Whitehead products and a spectral sequence of Quillen</i> .....	313
James Edward Arnold, Jr., <i>Attaching Hurewicz fibrations with fiber preserving maps</i> .....	325
Catherine Bandle and Moshe Marcus, <i>Radial averaging transformations with various metrics</i> .....	337
David Wilmot Barnette, <i>A proof of the lower bound conjecture for convex polytopes</i> .....	349
Louis Harvey Blake, <i>Simple extensions of measures and the preservation of regularity of conditional probabilities</i> .....	355
James W. Cannon, <i>New proofs of Bing's approximation theorems for surfaces</i> .....	361
C. D. Feustel and Robert John Gregorac, <i>On realizing HNN groups in 3-manifolds</i> .....	381
Theodore William Gamelin, <i>Iversen's theorem and fiber algebras</i> .....	389
Daniel H. Gottlieb, <i>The total space of universal fibrations</i> .....	415
Yoshimitsu Hasegawa, <i>Integrability theorems for power series expansions of two variables</i> .....	419
Dean Robert Hickerson, <i>Length of period simple continued fraction expansion of <math>\sqrt{d}</math></i> .....	429
Herbert Meyer Kamowitz, <i>The spectra of endomorphisms of the disc algebra</i> .....	433
Dong S. Kim, <i>Boundedly holomorphic convex domains</i> .....	441
Daniel Ralph Lewis, <i>Integral operators on <math>\mathcal{L}_p</math>-spaces</i> .....	451
John Eldon Mack, <i>Fields of topological spaces</i> .....	457
V. B. Moscatelli, <i>On a problem of completion in bornology</i> .....	467
Ellen Elizabeth Reed, <i>Proximity convergence structures</i> .....	471
Ronald C. Rosier, <i>Dual spaces of certain vector sequence spaces</i> .....	487
Robert A. Rubin, <i>Absolutely torsion-free rings</i> .....	503
Leo Sario and Cecilia Wang, <i>Radial quasiharmonic functions</i> .....	515
James Henry Schmerl, <i>Peano models with many generic classes</i> .....	523
H. J. Schmidt, <i>The <math>\mathcal{F}</math>-depth of an <math>\mathcal{F}</math>-projector</i> .....	537
Edward Silverman, <i>Strong quasi-convexity</i> .....	549
Barry Simon, <i>Uniform crossnorms</i> .....	555
Surjeet Singh, <i>(KE)-domains</i> .....	561
Ted Joe Suffridge, <i>Starlike and convex maps in Banach spaces</i> .....	575
Milton Don Ulmer, <i>C-embedded <math>\Sigma</math>-spaces</i> .....	591
Wolmer Vasconcelos, <i>Conductor, projectivity and injectivity</i> .....	603
Hidenobu Yoshida, <i>On some generalizations of Meier's theorems</i> .....	609