

Pacific Journal of Mathematics

THE TOTAL SPACE OF UNIVERSAL FIBRATIONS

DANIEL H. GOTTLIEB

THE TOTAL SPACE OF UNIVERSAL FIBRATIONS

DANIEL HENRY GOTTLIEB

It is shown that the total space of a universal fibration for a fibre F is a classifying space for the monoid of self homotopy equivalences of F which fix the base point.

For any space F , there exists a universal Hurewicz fibration $F \rightarrow E_\infty \rightarrow B_\infty$, where B_∞ is a CW complex which classifies Hurewicz fibrations over CW complexes (See Dold [2], Theorem 16.9.). Now B_∞ is the Dold-Lashof classifying space for the monoid of self homotopy equivalence of F , which we shall denote by F^F . At least, this is the case when F is a CW complex. (See Allaud [1], § IV.) The purpose of this note is to show that E_∞ is the classifying space for the monoid of self equivalences of F leaving the base point fixed, denoted F_0^F , when F is a CW complex with homotopy equivalent path components. In fact, we shall show E_∞ is the base space of a Serre fibration with fibre F_0^F and a total space which is essentially contractible. We need this characterization of E_∞ in order to calculate the induced homomorphism on integral cohomology of the evaluation map $\omega: X^X \rightarrow X$ where $X = CP^n$. This is done in [3].

Let $D = p^*(E_\infty)$, the pullback of E_∞ by $p: E_\infty \rightarrow B_\infty$. Thus $D = \{(e, e') \in E_\infty \times E_\infty \mid p(e) = p(e')\}$, and $\bar{p}: D \rightarrow E_\infty$ given by $\bar{p}(e, e') = e$ is the projection. Let D_0^F be the set of maps of $F \rightarrow D$ endowed with the $C-0$ topology such that:

- (a) Each map carries F into some fibre of $D \xrightarrow{\bar{p}} E_\infty$ and is a homotopy equivalence of F and the fibre.
- (b) Each map carries the base point, $*$, into a point of the form (e, e) .

Let $q: D_0^F \rightarrow E_\infty$ be given by $q(f) = \bar{p} \circ f(*)$.

THEOREM.

- (1) q is a Serre Fibration, and if F is locally compact q is a Hurewicz fibration.
- (2) The fibre of q is F_0^F .
- (3) There is a fibrewise action $D_0^F \times F_0^F \rightarrow D_0^F$ if F is locally compact.
- (4) D_0^F is essentially contractible.

Proof of (1). First note that q is onto since all the components of F have the same homotopy type.

We shall assume that F has a whisker. That is, assume F has

the form $J \vee F'$ where J is the unit interval with the base point $*$ being the 1 and with $0 \in F'$. Every space is homotopy equivalent to a space of this type, so we do not lose any generality.

Let X be a compact polyhedron (or F is locally compact). We must show that p has the covering homotopy property with respect to any map $X \rightarrow D_0^F$. Since X is compact (or F is locally compact) $X \times F$ is a CW complex; so we may consider the adjoint map

$$f: X \times F \longrightarrow D ,$$

where f is a fibre preserving map which carries $X \times *$ into $\Delta \subset D$, where $\Delta = \{(e, e) \mid e \in E_\infty\}$. Then the covering homotopy property translates into a statement involving a fibre homotopy of f which at each stage sends $X \times *$ into Δ . Reflecting on the definition of D , we see that the covering homotopy property is equivalent to the following statement: Let $f: X \times F \rightarrow E_\infty$ be a fibre map and let $h_i: X \rightarrow E_\infty$ be any homotopy such that $h_0(x) = f(x, *)$ for all $x \in X$. Then there exists a fibre homotopy $\tilde{h}_i: X \times F \rightarrow E_\infty$ such that $\tilde{h}_i(x, *) = h_i(x)$ and $\tilde{h}_0 = f$.

Now this statement is a special case of the statement that q has the covering homotopy extension property for the space $X \times F$ relative to $X \times *$. See Hu, page 62 [4] for the definition. But this follows from Satz 5.38, page 107 in [5]. (The fact that $*$ is on the end of a whisker allows us to satisfy the technical requirements of Satz 5.38 concerning a halo about $X \times *$.)

Proof of (4). First note that (2) and (3) are obviously true. The action in (3), $D_0^F \times F_0^F \rightarrow D_0^F$ is given by $(g, f) \rightarrow g \circ f$. (This action is continuous if F is locally compact.)

Now we shall show that D_0^F is essentially contractible. That is, any map $X \rightarrow D_0^F$ is homotopy trivial if X is a finite CW complex. Consider the adjoint map $g: X \times F \rightarrow D$. Then g is defined by, and defines, a fibre map $h: X \times F \rightarrow E_\infty$ by means of the relation

$$g(x, y) = (h(x, *), h(x, y)) \in D .$$

Now h can be extended to a fibre map $H: CX \times F \rightarrow E_\infty$, (because E_∞^F is essentially contractible, [1] Theorem 4.1). We define a fibre map $G: CX \times F \rightarrow D$ by

$$G(x, y) = (H(x, *), H(x, y)) , \quad x \in CX, y \in F .$$

Note that G extends g . The adjoint situation now shows that our original map $X \rightarrow D_0^F$ factors through $X \rightarrow CX \rightarrow D_0^F$.

COROLLARY.

$$\pi_i(E_\infty) \cong \pi_{i-1}(F_0^F) .$$

This corollary plays an important role in the computation of the homomorphisms induced in cohomology by the evaluation map $\omega: F^F \rightarrow F$. The program is based on the use of the Federer spectral sequence and obstruction theory to compute some homotopy groups of F_0^F , and hence of E_∞ by the corollary. From the homotopy group information, we obtain information about the cohomology of E_∞ . Then we use the Serre exact sequence and the slogan “ ω^* factors through the transgression” to recover information about ω^* .

This program has been used successfully to compute ω^* for $H^2(CP^n; Z)$. See Theorem 16 of [3].

REFERENCES

1. Guy Allaud, *On the classification of fibre spaces*, Math. Z., **92** (1966), 110-125.
2. Albrecht Dold, *Halbexakte Homotopiefunktoren*, Lecture Notes in Mathematics, No. 12, Springer-Verlag, 1966, (see p. 16. 8).
3. Daniel Henry Gottlieb, *Witnesses, transgressions, and the evaluation map*, mimeographed, Purdue Univ.
4. S. T. Hu, *Homotopy Theory*, Academic Press, New York, (1959).
5. T. Tom Dieck, K. H. Kamps, and D. Puppe, *Homotopietheorie*, Lecture Notes in Mathematics, vol. **157**, Springer-Verlag, 1970.

Received March 27, 1972 and in revised form June 21, 1972. This research was partially supported by a grant from the National Science Foundation.

PURDUE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$48.00 a year (6 Vols., 12 issues). Special rate: \$24.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics

Vol. 46, No. 2

December, 1973

Christopher Allday, <i>Rational Whitehead products and a spectral sequence of Quillen</i>	313
James Edward Arnold, Jr., <i>Attaching Hurewicz fibrations with fiber preserving maps</i>	325
Catherine Bandle and Moshe Marcus, <i>Radial averaging transformations with various metrics</i>	337
David Wilmot Barnette, <i>A proof of the lower bound conjecture for convex polytopes</i>	349
Louis Harvey Blake, <i>Simple extensions of measures and the preservation of regularity of conditional probabilities</i>	355
James W. Cannon, <i>New proofs of Bing's approximation theorems for surfaces</i>	361
C. D. Feustel and Robert John Gregorac, <i>On realizing HNN groups in 3-manifolds</i>	381
Theodore William Gamelin, <i>Iversen's theorem and fiber algebras</i>	389
Daniel H. Gottlieb, <i>The total space of universal fibrations</i>	415
Yoshimitsu Hasegawa, <i>Integrability theorems for power series expansions of two variables</i>	419
Dean Robert Hickerson, <i>Length of period simple continued fraction expansion of \sqrt{d}</i>	429
Herbert Meyer Kamowitz, <i>The spectra of endomorphisms of the disc algebra</i>	433
Dong S. Kim, <i>Boundedly holomorphic convex domains</i>	441
Daniel Ralph Lewis, <i>Integral operators on \mathcal{L}_p-spaces</i>	451
John Eldon Mack, <i>Fields of topological spaces</i>	457
V. B. Moscatelli, <i>On a problem of completion in bornology</i>	467
Ellen Elizabeth Reed, <i>Proximity convergence structures</i>	471
Ronald C. Rosier, <i>Dual spaces of certain vector sequence spaces</i>	487
Robert A. Rubin, <i>Absolutely torsion-free rings</i>	503
Leo Sario and Cecilia Wang, <i>Radial quasiharmonic functions</i>	515
James Henry Schmerl, <i>Peano models with many generic classes</i>	523
H. J. Schmidt, <i>The \mathcal{F}-depth of an \mathcal{F}-projector</i>	537
Edward Silverman, <i>Strong quasi-convexity</i>	549
Barry Simon, <i>Uniform crossnorms</i>	555
Surjeet Singh, <i>(KE)-domains</i>	561
Ted Joe Suffridge, <i>Starlike and convex maps in Banach spaces</i>	575
Milton Don Ulmer, <i>C-embedded Σ-spaces</i>	591
Wolmer Vasconcelos, <i>Conductor, projectivity and injectivity</i>	603
Hideobu Yoshida, <i>On some generalizations of Meier's theorems</i>	609