

Pacific Journal of Mathematics

**CONTINUATION OF HOLOMORPHIC MAPPINGS, WITH
VALUES IN A COMPLEX LIE GROUP**

K. ADACHI, MASUO SUZUKI AND M. YOSHIDA

CONTINUATION OF HOLOMORPHIC MAPPINGS, WITH VALUES IN A COMPLEX LIE GROUP

K. ADACHI, M. SUZUKI AND M. YOSHIDA

In this paper it is shown that any holomorphic mapping of a domain X over a Stein manifold in a complex Lie group L can be continued to a holomorphic mapping of the envelope of holomorphy of X in L .

1. Let M be a complex manifold. A pair (X, ψ) of a complex manifold X and a locally biholomorphic mapping ψ of X in M is called an open set over M . Moreover, if X is connected, (X, ψ) is called a domain over M . Let (X, ψ) and (X', ψ') be open sets over M . A holomorphic mapping λ of X in X' with $\psi = \psi' \circ \lambda$ is called a mapping of (X, ψ) in (X', ψ') . Consider domains (X, ψ) and (X', ψ') over M with a mapping λ of (X, ψ) in (X', ψ') such that each connected component of X' contains that of $\lambda(X)$. Let Y be a complex manifold. Let f be a holomorphic mapping of X in Y . A holomorphic mapping f' of X' in Y with $f = f' \circ \lambda$ is called a holomorphic continuation of f to (λ, X', ψ') . Let \mathcal{F} be a family of holomorphic mappings of X in Y . If any holomorphic mapping f of \mathcal{F} has a holomorphic continuation to (λ, X', ψ') , (λ, X', ψ') is called a holomorphic completion of (X, ψ) with respect to the family \mathcal{F} . A holomorphic completion $(\tilde{\lambda}_{\mathcal{F}}, \tilde{X}_{\mathcal{F}}, \tilde{\psi}_{\mathcal{F}})$ of (X, ψ) with respect to \mathcal{F} is called an envelope of holomorphy of (X, ψ) with respect to the family \mathcal{F} if the following conditions are satisfied:

Let (λ', X', ψ') be another holomorphic completion of (X, ψ) with respect to \mathcal{F} . There is a mapping φ of (X', ψ') in $(\tilde{X}_{\mathcal{F}}, \tilde{\psi}_{\mathcal{F}})$ with $\tilde{\lambda}_{\mathcal{F}} = \varphi \circ \lambda'$ such that $(\varphi, \tilde{X}_{\mathcal{F}}, \tilde{\psi}_{\mathcal{F}})$ is a holomorphic completion of (X', ψ') with respect to the family \mathcal{F}' of holomorphic continuations to (λ', X', ψ') of all holomorphic mappings of \mathcal{F} .

If \mathcal{F} is the family of all holomorphic functions in X , the envelope of holomorphy of (X, ψ) with respect to \mathcal{F} is called the envelope of holomorphy of (X, ψ) . If \mathcal{F} consists of only one holomorphic mapping f of X in Y , the envelope of holomorphy of (X, ψ) with respect to the family \mathcal{F} is called the open set of holomorphy of f . Malgrange [5] proved the unique existence of an envelope of holomorphy, considering a connected component of the sheaf of all germs of families of holomorphic mappings with the same index set. By this construction of an envelope of holomorphy we have the following lemma.

LEMMA 1. *Let (X, ψ) be a domain over a complex manifold, Y be a complex manifold and \mathcal{F} be a family of holomorphic mappings of X in Y . There is uniquely an envelope $(\tilde{\lambda}_{\mathcal{F}}, \tilde{X}_{\mathcal{F}}, \tilde{\psi}_{\mathcal{F}})$ of holomorphy of (X, ψ) with respect to \mathcal{F} . Moreover, let \mathcal{G} be a subfamily of \mathcal{F} and $(\tilde{\lambda}_{\mathcal{G}}, \tilde{X}_{\mathcal{G}}, \tilde{\psi}_{\mathcal{G}})$ be the envelope of holomorphy of (X, ψ) with respect to \mathcal{G} . There is a mapping φ of $(\tilde{X}_{\mathcal{F}}, \tilde{\psi}_{\mathcal{F}})$ in $(\tilde{X}_{\mathcal{G}}, \tilde{\psi}_{\mathcal{G}})$ such that $(\varphi, \tilde{X}_{\mathcal{G}}, \tilde{\psi}_{\mathcal{G}})$ is the holomorphic completion of $(\tilde{X}_{\mathcal{F}}, \tilde{\psi}_{\mathcal{F}})$ with respect to \mathcal{G} .*

2. We put

$$(1) \quad \begin{aligned} D = \{z = (z_1, z_2, \dots, z_n) \in C^n; |z_1| < 1 + \varepsilon, |z_j| < 1 (j = 2, 3, \dots, n)\} \\ \cup \{z = (z_1, z_2, \dots, z_n) \in C^n; 1 - \varepsilon < |z_1| < 1 + \varepsilon, |z_j| < 1 + \varepsilon \\ (j = 2, 3, \dots, n)\} \end{aligned}$$

and

$$(2) \quad E = \{z = (z_1, z_2, \dots, z_n) \in C^n; |z_j| < 1 + \varepsilon (j = 1, 2, \dots, n)\}$$

for a positive number ε with $\varepsilon < 1$. E is the envelope of holomorphy of D .

LEMMA 2. *Let L be a complex Lie group. For any holomorphic mapping f of D in L , there is a holomorphic mapping g of E in L such that $g = f$ in D .*

Proof. We may assume that L is connected. Let \mathcal{H} be the set of all holomorphic mappings of D in L . We introduce in \mathcal{H} the compact-open topology. As D is analytically contractible to a point, \mathcal{H} is a connected topological group. We put

$$\begin{aligned} K(\delta) = \{z \in C^n; |z_1| \leq 1 + \varepsilon - \delta, |z_j| \leq 1 - \delta (j = 2, 3, \dots, n)\} \\ \cup \{z \in C^n; 1 - \varepsilon + \delta \leq |z_1| \leq 1 + \varepsilon - \delta, |z_j| \leq 1 + \varepsilon - \delta \\ (j = 2, 3, \dots, n)\} \end{aligned}$$

and

$$E(\delta) = \{z \in C^n; |z_j| < 1 + \varepsilon - \delta (j = 1, 2, \dots, n)\}$$

for any positive number δ with $\delta < \varepsilon$. Let m be the complex dimension of L and \exp be the exponential mapping of C^m in L . \exp maps an open neighborhood $U = \{w \in C^m; |w_j| < a (j = 1, 2, \dots, m)\}$ of the origin in C^m biholomorphically on an open neighborhood W of the unit element e of L . Then $\log = (\exp|U)^{-1}$ is a biholomorphic mapping of W onto U . We put $\mathcal{V}(1) = \{h \in \mathcal{H}; h(K(\delta)) \subset W\}$. Then $\mathcal{V}(1)$ is a neighborhood of the unit element 1 of the topological

group \mathcal{H} . Since \mathcal{H} is connected, \mathcal{H} is generated by $\mathcal{V}(1)$. There is a finite number s of elements f_1, f_2, \dots, f_{s-1} and f_s of $\mathcal{V}(1)$ such that $f = f_1 f_2 \cdots f_s$ in D . Each $\log f_i$ is a holomorphic mapping of $K(\delta)$ in the polydisc U . There is a holomorphic mapping G_i of $E(\delta)$ in U such that $G_i = \log f_i$ in $K(\delta) \cap E(\delta)$ for $j = 1, 2, \dots, s$. We put $g = \exp G_1 \exp G_2 \cdots \exp G_s$ in $E(\delta)$. Then g is a holomorphic mapping of $E(\delta)$ in L such that $g = f$ in $K(\delta) \cap E(\delta)$. Since δ is arbitrary, we have Lemma 2 by the theorem of identity.

Let (X, ψ) be a domain over a Stein manifold S , L be a complex Lie group, $\mathcal{F} = \{f_i; i \in I\}$ be a family of holomorphic mappings of X in L and (λ, X', ψ') be the envelope of holomorphy of (X, ψ) with respect to \mathcal{F} .

LEMMA 3. (X', ψ') is p_r -convex in the sense of Docquier-Grauert [2].

Proof. Assume that (X', ψ') were not p_r -convex. There is a continuous mapping φ of the closure \bar{D} of D , defined in (1) for a positive number ε with $\varepsilon < 1$, in $X' \cup \tilde{\delta}X'$ such that $\varphi(D) \subset X'$, $\varphi(b, c) \in \tilde{\delta}X'$ for $b \in C$ with $|b| \leq 1 - \varepsilon$, $c = (c_2, \dots, c_n) \in C^{n-1}$ with $|c_j| = 1$ ($j = 2, 3, \dots, n$) and $\psi' \circ \varphi$ can be continued to a biholomorphic mapping π of E , given in (2) for the above ε , in S . Here $\tilde{\delta}X'$ is the boundary of the domain (X', ψ') over S defined in § 3.1d of [2].

Let f'_i be a holomorphic continuation of $f_i \in \mathcal{F}$ to (λ, X', ψ') for any $i \in I$. By Lemma 2 there is a holomorphic mapping g_i of E in L such that $g_i = f'_i \circ \varphi$ in D for any $i \in I$.

We consider the sum space $X' \cup E$. Let x_1 and x_2 be, respectively, points of X' and E such that $\psi'(x_1) = \pi(x_2)$. Let U_1 and U_2 be, respectively, neighborhoods of x_1 and x_2 in X' and E such that $\psi'|U_1$ and $\pi|U_2$ are, respectively, biholomorphic mappings of U_1 and U_2 onto an open neighborhood V in S . We shall identify x_1 and x_2 if and only if $f'_i \circ (\psi'|U_1)^{-1} = g_i \circ (\pi|U_2)^{-1}$ for any $i \in I$. Let Z be the quotient space of $X' \cup E$ by this equivalence relation. Let $\xi: X' \rightarrow Z$ and $\eta: E \rightarrow Z$ be the canonical mappings. There is a locally topological mapping ζ of Z in S such that $\zeta \circ \xi = \psi'$ and $\zeta \circ \eta = \pi$. Since Z is a Hausdorff space, we can introduce in Z a complex structure such that (Z, ζ) is a domain over S .

There is a holomorphic mapping h_i of Z in L such that $h_i \circ \xi = f'_i$ and $h_i \circ \eta = g_i$ for any $i \in I$. Then h_i is a continuation of f'_i to (ξ, Z, ζ) . Since (λ, X', ψ') is the envelope of holomorphy of X with respect to \mathcal{F} , there is a mapping μ of (Z, ζ) in (X', ψ') . Since $\zeta \circ \xi = \psi'$, $\xi \circ \mu$ and $\mu \circ \xi$ are, respectively, the identities of Z and X' . Hence ξ is a biholomorphic mapping of X' on Z . Since $(b, c) \in E$, we

have $\varphi(b, c) = (\xi^{-1} \circ \eta)(b, c) \in X'$. This is a contradiction.

3. Let (X, ψ) be a domain over a Stein manifold S , $(\tilde{\lambda}, \tilde{X}, \tilde{\psi})$ be its envelope of holomorphy, L be a complex Lie group and f be a holomorphic mapping of X in L . Let (λ, X', ψ') be the open set of holomorphy of f . By Lemma 3 and a theorem of Docquier-Grauert [2], X' is a Stein manifold. Hence (X', ψ') is a domain of holomorphy of a holomorphic function in X . Since $(\tilde{\lambda}, \tilde{X}, \tilde{\psi})$ is the envelope of holomorphy with respect to the family of all holomorphic functions in X , there is a mapping μ of $(\tilde{X}, \tilde{\psi})$ in (X', ψ') such that $\lambda = \mu \circ \tilde{\lambda}$ by Lemma 1. Let f' be the holomorphic continuation of f to (λ, X', ψ') . Then $f' \circ \mu$ is the holomorphic continuation of f to $(\tilde{\lambda}, \tilde{X}, \tilde{\psi})$. Thus we have proved the following theorem.

THEOREM. *Let (X, ψ) be a domain over a Stein manifold S and $(\tilde{\lambda}, \tilde{X}, \tilde{\psi})$ be its envelope of holomorphy. Any holomorphic mapping of X in a complex Lie group L has a holomorphic continuation to $(\tilde{\lambda}, \tilde{X}, \tilde{\psi})$.*

REFERENCES

1. H. Cartan und P. Thullen, *Zur Theorie der Singularitäten der Funktionen mehrerer komplexer Veränderlichen, Regularitäts- und Konvergenzbereiche*, Math. Ann., **106** (1932), 617–647.
2. F. Docquier und H. Grauert, *Levisches Problem und Rungescher Satz für Teilgebiete Steinscher Mannigfaltigkeiten*, Math. Ann., **140** (1960), 94–123.
3. J. Kajiwara, *On the limit of a monotonous sequence of Cousin's domain*, J. Math. Soc. Japan, **17** (1965), 36–46.
4. J. Kajiwara and E. Sakai, *Generalization of Levi-Oka's theorem concerning meromorphic functions*, Nagoya Math. J., **29** (1967), 75–84.
5. B. Malgrange, *Lectures on the theory of functions of several complex variables*, Tata. Inst. Bombay (1958).

Received March 6, 1972 and revised form May 25, 1972.

IBARAKI UNIVERSITY
 TOYAMA UNIVERSITY
 AND
 FUKUOKA UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

* C. DePrima will replace J. Dugundji until August 1974.

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

K. Adachi, Masuo Suzuki and M. Yoshida, <i>Continuation of holomorphic mappings, with values in a complex Lie group</i>	1
Michael Aschbacher, <i>A characterization of the unitary and symplectic groups over finite fields of characteristic at least 5</i>	5
Larry Eugene Bobisud and James Calvert, <i>Energy bounds and virial theorems for abstract wave equations</i>	27
Christer Borell, <i>A note on an inequality for rearrangements</i>	39
Peter Southcott Bullen and S. N. Mukhopadhyay, <i>Peano derivatives and general integrals</i>	43
Wendell Dan Curtis, Yu-Lee Lee and Forrest Miller, <i>A class of infinite dimensional subgroups of $\text{Diff}^r(X)$ which are Banach Lie groups</i>	59
Paul C. Eklof, <i>The structure of ultraproducts of abelian groups</i>	67
William Alan Feldman, <i>Axioms of countability and the algebra $C(X)$</i>	81
Jack Tilden Goodykoontz, Jr., <i>Aposyndetic properties of hyperspaces</i>	91
George Grätzer and J. Płonka, <i>On the number of polynomials of an idempotent algebra. II</i>	99
Alan Trinler Huckleberry, <i>The weak envelope of holomorphy for algebras of holomorphic functions</i>	115
John Joseph Hutchinson and Julius Martin Zelmanowitz, <i>Subdirect sum decompositions of endomorphism rings</i>	129
Gary Douglas Jones, <i>An asymptotic property of solutions of $y''' + py' + qy = 0$</i>	135
Howard E. Lacey, <i>On the classification of Lindenstrauss spaces</i>	139
Charles Dwight Lahr, <i>Approximate identities for convolution measure algebras</i>	147
George William Luna, <i>Subdifferentials of convex functions on Banach spaces</i>	161
Nelson Groh Markley, <i>Locally circular minimal sets</i>	177
Robert Wilmer Miller, <i>Endomorphism rings of finitely generated projective modules</i>	199
Donald Steven Passman, <i>On the semisimplicity of group rings of linear groups</i>	221
Bennie Jake Pearson, <i>Dendritic compactifications of certain dendritic spaces</i>	229
Ryōtarō Satō, <i>Abel-ergodic theorems for subsequences</i>	233
Henry S. Sharp, Jr., <i>Locally complete graphs</i>	243
Harris Samuel Shultz, <i>A very weak topology for the Mikusinski field of operators</i>	251
Elena Stroescu, <i>Isometric dilations of contractions on Banach spaces</i>	257
Charles W. Trigg, <i>Versum sequences in the binary system</i>	263
William L. Voxman, <i>On the countable union of cellular decompositions of n-manifolds</i>	277
Robert Francis Wheeler, <i>The strict topology, separable measures, and paracompactness</i>	287