

Pacific Journal of Mathematics

ON THE CLASSIFICATION OF LINDENSTRAUSS SPACES

HOWARD E. LACEY

ON THE CLASSIFICATION OF LINDENSTRAUSS SPACES

H. ELTON LACEY

A Lindenstrauss space is a real Banach space X such that its dual X^* is linearly isometric to $L_1(\mu)$ for some measure μ . The purpose of this paper is to describe how certain classical types of Lindenstrauss spaces are characterized by mappings from a compact Hausdorff space S into $C(S)^*$.

Let S be a compact Hausdorff space and $\rho: S \rightarrow C(S)^*$ a bounded function such that for each $f \in C(S)$, the function f_ρ defined by $f_\rho(s) = \int f d\rho(s)$ is integrable with respect to each regular Borel measure on S . Thus ρ induces a natural bounded linear operator P on $C(S)^*$ defined by $(P\mu)(f) = \int f_\rho d\mu$ for all $\mu \in C(S)^*$ and $f \in C(S)$. If (i) $\|\rho(s)\| \leq 1$ for all $s \in S$, and (ii) whenever $\mu \in C(S)$ and $\int f d\mu = 0$ for all $f \in C(S)$ with $f = f_\rho$, then $\int f_\rho d\mu = 0$ for all $f \in C(S)$, then ρ is said to be an *affine mapping*.

It was shown in [1] that if ρ is affine, then $X_\rho = \{f \in C(S): f = f_\rho\}$ is a Lindenstrauss space, P is a contractive projection on $C(S)^*$ with kernel equal to $X_\rho^\perp = \{\mu \in C(S)^*: \mu(f) = 0 \text{ for all } f \in X_\rho\}$. Moreover, the restriction mapping $\mu \rightarrow \mu|_{X_\rho}$ is a linear isometry from the range of P onto X_ρ^* .

Condition (ii) of the definition of an affine mapping is usually the hardest to verify. In [7] Gleit gives a nontrivial example when $\rho(s) \geq 0$ for all $s \in S$. Although he did not actually use this terminology, careful inspection of his proof yields that the mapping he postulates is indeed an affine mapping. In fact, slight modifications in his proof yields the following general result.

THEOREM 1. (Gleit). *Let S be a compact Hausdorff space and T a closed subset of S . Let $\rho: S \rightarrow C(S)^*$ be a mapping such that*

- (a) $\|\rho(s)\| \leq 1$ for all $s \in S$,
- (b) $\rho(s) = \varepsilon_s$ (i.e., point evaluation) for $s \in S \setminus T$,
- (c) for $T_2 = \{s \in T: \rho(s) = \varepsilon_s\}$ and $T_1 = T \setminus T_2$, we suppose that $T_1 \neq S$ and $|\rho(s)|(T_1) = 0$ for all $s \in T_1$,
- (d) $\rho|_T$ is weak* continuous.

Then ρ is an affine mapping and X_ρ^ is, in fact, linearly isometric to $\{\mu \in C(S)^*: |\mu|(T_1) = 0\}$.*

As mentioned above, Gleit assumes in addition that $\rho(s) \geq 0$ for

all $s \in S$. This assumption yields that X_ρ is ordered and X_ρ^* is an L space in its dual ordering. This will be proved in general in Theorem 2. For this and other results certain classical types of Lindenstrauss spaces are recalled.

$C(S)$: the Banach space of all continuous real valued functions on a compact Hausdorff space S .

$C_0(LS)$: the Banach space of all continuous real valued functions which vanish at infinity on a locally compact Hausdorff space LS .

$A(K)$: the Banach space of all affine continuous functions on a compact Choquet simplex K .

$A_0(K)$: the Banach space of all affine continuous functions on a compact Choquet simplex space K which vanish at a fixed extreme point of K .

$C_\sigma(S)$: the set of all $f \in C(S)$ such that $f(s) = -f(\sigma s)$ for all $s \in S$, where σ is a homeomorphism on S such that σ^2 is the identity.

M space: the set of all $f \in C(S)$ satisfying a fixed set of relations $\{s_i, s'_i, \lambda_i\}$ where $s_i, s'_i \in S, \lambda_i \in [0, 1]$ and f satisfies $f(s_i) = \lambda_i f(s'_i)$ for all i in some index set.

G space: the set of all $f \in C(S)$ satisfying a fixed set of relations $\{s_i, s'_i, \lambda_i\}$ where s_i, s'_i are in $S, \lambda_i \in [-1, 1]$ and f satisfies $f(s_i) = \lambda_i f(s'_i)$ for all i in some index set.

THEOREM 2. *Let S be a compact Hausdorff space and $\rho: S \rightarrow C(S)^*$ be an affine mapping. If $\rho(s) \geq 0$ for all $s \in S$, then X_ρ is an $A_0(K)$ space. If $\rho(s) \geq 0$ and $\|\rho(s)\| = 1$ for all $s \in S$, then X_ρ is an $A(K)$ space. Conversely, each $A_0(K)$ and $A(K)$ space can be so represented.*

Proof. Let P be the projection determined by ρ . Let r be the restriction mapping from the range N of P onto X^* . Then r is a linear isometry and is clearly positive from N to X_ρ^* where X_ρ^* has the dual ordering from the induced ordering on X_ρ from $C(S)$.

Now N is a sublattice of $C(S)^*$. For, if μ, ν are in N , then since P is positive, $P(\mu \vee \nu) \geq \mu \vee \nu$ and since P is contractive, $\|\mu \vee \nu\| \geq \|P(\mu \vee \nu)\|$. Thus it follows that $\mu \vee \nu = P(\mu \vee \nu)$. Hence it remains only to show that r is order preserving to establish that X_ρ is an $A_0(K)$ space (see [2]). Let $x^* \in X_\rho^*$ be positive and have norm one and let $W = \{y^* \in X^*: 0 \leq y^*, \|y^*\| \leq 1\}$. Clearly W is a weak* closed convex set in X_ρ^* . Let S_0 be the weak* closure of the extreme points of W . Since each nonzero extreme point of W is the image (under restriction) of an extreme point of the unit sphere of $C(S)^*$ and since the positive cone of X_ρ is nonzero, it follows that S_0 is contained in the image of $\{\varepsilon_s: s \in S\} \cup \{0\} = S_1$

under restriction. Now, there is a probability measure μ on S_0 such that $\int y^*(f)d\mu(y^*) = x^*(f)$ for all $f \in X_\rho$ (see [19]) and by [8] there is a probability measure ν on S_1 such that $\int \hat{f}d\nu = \int y^*(f)d\mu(y^*) = x^*(f)$ for all $f \in X_\rho$, where \hat{f} is defined by $\hat{f}(\varepsilon_s) = f(s)$ and $\hat{f}(0) = 0$. Thus ν' defined by $\nu'(A) = \nu(\{\varepsilon_s: s \in A\})$ is a nonnegative measure on S and $\int f d\nu' = \int \hat{f} d\nu$ for all $f \in X_\rho$ (since $\hat{f}(0) = 0$). If $rP\nu' = x^*$, then since $P\nu' \geq 0$, the proof is complete. But, $\int f d\nu' = \int \hat{f} d\nu = \int y^*(f)d\mu(y^*) = x^*(f)$ for $f \in X_\rho$.

If, in addition, $\|\rho(s)\| = 1$ for all $s \in S$, then $1 \in X_\rho$ and it follows that X_ρ has a strong order unit and so X_ρ is an $A(K)$ space where K is a compact Choquet simplex.

Conversely, if we are given a simplex space $A_0(K)$, then without loss of generality $K = \{x^* \in A_0(K)^*: 0 \leq x^*, \|x^*\| \leq 1\}$. For each $x^* \in K$, let μ_{x^*} denote the unique maximal probability measure representing x^* (see [21]). It is well known that if S is equal to the weak* closure of the extreme points of K and $\rho(x^*)$ is the unique maximal probability measure on S representing x^* , ρ is an affine mapping and $A(K) = X_\rho$. Let P be the projection associated with ρ . Then it is also well known that $\text{kernel } P = A(K)^\perp$ (the reader can see [1] for a proof of this).

Now, let $\tilde{\rho}(x^*) = \rho(x^*)$ for $x^* \neq 0$ and $\tilde{\rho}(0) = 0$. Then $\tilde{\rho}$ is an affine mapping on S and the associated projection \tilde{P} is given by $\tilde{P}\mu = \mu - \mu(0) \cdot \varepsilon_0$. From this it follows that the kernel of \tilde{P} is equal to $\{\mu: (\mu - \mu(0) \cdot \varepsilon_0) \in A(K)^\perp\} = \{\mu: \mu \in A_0(K)^\perp\}$. Thus it follows that $X_{\tilde{\rho}} = A_0(K)$.

Let X be a Banach space, V the unit sphere of X^* , S the weak* closure of the set of extreme points of V , and $\sigma: V \rightarrow V$ be defined by $\sigma(x^*) = -x^*$ for all $x^* \in V$. The following theorem was also proved in [1]. The notation is as above.

THEOREM 3. *Let X be a Lindenstrauss space. Then $\rho: S \rightarrow C(S)^*$ defined by $\rho(x^*) = (1/2)[\mu - \mu \circ \sigma]$, where μ is a maximal probability measure on V (supported on S) representing x^* , is an affine mapping and $X = X_\rho$.*

Theorem 3 will be used to characterize some of the types of Lindenstrauss spaces in terms of affine mappings.

THEOREM 4. *Let S be a compact Hausdorff space and $\rho: S \rightarrow C(S)^*$ be a weak* continuous affine mapping. Then X_ρ is a $C_0(S_0)$ space for some compact Hausdorff space S_0 . Conversely, any $C_0(S_0)$ space can be so represented.*

Proof. Let $\mu \in C(S)^*$. Then $\int fd\mu = \int fdP\mu = \int f_\rho d\mu$ for all $f \in X_\rho$.

Thus $\int f_\rho d\mu = \int f_\rho dP\mu = \int f_{\rho\rho} d\mu$ for all $f \in C(S)$. Since ρ is weak* continuous, f_ρ and $f_{\rho\rho}$ are in $C(S)$. Thus $f_\rho = f_{\rho\rho}$ for all $f \in C(S)$ and $X_\rho = \{f_\rho: f \in C(S)\}$. Moreover, the operator Q defined by $Q(f) = f_\rho$ for $f \in C(S)$ is a constructive projection of $C(S)$ onto X_ρ . Consequently X_ρ is a $C_\sigma(S_0)$ space for some compact Hausdorff space S_0 by [1].

On the other hand, if $X = C_\sigma(S_0)$ for some compact Hausdorff space S_0 , then the weak* closure, S , of the set E of extreme points of the unit sphere of X^* is either E or $E \cup \{0\}$ depending on whether or not σ is fixed point free (see [16]). In particular for $s \in S_0$ and $\sigma(s) \neq s$, $(\varepsilon_s | X) \in E$. Thus by Theorem 3 the mapping defined by $\rho(s) = (1/2)(\varepsilon_s - \varepsilon_{\sigma(s)})$ defines an affine mapping and $X = X_\rho$. Clearly ρ is weak* continuous.

From the definition of an M space, if $\rho: S \rightarrow C(S)^*$ is any function which takes its values in $[0, 1]S = \{\lambda\varepsilon_s: 0 \leq \lambda \leq 1, s \in S\}$, then $X_\rho = \{f \in C(S): f(s) = \int fd\rho(s) \text{ for all } s \in S\}$ is an M space.

THEOREM 5. *Let X be an M space. Then there is a compact Hausdorff space S and an affine mapping $\rho: S \rightarrow C(S)^*$ such that ρ takes its values in $[0, 1]S$ and $X = X_\rho$.*

Proof. Let S be the weak* closure of the set E of positive extreme points of the unit sphere of X^* . It is well known that each element $x^* \in S$ is of the form λy^* with $y^* \in E$ and $0 \leq \lambda \leq 1$ [9]. Moreover this representation is unique if $x^* \neq 0$. Thus $\rho(x^*) = \lambda\varepsilon_{y^*}$ defines an affine mapping and $X_\rho = X$ (see the proof of Theorem 2).

Similarly, if $\rho: S \rightarrow C(S)^*$ takes its values in $[-1, 1]S = \{\lambda\varepsilon_s: -1 \leq \lambda \leq 1, s \in S\}$, then $X_\rho = \{f \in C(S): f(s) = \int fd\rho(s) \text{ for all } s \in S\}$ in a G space. On the other hand, if X is a G space and S is the weak* closure of the set E of extreme points of the unit sphere of X^* , then Fakhoury has shown in [3] that $S \subset [-1, 1]E$. Thus for any $x^* \in S$, $x^* = \alpha y^* + (1 - \alpha)(-y^*)$ for some $0 \leq \alpha \leq 1$ and $y^* \in E$. Moreover, $\mu = \alpha\varepsilon_{y^*} + (1 - \alpha)\varepsilon_{-y^*}$ is a maximal measure representing x^* and $(1/2)[\mu - \mu \circ \sigma] = (1/2)[(2\alpha - 1)\varepsilon_{y^*} + (1 - 2\alpha)\varepsilon_{-y^*}] = \rho(x^*)$ defines an affine mapping with $X = X_\rho$ by Theorem 3.

If S is a compact Hausdorff space, then ρ defined by $\rho(s) = \varepsilon_s$ is clearly an affine mapping and $C(S) = X_\rho$. On the other hand, if LS is a locally compact noncompact Hausdorff space, then the weak* closure S_0 of the set E of positive extreme points of the unit sphere of $C_0(LS)^+$ is $E \cup \{0\}$. Clearly S_0 can be thought of as $S \cup \{0\}$ since

E is homeomorphic to S . Let $\rho(s) = \varepsilon_s$ for $s \in S$ and $\rho(0) = 0$. Then as in Theorem 5, ρ is an affine mapping and $X_\rho = C_0(LS)$.

THEOREM 6. *Let S be a compact Hausdorff space and $\rho: S \rightarrow C(S)^*$ be a positive weak* continuous affine mapping. Then X_ρ is a $C_0(LS)$ space.*

Proof. By Theorems 2 and 4, X_ρ is simultaneously a $C_0(S_0)$ space and an $A_0(K)$ space. Thus by [16] X_ρ is a $C_0(LS)$ space for some locally compact Hausdorff space LS .

Open questions. The results in this paper are in the isometric theory of Banach spaces. It is natural to ask about the isomorphic theory. The isomorphic analogue to a Lindenstrauss space is a \mathcal{L}_∞ space. A Banach space X is said to be a $\mathcal{L}_{\infty, \lambda}$ space if for each finite dimensional subspace Y of X there is finite dimensional subspace Z of X with $Y \subset Z$ and $d(Z, l_\infty(\dim Z)) = \inf \{ \|T\| \|T^{-1}\| : T: Z \rightarrow l_\infty(\dim Z) \text{ is 1-1} \} \leq \lambda$.

Using proofs similar to those in [1] one can show that if F is a compact Hausdorff space and $\rho: S \rightarrow C(S)^*$ has the properties of an affine mapping except that we only assume that $\sup \|\rho(s)\| < \infty$ instead of $\|\rho(s)\| \leq 1$ for all $s \in S$, then X_ρ is an \mathcal{L}_∞ space.

Question 1. Is it possible to construct each \mathcal{L}_∞ space from such a mapping?

Another question which is well known is the following.

Question 2. Is every \mathcal{L}_∞ space isomorphic to a Lindenstrauss space? Recently, Benyamini and Lindenstrauss [21] have shown that there are separable Lindenstrauss spaces X and Y such that X^* is separable and Y^* is nonseparable and both X and Y are not isomorphic to a complemented subspace of a $C(S)$ space.

Some questions which arise in the isometric theory are discussed below. Let S be a compact Hausdorff space and $X \subset C(S)$ a closed linear subspace containing 1. Suppose further that $\{s \in S: \varepsilon_s \in X\}$ is an extreme point of the unit sphere of X^* is dense in X . Then $X = A(K)$ where $K = \{x^* \in X^*: x^*(1) = 1 = \|x^*\|\}$ has the weak* topology and the extreme points of K are homeomorphic to $\partial_X S$ and their closure is homeomorphic to S . In particular, each maximal measure on K is supported on S .

Question 3. Is there an affine mapping ρ of S into the maximal measures on K such that $\rho(s) = \varepsilon_s$ if and only if $s \in \partial_X S$?

In [20] Rao has shown that there is a Borel measurable mapping when S is metrizable metrizable.

This is related to the following question.

Question 4. If S is a metrizable compact Hausdorff space and $T \subset S$ is a dense G_δ set, is there a compact Choquet simplex with extreme points homeomorphic to T and their closure homeomorphic to S ?

Let $X \subset C(S)$ be as above. It is shown in [6] that if X is a Lindenstrauss space, then X is maximal with respect to $\partial_X S$. That is, if $X \subset Y$ and $\partial_Y S = \partial_X S$, then $X = Y$. An easy application of Zorn's lemma shows that for any such $X \subset C(S)$ there is a maximal $Y \supset X$ with respect to $\partial_X S$.

Question 5. Is any X maximal with respect to $\partial_X S$ a Lindenstrauss space?

REFERENCES

1. J. B. Bednar and H. E. Lacey, *Concerning Banach spaces whose duals are L spaces*, Pacific J. Math., **41** (1972), 13-24.
2. E. B. Davies, *On the Banach space duals of certain spaces with the Riesz decomposition property*, Quart. J. Math., Oxford **18** (1967), 109-111.
3. H. Fakhoury, *Preduax de L espaces; Proprieties du G espaces et des C espaces*, C. R. Acad. Sci. Paris t **271** (1970), Series A, 941-944.
4. ———, *Sélections Linéaires Associees au Théorème de Hahn-Banach*, (preprint).
5. ———, *Une Characterisation des L espaces duax*, (to appear).
6. D. A. Edwards and G. Vincent-Smith, *A Weierstrass Stone theorem for Choquet simplexes*, Ann. Inst. Fourier Grenoble **18** (1968), 261-282.
7. A. Gleit, *On the existence of simplex spaces*, Israel J. Math., **9** (1971), 199-209.
8. J. Hardy and H. E. Lacey, *Extensions of regular Borel measures*, Pacific J. Math., **24** (1968), 277-282.
9. S. Kakutani, *Concrete representations of abstract M spaces*, Ann. Math., **42** (1941), 994-1024.
10. H. E. Lacey, *A note concerning $A^* = L_1(\mu)$* , Proc. Amer. Math. Soc., **29** (1971), 525-528.
11. H. E. Lacey, and P. D. Morris, *Continuous linear operators on spaces of continuous functions*, Proc. Amer. Math. Soc., **17** (1966), 848-853.
12. ———, *On spaces of type $A(K)$ and their duals*, Proc. Amer. Math. Soc., **23** (1969), 151-157.
13. A. J. Lazar, *The unit ball in conjugate L_1 spaces*, Duke Math. J., **39** (1972), 1-8.
14. J. Lindenstrauss and A. Pelczynski, *Absolutely summing operators in L_p spaces and their applications*, Studia Math., **29** (1968), 275-326.
15. J. Lindenstrauss and H. P. Rosenthal, *The \mathcal{L}_p spaces*, Israel J. Math., **7** (1969), 325-349.
16. J. Lindenstrauss and D. E. Wulbert, *On the classification of the Banach spaces whose duals are L_1 spaces*, J. Functional Anal., (1969), 332-349.
17. A. Pelczynski, *On Banach spaces containing $L_1(\mu)$* , Studia Math., **30** (1968), 231-246.

18. A. Pelczynski and Z. Semaderi, *Spaces of continuous functions III*, *Studia Math.*, **18** (1959), 211-222.
19. R. R. Phelps, *Lectures on Choquet's Theorem*, *Math. Studies*, D. Van Nostrand, Princeton, 1966.
20. M. M. Rao, *Borel Selections of Representing Measures*, Aarhus University notes, 1970.
21. Y. Benyamini and J. Lindenstrauss, *A predual of l_1 which is not isomorphic to a $C(K)$ spaces*, (preprint).

Received December 14, 1971 and in revised form January 18, 1973.

UNIVERSITY OF TEXAS AT AUSTIN

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

* C. DePrima will replace J. Dugundji until August 1974.

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

K. Adachi, Masuo Suzuki and M. Yoshida, <i>Continuation of holomorphic mappings, with values in a complex Lie group</i>	1
Michael Aschbacher, <i>A characterization of the unitary and symplectic groups over finite fields of characteristic at least 5</i>	5
Larry Eugene Bobisud and James Calvert, <i>Energy bounds and virial theorems for abstract wave equations</i>	27
Christer Borell, <i>A note on an inequality for rearrangements</i>	39
Peter Southcott Bullen and S. N. Mukhopadhyay, <i>Peano derivatives and general integrals</i>	43
Wendell Dan Curtis, Yu-Lee Lee and Forrest Miller, <i>A class of infinite dimensional subgroups of $\text{Diff}^r(X)$ which are Banach Lie groups</i>	59
Paul C. Eklof, <i>The structure of ultraproducts of abelian groups</i>	67
William Alan Feldman, <i>Axioms of countability and the algebra $C(X)$</i>	81
Jack Tilden Goodykoontz, Jr., <i>Aposyndetic properties of hyperspaces</i>	91
George Grätzer and J. Płonka, <i>On the number of polynomials of an idempotent algebra. II</i>	99
Alan Trinler Huckleberry, <i>The weak envelope of holomorphy for algebras of holomorphic functions</i>	115
John Joseph Hutchinson and Julius Martin Zelmanowitz, <i>Subdirect sum decompositions of endomorphism rings</i>	129
Gary Douglas Jones, <i>An asymptotic property of solutions of $y''' + py' + qy = 0$</i>	135
Howard E. Lacey, <i>On the classification of Lindenstrauss spaces</i>	139
Charles Dwight Lahr, <i>Approximate identities for convolution measure algebras</i>	147
George William Luna, <i>Subdifferentials of convex functions on Banach spaces</i>	161
Nelson Groh Markley, <i>Locally circular minimal sets</i>	177
Robert Wilmer Miller, <i>Endomorphism rings of finitely generated projective modules</i>	199
Donald Steven Passman, <i>On the semisimplicity of group rings of linear groups</i>	221
Bennie Jake Pearson, <i>Dendritic compactifications of certain dendritic spaces</i>	229
Ryōtarō Satō, <i>Abel-ergodic theorems for subsequences</i>	233
Henry S. Sharp, Jr., <i>Locally complete graphs</i>	243
Harris Samuel Shultz, <i>A very weak topology for the Mikusinski field of operators</i>	251
Elena Stroescu, <i>Isometric dilations of contractions on Banach spaces</i>	257
Charles W. Trigg, <i>Versum sequences in the binary system</i>	263
William L. Voxman, <i>On the countable union of cellular decompositions of n-manifolds</i>	277
Robert Francis Wheeler, <i>The strict topology, separable measures, and paracompactness</i>	287