

Pacific Journal of Mathematics

ANGULAR LIMITS OF LOCALLY FINITELY VALENT HOLOMORPHIC FUNCTIONS

DAVID CHARLES HADDAD

ANGULAR LIMITS OF LOCALLY FINITELY VALENT HOLOMORPHIC FUNCTIONS

DAVID C. HADDAD

A function f defined in a domain D is n -valent in D if $f(z) - w_0$ has at most n zeros in D for each complex number w_0 . The purpose of this paper is to show that a sufficient condition for a holomorphic function f in $|z| < 1$ to have angular limits almost everywhere on $|z| = 1$ is that there exist a positive integer n and a positive number r_0 such that f is n -valent in each component of the set $\{z: |f(z)| > r_0\}$.

We have previously shown that the same conditions on f imply that f is a quasi-normal function of order at most $n - 1$ [3, Theorem 2], and f has angular limits at a dense subset of $|z| = 1$ [3, Corollary 1]. Note that the bound n on the valence of f is the same for each component of $\{z: |f(z)| > r_0\}$. This uniformity on n is essential to the conclusion that f has angular limits almost everywhere on $|z| = 1$; for we have shown in the example in [2] that if the uniformity is dropped, then f need not even have asymptotic values at a dense subset of $|z| = 1$.

If $w = f(z)$ is a nonconstant, holomorphic function in $|z| < 1$, we denote by F the Riemann surface of f^{-1} (as a covering surface over the w -plane). If S is a subset of $|z| = 1$, then $m(S)$ denotes the Lebesgue measure of S .

A Jordan arc $T = \{z = h(t): 0 < t < 1\}$ lying in a domain D is a *crosscut* of D if $h(t) \rightarrow z_0 \in \partial D$ as $t \downarrow 0$, $h(t) \rightarrow z_1 \in \partial D$ as $t \uparrow 1$, and $z_0 \neq z_1$. If $z_0 = z_1$, then T is a *loopcut* of D .

If a holomorphic function f in $|z| < 1$ is n -valent in a component $D(r)$ of the set $\{z: |f(z)| > r\}$ then the connectivity of $D(r)$ is at most $n + 1$ [3, Lemma 3]. We denote by $D^*(r)$ the simply connected domain obtained by adding to $D(r)$ those (at most n) components of $\{z: |f(z)| \leq r\}$ that punch holes in $D(r)$.

LEMMA 1. *Let f be a nonconstant, holomorphic function in $|z| < 1$ that is n -valent in each component of the set $\{z: |f(z)| > r_0\}$. For each $r > r_0$, let $\{D_k(r)\}$ denote the at most countable collection of components of $\{z: |f(z)| > r\}$. Then there exists a countable subset E of (r_0, ∞) such that $\partial D_k^*(r)$ is a Jordan curve for all k and all $r \in (r_0, \infty) - E$.*

Proof. Define a set $R = \{r: r > r_0, \text{ and } F \text{ has no branch points lying over the circle } |w| = r\}$. Then the set $(r_0, \infty) - R$ is at most countable. If $r \in R$, then for each k , $\partial D_k^*(r) \cap \{|z| < 1\}$ consists of at

most countably many crosscuts and loopcuts T_j^k of $|z| < 1$ by [2, Corollary 1].

We show that if for a fixed k there are infinitely many curves T_j^k , then their diameters tend to zero as $j \rightarrow \infty$. If the diameters did not tend to zero, then the sequence $\{T_j^k\}$ would have an accumulation continuum in $|z| \leq 1$. Since f is a nonconstant, holomorphic function, $\{T_j^k\}$ cannot have an accumulation continuum in $|z| < 1$. By [2, Theorem 3], f has asymptotic values at a dense subset of $|z| = 1$, and hence, by a theorem of MacLane [4, Theorem 1], the sequence $\{T_j^k\}$ of level curves cannot have an arc of $|z| = 1$ for an accumulation continuum. Hence, the diameters of the curves T_j^k tend to zero as $j \rightarrow \infty$.

We still must show that there exists a countable subset E of (r_0, ∞) such that $\partial D_k^*(r)$ has no double points for all k and all $r \in (r_0, \infty) - E$. Suppose to the contrary that S is an uncountable subset of R and that for each $r \in S$ there exists a component $D(r)$ of the set $\{z: |f(z)| > r_0\}$ such that $\partial D^*(r)$ has double points. This implies that for each $r \in S$, $\partial D^*(r)$ contains a loopcut T_r , since the curves comprising $D^*(r) \cap \{|z| < 1\}$ are Jordan arcs for all $r \in R$. The domain $D^*(r)$ cannot be interior to a loopcut; for if it were, f would be unbounded in $D^*(r)$ by the extended maximum principle, and, consequently, the loopcut would determine two distinct asymptotic tracts ending at one point contradicting [2, Theorem 2]. (See [4] or [2] for the definition of an asymptotic tract.) Let G_r denote the domain interior to the loopcut T_r . The uncountable collection of open sets G_r must contain a pair that intersect, say G_q and G_s where $q < s$. Since the loopcuts T_q and T_s cannot intersect inside $|z| < 1$, then $G_q \subset G_s$, and T_q and T_s end at the same point of $|z| = 1$. By [2, Corollary 1], T_q and T_s determine at least two (since $q \neq s$) asymptotic tracts ending at one point contradicting [2, Theorem 2]. Thus, there must exist a countable subset E of (r_0, ∞) such that $\partial D_k^*(r)$ is a Jordan curve for all k and all $r \in (r_0, \infty) - E$.

LEMMA 2. *Let f be a nonconstant, holomorphic function in $|z| < 1$ that is n -valent in each component of $\{z: |f(z)| > r_0\}$. If $r_1 > r_0$ and $D(r_1)$ is a component of $\{z: |f(z)| > r_1\}$, then f has angular limits almost everywhere on $E(r_1) = \overline{D(r_1)} \cap \{|z| = 1\}$.*

Proof. We assume $m(E(r_1)) > 0$, for, otherwise, there is nothing to prove. For $r < r_1$, we denote by $D(r)$ the component of $\{z: |f(z)| > r\}$ containing $D(r_1)$, and we write $E(r) = \overline{D(r)} \cap \{|z| = 1\}$. We first show that there exists $s \in (r_0, r_1)$ such that $\partial D^*(s)$ is a rectifiable Jordan curve.

By Lemma 1, the set $R = \{r \in (r_0, r_1) \text{ such that } F \text{ has no branch}$

points over $|w| = r$ and $\partial D^*(r)$ is a Jordan curve} is the whole interval (r_0, r_1) minus possibly a set of measure zero. Let $C(r) = \partial D(r) \cap \{|z| < 1\}$, and let Γ be the family $\{C(r): r \in R\}$. By passing to the Riemann surface F , it is not hard to show that the extremal length of the family Γ is bounded by $2n\pi \log r_1/r_0$, and this implies $\partial D(r)$ is rectifiable for infinitely many values $r \in R$ (for example, see [2, Theorem 1]). Thus, we can choose $s \in (r_0, r_1)$ such that $\partial D^*(s)$ is a rectifiable Jordan curve.

By the Riemann mapping theorem and Carathéodory's theorem on boundary correspondence there exists a homeomorphism g of $\overline{D^*(s)}$ onto $|\zeta| \leq 1$ that is a conformal mapping of $D^*(s)$ onto $|\zeta| < 1$. Since the connectivity of $D(s)$ is finite, $|f(g^{-1}(\zeta))| > s$ in some annulus $t < |\zeta| < 1$. Hence, $f \circ g^{-1}$ has angular limits almost everywhere on $|\zeta| = 1$ by a simple extension of theorems of Fatou [1, p. 19] and F. and M. Riesz [1, p. 22] on angular limits. Since $\partial D^*(s)$ is a rectifiable Jordan curve, g^{-1} maps a set of measure zero on $|\zeta| = 1$ onto a set of measure zero on $\partial D^*(s)$ by a theorem of F. and M. Riesz [1, p. 50]. Thus f has asymptotic values almost everywhere on $E(s)$ and hence angular limits almost everywhere on $E(s)$ by [3, Theorem 3]. This completes the proof of the lemma since $E(s) \supset E(r_1)$.

LEMMA 3. *Let $\{I_j\}$ be a sequence of mutually disjoint open arcs on $|z| = 1$, and let $C = \bigcup_j I_j$. Let f be a continuous function on $\{|z| < 1\} \cup C$ that is holomorphic in $|z| < 1$. Let $|f(z)| = r_0$ for $z \in C$, $|f(0)| > r_0$, and the set $D = \{z: |z| < 1, |f(z)| > r_0\}$ be a connected set whose boundary contains the circle $|z| = 1$. If f is n -valent in D , then $|f(0)| \leq r_0 \exp [2\pi^4 n/m(C)^2]$.*

Proof. Let $\gamma(r)$ be the level set $\{z: |f(z)| = r\}$. The proof consists of finding bounds on the extremal length $\lambda(\Gamma)$ of the family $\Gamma = \{\gamma(r): r_0 < r < |f(0)|\}$, and F has no branch points lying over $|w| = r$. By passing to the Riemann surface F , it can be shown that $\lambda(\Gamma) \leq 2\pi n/\log |f(0)|/r_0$ (for example, see [2, Theorem 1]).

By our hypotheses on f , each arc I_j must be separated from the point $z = 0$ by a level curve of $\{z: |f(z)| = r\}$ for each r in the interval $(r_0, |f(0)|)$. None of these curves can be relatively compact curves encircling the point $z = 0$ by the maximum principle. Thus, the Euclidean length of a level curve separating I_j from $z = 0$ is bounded below by $\min(2, (2/\pi)m(I_j))$. Hence, the Euclidean length of each $\gamma(r) \in \Gamma$ is bounded below by $(1/\pi)m(C)$. By considering the linear density $\rho(z)$ defined to be 1 on D and 0 elsewhere, we can easily obtain the inequality $\lambda(\Gamma) \geq (1/\pi^3)m(C)^2$. Combining the two bounds on $\lambda(\Gamma)$ we have $|f(0)| < r_0 \exp [2\pi^4 n/m(C)^2]$, which completes the proof of the lemma.

A point $e^{i\theta}$ is a *Plessner point* for a function f defined in $|z| < 1$ if for every Stolz angle S at $e^{i\theta}$, the cluster set of f at $e^{i\theta}$ with respect to the domain S is total.

THEOREM. *A sufficient condition for a holomorphic function f in $|z| < 1$ to have finite angular limits almost everywhere on $|z| = 1$ is that there exist a positive number r_0 and a positive integer n such that f is n -valent in each component of the set $\{z: |f(z)| > r_0\}$.*

Proof. Suppose to the contrary that the set of points of $|z| = 1$ at which f does not have finite angular limits has positive measure. Then, by a theorem of Plessner [1, p. 147] and a theorem of Priwalow [1, p. 146], f must be a nonconstant function whose set of Plessner points P has positive measure.

For each $r > 0$, let $\{D_j(r)\}$ denote the at most countable collection of components of the set $\{z: |f(z)| > r\}$. By Lemma 1, there exists $r_1 > r_0$ such that $\partial D_j^*(r_1)$ is a Jordan curve for each j and F has no branch points over the circle $|w| = r_1$. Thus, $\partial D_j^*(r_1) \cap \{|z| < 1\}$ consists of at most countably many level curves which are crosscuts of $|z| < 1$. Write $D_j = D_j(r_1)$, $E_j = \overline{D_j^*} \cap \{|z| = 1\}$, and $E'_j = \{|z| = 1\} - E_j$. Since by Lemma 2, f has angular limits almost everywhere on $\bigcup_j E_j$, we can assume $P \subset \bigcap_j E'_j$. Let ω_j be the harmonic measure in D_j^* of the set $\partial D_j^* \cap \{|z| < 1\}$. We need the following lemma whose proof we postpone.

LEMMA 4. *There exists a harmonic function v in $|z| < 1$ having angular limit 0 almost everywhere on $\bigcap_j E'_j$, and $\omega_j(z) \geq 1 - v(z)$ for $z \in D_j^*$ ($j = 1, 2, \dots$).*

Thus, there exists a point $z_0 \in P$ at which v has angular limit 0. Then, by the definition of P , there exists a sequence $\{z_k\}$ of points lying inside a Stolz angle at z_0 and converging to z_0 such that $|f(z_k)| > r_1$ for each k and $f(z_k) \rightarrow \infty$ as $k \rightarrow \infty$. At most finitely many z_k can lie in the same component D_j since $z_0 \in P \subset \bigcap_j E'_j$. Hence, we can assume (by taking subsequences if necessary) that $z_j \in D_j$ ($j = 1, 2, \dots$) and $D_j \cap D_k = \phi$ for $j \neq k$. By Lemma 4, $\omega_j(z_j) \rightarrow 1$ as $j \rightarrow \infty$.

By the Riemann mapping theorem and Carathéodory's theorem on boundary correspondence, there exists a homeomorphism g_j of $\overline{D_j^*}$ onto $|\zeta| < 1$ that is a holomorphic map of D_j^* onto $|\zeta| < 1$ sending z_j into 0. Applying Lemma 3 to the function $h_j = f \circ g_j^{-1}$ and the set $C_j = g_j(\partial D_j^* \cap |z| < 1)$ we have $|h_j(0)| \leq r_1 \exp[2\pi^4 n/m(C_j)^2]$. On the one hand, $h_j(0) = f(z_j) \rightarrow \infty$ as $j \rightarrow \infty$. On the other hand, $m(C_j) = 2\pi\omega_j(g_j^{-1}(0)) = 2\pi\omega_j(z_j) \rightarrow 2\pi$ as $j \rightarrow \infty$, and this implies $h_j(0) \rightarrow \infty$ as $j \rightarrow \infty$. Thus to complete the proof of the theorem, we need only

prove Lemma 4.

Let u_j be the harmonic measure in $|z| < 1$ of the set E'_j , and let $v_k(z) = \sum_{j=1}^k (1 - u_j(z))$. Clearly, $\{v_k\}$ is an increasing sequence of nonnegative harmonic functions, and v_k has angular limit 0 at each point of the set $\bigcap_{j=1}^k E'_j$. Since the set $\overline{D_j^*} \cap \overline{D_q^*}$ can contain at most two points for $j \neq q$, each point $e^{i\theta}$ lies in at most one of the sets E_1, E_2, \dots, E_k for all but finitely many values of θ in the interval $[0, 2\pi)$. Hence, $\overline{\lim}_{z \rightarrow e^{i\theta}} v_k(z) \leq 1$ ($k = 1, 2, \dots$) for all but finitely many values $\theta \in [0, 2\pi)$. It follows from the extended maximum principle that $v_k(z) \leq 1$ for $|z| \leq 1$ ($k = 1, 2, \dots$). By Harnack's theorem, the sequence $\{v_k\}$ converges in $|z| < 1$ to a bounded harmonic function $v(z)$. Let $I = \{\theta: 0 \leq \theta < 2\pi, e^{i\theta} \in \bigcap_j E'_j\}$, and v, v_1, v_2, \dots have angular limits at $e^{i\theta}$. Then, writing $v(e^{i\theta})$ for the angular limit of v at $e^{i\theta}$, we have

$$\begin{aligned} \int_I v(e^{i\theta}) d\theta &= \int_I v(e^{i\theta}) - v_k(e^{i\theta}) d\theta \\ &\leq \int_0^{2\pi} v(e^{i\theta}) - v_k(e^{i\theta}) d\theta \\ &= v(0) - v_k(0) . \end{aligned}$$

Thus, v has angular limit 0 at $e^{i\theta}$ for almost all $\theta \in I$, since $v(0) - v_k(0) \rightarrow 0$ as $k \rightarrow \infty$. Since the set $\bigcap_j E'_j - \{e^{i\theta}: \theta \in I\}$ has measure zero by Fatou's theorem, v has angular limit 0 almost everywhere on $\bigcap_j E'_j$. Clearly, $v(z) \geq v_j(z) \geq 1 - u_j(z)$ for all j and $|z| < 1$, and by Carleman's principle of domain extension, $\omega_j(z) \geq u_j(z)$ for $z \in D_j^*$ ($j = 1, 2, \dots$). This completes the proof of Lemma 4 and hence of the theorem.

REMARK. The conclusion of the theorem raises the following question. Are all functions that satisfy the hypotheses of the theorem of bounded characteristic? This seems to be a difficult question to answer. The best we can presently show is that $T(r) = o(1/1 - r)$, where T is the Nevanlinna characteristic of f .

REFERENCES

1. E. F. Collingwood and A. J. Lohwater, *The Theory of Cluster Sets*, Cambridge University Press, 1966.
2. D. C. Haddad, *Asymptotic values of finitely valent functions*, Duke Math. J., **39** (1972), 391-367.
3. ———, *Boundary behavior and quasi-normality of finitely valent holomorphic functions*, to appear, Canad. J. Math., (1973).
4. G. R. MacLane, *Asymptotic Values of Holomorphic Functions*, Rice University Studies, **49** No. 1 (1963).

Received June 22, 1972. The author is indebted to the late Professor G. R. MacLane for his considerable assistance.

WEST VIRGINIA COLLEGE OF GRADUATE STUDIES

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Jan Aarts and David John Lutzer, <i>Pseudo-completeness and the product of Baire spaces</i>	1
Gordon Owen Berg, <i>Metric characterizations of Euclidean spaces</i>	11
Ajit Kaur Chilana, <i>The space of bounded sequences with the mixed topology</i>	29
Philip Throop Church and James Timourian, <i>Differentiable open maps of $(p + 1)$-manifold to p-manifold</i>	35
P. D. T. A. Elliott, <i>On additive functions whose limiting distributions possess a finite mean and variance</i>	47
M. Solveig Espelie, <i>Multiplicative and extreme positive operators</i>	57
Jacques A. Ferland, <i>Domains of negativity and application to generalized convexity on a real topological vector space</i>	67
Michael Benton Freeman and Reese Harvey, <i>A compact set that is locally holomorphically convex but not holomorphically convex</i>	77
Roe William Goodman, <i>Positive-definite distributions and intertwining operators</i>	83
Elliot Charles Gootman, <i>The type of some C^* and W^*-algebras associated with transformation groups</i>	93
David Charles Haddad, <i>Angular limits of locally finitely valent holomorphic functions</i>	107
William Buhmann Johnson, <i>On quasi-complements</i>	113
William M. Kantor, <i>On 2-transitive collineation groups of finite projective spaces</i>	119
Joachim Lambek and Gerhard O. Michler, <i>Completions and classical localizations of right Noetherian rings</i>	133
Kenneth Lamar Lange, <i>Borel sets of probability measures</i>	141
David Lowell Lovelady, <i>Product integrals for an ordinary differential equation in a Banach space</i>	163
Jorge Martinez, <i>A hom-functor for lattice-ordered groups</i>	169
W. K. Mason, <i>Weakly almost periodic homeomorphisms of the two sphere</i>	185
Anthony G. Mucci, <i>Limits for martingale-like sequences</i>	197
Eugene Michael Norris, <i>Relationally induced semigroups</i>	203
Arthur E. Olson, <i>A comparison of c-density and k-density</i>	209
Donald Steven Passman, <i>On the semisimplicity of group rings of linear groups. II</i>	215
Charles Radin, <i>Ergodicity in von Neumann algebras</i>	235
P. Rosenthal, <i>On the singularities of the function generated by the Bergman operator of the second kind</i>	241
Arthur Argyle Sagle and J. R. Schumi, <i>Multiplications on homogeneous spaces, nonassociative algebras and connections</i>	247
Leo Sario and Cecilia Wang, <i>Existence of Dirichlet finite biharmonic functions on the Poincaré 3-ball</i>	267
Ramachandran Subramanian, <i>On a generalization of martingales due to Blake</i>	275
Bui An Ton, <i>On strongly nonlinear elliptic variational inequalities</i>	279
Seth Warner, <i>A topological characterization of complete, discretely valued fields</i>	293
Chi Song Wong, <i>Common fixed points of two mappings</i>	299