

# Pacific Journal of Mathematics

**DECOMPOSITION OF PLANE CONVEX SETS. II. SETS  
ASSOCIATED WITH A WIDTH FUNCTION**

RUTH SILVERMAN

DECOMPOSITION OF PLANE CONVEX SETS,  
PART II:  
SETS ASSOCIATED WITH A WIDTH FUNCTION

RUTH SILVERMAN

**This paper treats several classes  $K$  of plane convex bodies such that the sum of any two members of the class is again a member of the class. In each case  $K$  is a class of bodies associated with a certain width function. An explicit characterization is provided for the corresponding subclass  $I(K)$  consisting of all indecomposable members of  $K$ .**

This result was proved by the author [4].

The results and methods of that paper are used to characterize  $I(K)$  when  $K$  consists of all plane convex bodies of constant width, also when  $K$  consists of all plane convex bodies whose associated width functions are a multiple of a given width function, known as bodies of constant relative width. It is supposed without further mention that all members of  $K$  have piecewise continuously second differentiable support functions.

The class  $K$  of all plane convex bodies has the property that the sum of any two members of the class is again a member of the class. The indecomposable subclass  $I(K)$  is the class of all triangles and line segments.

The reader is referred to [4] for the definitions used in this paper, which are not repeated, for the sake of brevity. For clarity it is, however, pointed out that as in [4], a *body* is a compact set.

This paper shows that, relative to the class  $K$  of all plane convex bodies of constant width, the  $K$ -indecomposable bodies are those whose boundaries are composed solely of circular arcs and have a corner point opposite every arc, or equivalently, are composed solely of circular arcs with radius equal to the width of the body. A more general problem is also considered; what are the  $K$ -indecomposable members of the family  $K$  of sets with width function a multiple of a given continuously second differentiable width function? The  $K$ -indecomposable members of  $K$  are these bodies which have a corner point in every direction, that is, at least one of the two support lines in every direction goes through a corner point.

Let  $K$  be a convex body; by the *width* of  $K$  in the direction  $\theta$ , denoted  $w(\theta)$ , we mean the distance between the two parallel supporting lines of  $K$  in the direction perpendicular to  $\theta$ .

Let  $w$  be a positive function, and  $K(w)$  be the associated family of sets whose width functions are a multiple of  $w$ . The machinery

we will set up requires that the sets to be decomposed, or proved indecomposable, have the origin in their interior.

The following remark is elementary but useful. Its proof is left to the reader.

**LEMMA 1.** *Let  $w$  be a positive function, and define the family of sets  $K(w)$  as above. A set  $B$  in  $K(w)$  is  $K$ -decomposable if and only if it has a  $K$ -decomposable translate whose interior contains the origin.*

Let  $w(\theta)$  be a width function with continuous second derivative, and  $K(w)$  the associated family of sets whose width functions are a multiple of  $w(\theta)$ . We will determine the  $K$ -indecomposable members of  $K(w)$ .

**LEMMA 2.** *An admissible pair  $\{\varphi_1, \varphi_2\}$  is the restriction of the support function of a set with width function  $w(\theta)$ , if*

$$\varphi_1(-t) + \varphi_2(t) = \sqrt{1+t^2} w(\text{arc cot } t),$$

for all real  $t$ .

*Proof.* Let  $K$  be a convex body in  $R^2$ , whose interior contains the origin, with width function  $w(\theta)$ . Then

$$\varphi_2(t) = \max_{(x,y) \in K} (tx + y),$$

and

$$\varphi_1(t) = \max_{(x,y) \in K} (tx - y),$$

so

$$\varphi_1(-t) = \max_{(x,y) \in K} (-tx - y) = -\min_{(x,y) \in K} (tx + y).$$

Clearly the extrema of  $(tx + y)$  will occur on the boundary of  $K$ . For a specified value of  $t$ , the extrema will occur when  $(dy/dx) = -t$ , or where the one sided derivatives have the property

$$\left(\frac{dy}{dx}\right)_- \leq -t \leq \left(\frac{dy}{dx}\right)_+.$$

When we require that  $(dy/dx) = -t$ , we are actually looking for support lines for  $K$  in the direction  $\theta$  such that  $\cot \theta = -(dy/dx) = t$ . There exist exactly two points on  $K$  with support lines in this direction. Let us denote by  $P_2 = (x_2, y_2)$  the one minimizing  $(tx + y)$ , and by  $P_1 = (x_1, y_1)$  the one maximizing  $(tx + y)$ . Then

$$\varphi_2(t) = x_2 t + y_2, \quad \varphi_1(-t) = -x_1 t - y_1,$$

where

$$t = \cot \theta, \quad \cot \varphi = \frac{x_2 - x_1}{y_2 - y_1},$$

and

$$w(\theta) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} \cos(\theta - \varphi).$$

Substituting all the above, adding, and simplifying,

$$\varphi_2(t) + \varphi_1(-t) = w(\arccot t) \cdot \sqrt{1 + t^2}.$$

On the other hand, if  $\{\varphi_1, \varphi_2\}$  is an admissible pair satisfying

$$\varphi_2(t) + \varphi_1(-t) = h(t)$$

for all real  $t$ , let  $K$  be a set containing the origin in its interior, whose support function corresponds to  $\{\varphi_1, \varphi_2\}$ .

For a fixed direction  $\theta$ , suppose  $P_1 = (x_1, y_1)$  and  $P_2 = (x_2, y_2)$  are the points where the parallel support lines in direction perpendicular to  $\theta$  intersect  $K$ . Then

$$w(\theta) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} \cos(\theta - \varphi), \text{ where } \cot \varphi = \frac{x_2 - x_1}{y_2 - y_1}.$$

But  $P_1$  and  $P_2$  are extrema for  $(tx + y)$  on the set  $K$ , when  $t = \cot \theta$ ; suppose the maximum occurs at  $P_1$  and the minimum at  $P_2$ . Then  $\varphi_2(t) = x_1 t + y_1$ , and  $\varphi_1(-t) = -x_2 t - y_2$ . Then

$$h(\cot \theta) = \frac{w(\theta)}{|\sin \theta|}.$$

Therefore,

$$w(\theta) = h(\cot \theta) \cdot |\sin \theta|.$$

**LEMMA 3.** *An admissible pair  $\{\varphi_1, \varphi_2\}$  satisfying  $\varphi_1(-t) + \varphi_2(t) = h(t) = \sqrt{1 + t^2} w(\arccot t)$ , for all real  $t$ , where  $w(\theta)$  has continuous second derivative, can be nontrivially decomposed into the sum of two admissible pairs  $\{\sigma_1, \sigma_2\}$ ,  $\{\psi_1, \psi_2\}$ , satisfying  $\sigma_1(-t) + \sigma_2(t) = 1/2 h(t)$ ,  $\psi_1(-t) + \psi_2(t) = 1/2 h(t)$ , if there is an interval  $I = (a, b)$  on the real line such that neither  $\varphi_2(t)$  nor  $\varphi_1(-t)$  is linear on  $I$ , or on any proper subinterval of  $I$ .*

*Proof.* Defining  $\sigma_i(t) = 1/2[\varphi_i(t) + y_i(t)]$ , and  $\psi_i(t) = 1/2[\varphi_i(t) - y_i(t)]$ , as in Theorem 2, [4], we must have  $y_1(-t) + y_2(t) = 0$  for all real  $t$ .

Let  $h(t) = \sqrt{1 + t^2} w(\arccot t)$ . Then

$$h''(t) = \frac{w(\arccot t) + w''(\arccot t)}{(1 + t^2)^{3/2}}$$

where  $w''(\arccot t)$  denotes

$$\frac{d^2 w}{d\theta^2} \Big|_{\theta = \text{arc cot } t} .$$

$w$  and  $w''$  are both defined on  $[0, \pi]$  and continuous, so are bounded on  $[0, \pi]$ . Therefore,  $h''(t)$  is bounded by some number  $M$ , for all real  $t$ . Note that  $h''(t)$  is also continuous.

Let  $f_1(t) = \varphi_1'(-t)$ ,  $f_2(t) = \varphi_2'(t)$ , and let  $p_1(t) = \varphi_1''(-t)$ ,  $p_2(t) = \varphi_1''(t)$ . Consider a closed subinterval  $I_1 = [a_1, b_1]$  of  $I$  on which  $p_i$  is continuous for  $i = 1, 2$ . Let  $T_i = \{x \in I_1 \mid p_i(t) = 0\}$ .  $T_i$  is compact for  $i = 1, 2$ , and  $T_i \neq I_1$  as  $\varphi_i$  is not linear on  $I_1$ . If  $I_1 = T_1 \cup T_2$ , the connectedness of  $I_1$  implies that  $T_1 \cap T_2 \neq \emptyset$ . But this is impossible, since  $p_1(t) + p_2(t) = h''(t) \neq 0$ . Therefore there exists a point  $t_0 \in I_1$  such that  $p_i(t_0) \neq 0$  for  $i = 1, 2$ . Let  $p(t) = \min_{i=1,2} p_i(t)$ . Then  $p(t_0) > 0$ , and by piecewise continuity of  $p_i$ , there is an interval  $I_2$  such that  $t_0 \in I_2 \subset I$  and  $t \in I_2 \Rightarrow p(t) > 0$ . Therefore,  $p = \min_{i=1,2} p_i > 0$  almost everywhere. Therefore,  $\int_a^x p(t) dt$  is a strictly increasing function of  $x$ , on  $I_2$ . As in the theorem cited above, select  $x_1, \dots, x_4$  so that  $f_0$  assumes four different positive values, and define  $y_1'(x)$  from  $f_0(x)$  rather than  $\varphi_1'(x)$ , as in that theorem. Letting  $y_2'(-t) = -y_1'(t)$ , we then define  $\{\sigma_1, \sigma_2\}$  and  $\{\psi_1, \psi_2\}$ . It is clear that  $f_0(t) \leq f_i(t)$ ,  $i = 1, 2$ , and

$$0 \leq f_0(t') - f_0(t) \leq f_i(t') - f_i(t)$$

for  $t < t'$ , so  $y$  certainly satisfies the necessary conditions for admissibility. Thus, the nontrivial decomposition has been accomplished.

**LEMMA 4.** *If an admissible pair  $\{\varphi_1, \varphi_2\}$  satisfying  $\varphi_1(-t) + \varphi_2(t) = h(t)$ , for all real  $t$ , where  $h(t) = \sqrt{1+t^2} w(\text{arc cot } t)$ , and  $w(\theta)$  has continuous second derivative is decomposable, there is an interval on which both  $\varphi_1(-t)$  and  $\varphi_2(t)$  are nonlinear.*

*Proof.* If  $y_i(t)$  is a function such that  $\sigma_i(t) = \varphi_i(t) + y_i(t)$  and  $\psi_i(t) = \varphi_i(t) - y_i(t)$  are convex and satisfy

$$\sigma_2(t) + \sigma_1(-t) = h(t) ,$$

and

$$\psi_2(t) + \psi_1(-t) = h(t) ,$$

then  $y_1(t) + y_2(-t) = 0$ . This implies that, if on an interval  $I$ , say  $\varphi_1(t)$  is linear, then for every  $x, x' \in I$  with  $x' > x$ ,

$$|y_1'(x') - y_1'(x)| \leq \varphi_1'(x') - \varphi_1'(x) = 0 ,$$

and also  $|y'_2(-x') - y'_2(-x)| = 0$ . Therefore,  $y'_1$  is constant on  $I$ , and  $y'_2$  is constant on  $-I$ . Suppose on every interval  $\varphi_1(-t)$  or  $\varphi_2(t)$  is linear. Then  $y'_i$  is constant on every interval, and continuous on the line, so it is identically a constant. Therefore,  $\sigma_i(t)$  and  $\psi_i(t)$  differ from a multiple of  $\varphi_i(t)$  at most by a linear function, and are therefore translates of multiples of  $\varphi_i(t)$ . Therefore,  $\{\varphi_1, \varphi_2\}$  is indecomposable.

The following theorem is therefore immediate.

**THEOREM 1.** *An admissible pair  $\{\varphi_1, \varphi_2\}$  satisfying  $\varphi_1(-t) + \varphi_2(t) = h(t)$ , for all real  $t$ , where  $h(t) = \sqrt{1 + t^2} w(\text{arc cot } t)$ ,  $w(\theta)$  having continuous second derivative, is indecomposable if and only if there is a family  $\{I_\alpha\}$  of disjoint intervals, such that  $\bigcup_\alpha I_\alpha = R^1$  and on every  $I_\alpha$  exactly one of  $\varphi_1(t), \varphi_2(-t)$  is linear.*

We can now characterize the indecomposable sets with a given width function.

**THEOREM 2.** *Let  $K(w)$  be the family of plane convex bodies with width function a multiple of  $w(\theta)$ , where  $w(\theta)$  has a continuous second derivative. The  $K$ -indecomposable members of  $K$  are those sets with the property that at least one of the support lines of the set in every direction goes through a corner point.*

*Proof.* Let  $K$  be a body with width function  $w(\theta)$ , and with corresponding supporting admissible pair  $\{\varphi_1, \varphi_2\}$ . Let  $y = f(x)$  be the function satisfied by the boundary of  $K$  in a neighborhood of a corner point  $(x_0, y_0)$  of  $K$ . Then the one-sided tangents, or extreme rays of the bundle of support lines, will have slopes  $f'_-(x_0), f'_+(x_0)$ . The support function  $\varphi_2(t) = \max_{(x,y) \in K} [xt + y]$  is linear on  $[t_1, t_2]$  if and only if for every  $t \in [t_1, t_2]$  there is a point  $(x_0, y_0)$  such that  $f'_-(x_0) + t \leq 0 \leq f'_+(x_0) + t$ , or  $f'_-(x_0) = t_2$ , and  $f'_+(x_0) = -t_1$ . The value of the function  $\varphi_2(t)$  on  $[t_1, t_2]$  is then  $\varphi_2(t) = x_0 t + y_0$ . The function  $\varphi_1(t) = \max_{(x,y) \in K} [xt - y]$  is, correspondingly, linear on a line segment  $[t_1, t_2]$  when there is a point  $(x_0, y_0) \in K$  such that  $f'_-(x_0) = t_1, f'_+(x_0) = t_2$ . Therefore by Theorem 1,  $K$  is  $K$ -indecomposable if and only if at least one of the two support lines in every direction goes through a corner point.

Of special interest is the case where  $w(\theta)$  is independent of the direction, i.e.,  $K(w)$  is the family of bodies of constant width.

Applying the results of the preceding discussion we note that the admissible pair  $\{\varphi_1, \varphi_2\}$  of support functions corresponding to a body of constant width  $r$  with origin in its interior satisfies  $\varphi_1(-t) + \varphi_2(t) = r\sqrt{1 + t^2}$ , for all real  $t$ .

Applying the results of Theorem 1 a body of constant width is  $K$ -indecomposable if and only if at least one of the two support lines in every direction goes through a corner point.

We note that if  $K$ , a body of constant width  $r$ , has a corner point  $Q$ , where the bundle of supporting lines occupies an angle equal to  $\pi - \theta$ , then opposite  $Q$  is an arc  $\widehat{pp'}$  with the property that every point on  $\widehat{pp'}$  is at distance  $r$  from  $Q$ , i.e.,  $\widehat{pp'}$  is a circular arc of radius  $r$  and angle  $\theta$ . Therefore, the boundary of  $K$  is entirely composed of circular arcs of radius equal to the width.

**THEOREM 3.** *Let  $K$  be the family of bodies of constant width. The  $K$ -indecomposable sets are precisely those whose boundary is entirely composed of circular arcs whose radius is equal to the width.*

Since a plane convex body of constant width  $r$ , whose boundary is composed entirely of arcs of radius  $r$ , must have a corner point opposite every arc, this result is immediate.

The author wishes to express her appreciation to her thesis advisor, V. Klee, to M. Kallay, and to others too numerous to mention, for their helpful suggestions and criticisms.

#### REFERENCES

1. W. Fenchel, *Convex Cones, Sets and Functions*, Princeton University, Princeton, 1943
2. P. R. Halmos, *Measure Theory*, D. Van Nostrand and Company, New York, 1950.
3. H. Rademacher and O. Toeplitz, *The Enjoyment of Mathematics*, translated by H. Zuckerman, Princeton University Press, Princeton, Chapter 25, pp. 163-176, 1957.
4. R. Silverman, *Decomposition of Plane Convex Sets, Part I: Machinery and Preliminary Results*, to appear.

Received November 6, 1970 and in revised form July 7, 1972.

LEHIGH UNIVERSITY

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)  
University of California  
Los Angeles, California 90024

J. DUGUNDJI\*  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT  
University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM  
Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$48.00 a year (6 Vols., 12 issues). Special rate: \$24.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

\* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by  
Pacific Journal of Mathematics  
All Rights Reserved

Mir Maswood Ali, <i>Content of the frustum of a simplex</i> .....	313
Mieczyslaw Altman, <i>Contractors, approximate identities and factorization in Banach algebras</i> .....	323
Charles Francis Amelin, <i>A numerical range for two linear operators</i> .....	335
John Robert Baxter and Rafael Van Severen Chacon, <i>Nonlinear functionals on <math>C([0, 1] \times [0, 1])</math></i> .....	347
Stephen Dale Bronn, <i>Cotorsion theories</i> .....	355
Peter A. Fowler, <i>Capacity theory in Banach spaces</i> .....	365
Jerome A. Goldstein, <i>Groups of isometries on Orlicz spaces</i> .....	387
Kenneth R. Goodearl, <i>Idealizers and nonsingular rings</i> .....	395
Robert L. Griess, Jr., <i>Automorphisms of extra special groups and nonvanishing degree 2 cohomology</i> .....	403
Paul M. Krajcikiewicz, <i>The Picard theorem for multianalytic functions</i> .....	423
Peter A. McCoy, <i>Value distribution of linear combinations of axisymmetric harmonic polynomials and their derivatives</i> .....	441
A. P. Morse and Donald Chesley Pfaff, <i>Separative relations for measures</i> .....	451
Albert David Polimeni, <i>Groups in which <math>\text{Aut}(G)</math> is transitive on the isomorphism classes of <math>G</math></i> .....	473
Aribindi Satyanarayan Rao, <i>Matrix summability of a class of derived Fourier series</i> .....	481
Thomas Jay Sanders, <i>Shape groups and products</i> .....	485
Ruth Silverman, <i>Decomposition of plane convex sets. II. Sets associated with a width function</i> .....	497
Richard Snay, <i>Decompositions of <math>E^3</math> into points and countably many flexible dendrites</i> .....	503
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable, IV</i> .....	511
Robert E. Waterman, <i>Invariant subspaces, similarity and isometric equivalence of certain commuting operators in <math>L_p</math></i> .....	593
James Chin-Sze Wong, <i>An ergodic property of locally compact amenable semigroups</i> .....	615
Julius Martin Zelmanowitz, <i>Orders in simple Artinian rings are strongly equivalent to matrix rings</i> .....	621