Pacific Journal of Mathematics

ON NUMERICAL RANGES OF ELEMENTS OF LOCALLY *m*-CONVEX ALGEBRAS

JOHN R. GILES AND DONALD OTTO KOEHLER

ON NUMERICAL RANGES OF ELEMENTS OF LOCALLY *m*-CONVEX ALGEBRAS

J. R. GILES AND D. O. KOEHLER

The concept of numerical range is extended from normed algebras to locally *m*-convex algebras. It is shown that the approximating relations between the numerical range and the spectrum of an element are preserved in the generalization. The set of elements with bounded numerical range is characterized and the relation between boundedness of the spectrum and of the numerical range is discussed. The Vidav-Palmer theory is generalized to give a characterization of b^* -algebras by numerical range.

In a complex unital Banach algebra the numerical range of an element is a set of complex numbers which can be used to approximate the spectrum of an element. In a complex locally *m*-convex algebra with identity, for each element we define a set of numerical ranges and establish similar approximation to the spectrum of the element. In a normed algebra the spectrum and the numerical range of each element are bounded sets, but in a locally m-convex algebra the spectrum and the numerical ranges of an element may be unbounded. For a locally m-convex algebra with identity we characterize those elements with a bounded numerical range as an important normed subalgebra, and we discuss the relation between boundedness of the spectrum and the numerical ranges. In the normed algebra theory the study of hermitian elements, those with real numerical range, has led to the important Vidav-Palmer theory characterizing unital B^* -algebras among unital Banach algebras. We generalize the results of this theory to a characterization of b^* -algebras by numerical range.

We would like to thank Dr. T. Husain for the valuable discussions we have had with him on this subject. We would also like to express our appreciation to the referee for his valuable suggestions.

1. The numerical ranges of an element. For a complex normed algebra $(A, || \cdot ||)$ with identity 1 where ||1|| = 1, i.e., a complex unital normed algebra, we define the set

$$D(A, || \cdot ||; 1) \equiv \{f \in A' : f(1) = 1 \text{ and } ||f|| = 1\}.$$

For each $a \in A$ we define the numerical range of a as the set

$$V(A, || \cdot ||; a) \equiv \{f(a): f \in D(A, || \cdot ||; 1)\},\$$

and the numerical radius of a as

$$v(A, || \cdot ||; a) \equiv \sup \{|\lambda| : \lambda \in V(A, || \cdot ||; a) .$$

The set $D(A, ||\cdot||; 1)$ is a convex weak * compact subset of A' and the numerical range $V(A, ||\cdot||; a)$ is also a convex compact subset of the complex numbers, [2, p. 16]. The properties and applications of normed algebra numerical ranges have been studied extensively and the main results are conveniently presented by F. F. Bonsall and J. Duncan in [2].

Locally *m*-convex algebras, i.e., l.m.c. algebras, are examined in some detail by E. A. Michael in [4]. We call a l.m.c. algebra with identity a *unital l.m.c. algebra*. It is our aim to extend the concept of numerical range from complex unital normed algebras to complex unital l.m.c. algebras. It is sufficient for our purpose to note that, for a given l.m.c. algebra A with identity 1 there exists a separating family of submultiplicative semi-norms $\{p_{\alpha}\}$ on A which generates the topology and is such that $p_{\alpha}(1) = 1$ for all α , [3, p. 7]. Given such an algebra, we denote by P(A) the class of all such families of seminorms on A, and by $(A, \{p_{\alpha}\})$ the algebra A with a particular family of semi-norms $\{p_{\alpha}\} \in P(A)$.

Given $(A, \{p_{\alpha}\})$, for each α let N_{α} denote the nullspace of p_{α} , A_{α} denote the quotient space A/N_{α} , and $|| \cdot ||_{\alpha}$ denote the norm on A_{α} defined by $|| x + N_{\alpha} ||_{\alpha} = p_{\alpha}(x)$. For each α consider the natural linear mapping $x \mapsto x_{\alpha} \equiv x + N_{\alpha}$ of A onto A_{α} . We note that 1_{α} is the identity in A_{α} and that $|| 1_{\alpha} ||_{\alpha} = 1$ for each α . Michael has given the significant result that A is isomorphic to a subalgebra of the product of the normed algebras $(A_{\alpha}, || \cdot ||_{\alpha})$, Proposition 2.7, [4, p. 13]. Using this characterization of 1.m.c. algebras we are able to generalize much of the numerical range theory for normed algebras directly to a theory of numerical range for 1.m.c. algebras.

Given $(A, \{p_{\alpha}\})$, we define the set

$$D_{lpha}(A,\ p_{lpha};1)\equiv\{f\in A'\colon f(1)=1 \ ext{ and } |f(x)|\leq p_{lpha}(x) \ ext{for all } x\in A\} \ ,$$

and we write

$$D(A, \{p_{\alpha}\}; 1) \equiv \bigcup_{\alpha} \{D_{\alpha}(A, p_{\alpha}; 1)\}$$
.

For each $a \in A$ we write

$$V_{\alpha}(A, p_{\alpha}; a) \equiv \{f(a): f \in D_{\alpha}(A, p_{\alpha}; 1)\},\$$

and define the numerical range of a as the set

$$V(A, \{p_{\alpha}\}; a) \equiv \bigcup_{\alpha} \{V_{\alpha}(A, p_{\alpha}; a)\}$$
.

To each linear functional f on (A, p_{α}) which annihilates N_{α} , we can define the linear functional F on A_{α} by $F(x_{\alpha}) = f(x)$, and to each linear functional F on A_{α} we can define the linear functional f on (A, p_{α}) by $f(x) = F(x_{\alpha})$. Consequently, from the definition of the norm in A_{α} we see that $D_{\alpha}(A, p_{\alpha}; 1)$ is isomorphic to $D(A_{\alpha}, || \cdot ||_{\alpha}; 1_{\alpha})$, and for $a \in A$

$$V_{\alpha}(A, p_{\alpha}; a) = V(A_{\alpha}, || \cdot ||_{\alpha}; a_{\alpha})$$
.

Hence, we have the numerical range of a characterized by the normed algebra numerical ranges of the a_{α} in that

$$V(A, \{p_{lpha}\}; a) = igcup_{lpha} \left\{ V(A_{lpha}, || \cdot ||_{lpha}; a_{lpha})
ight\}$$
 .

Both $D(A, \{p_{\alpha}\}; 1)$ and $V(A, \{p_{\alpha}\}; a)$ depend upon the particular family of semi-norms $\{p_{\alpha}\} \in P(A)$ chosen to associate with A. It is clear that when $\{p_{\alpha}\}$ is a directed family, $D(A, \{p_{\alpha}\}; 1)$ is a convex subset of A' and the numerical range $V(A, \{p_{\alpha}\}; a)$ is a convex subset of the complex numbers.

For each $a \in A$ we write

$$v_{lpha}(A,\ p_{lpha}; a) \equiv \sup \left\{ \mid \lambda \mid : \lambda \in V_{lpha}(A,\ p_{lpha}; a)
ight\}$$
 ,

and we define the numerical radius of a as

$$v(A, \{p_{\alpha}\}; a) \equiv \sup \{|\lambda| : \lambda \in V(A, \{p_{\alpha}\}; a)\}$$
.

We note that $v_{\alpha}(A, p_{\alpha}; a) \leq p_{\alpha}(a)$ for each α , and we allow $v(A, \{p_{\alpha}\}; a) = \infty$. We have that

$$egin{aligned} v(A, \{p_lpha\}; a) &= \sup_lpha v_lpha(A, p_lpha; a) \ &= \sup_lpha v(A_lpha, || \cdot ||_lpha; a_lpha) \ . \end{aligned}$$

It is clear that the numerical range and the numerical radius have the following properties. For $a \in A$ and λ , μ complex

$$V(A, \{p_{\alpha}\}; \lambda a + \mu) = \lambda V(A, \{p_{\alpha}\}; a) + \mu$$

and

$$v(A, \{p_{\alpha}\}; \lambda a + \mu) \leq |\lambda| v(A, \{p_{\alpha}\}; a) + |\mu|$$
,

and for $a, b \in A$

$$V(A, \{p_{\alpha}\}; a + b) \subseteq V(A, \{p_{\alpha}\}; a) + V(A, \{p_{\alpha}\}; b)$$

and

$$v(A, \{p_{\alpha}\}; a + b) \leq v(A, \{p_{\alpha}\}; a) + v(A, \{p_{\alpha}\}; b)$$
.

2. The numerical ranges and the spectrum. In a unital Banach

algebra the numerical range of an element approximates its spectrum. We now establish similar approximating relations between the numerical ranges and the spectrum of an element in a complete unital l.m.c. algebra.

We recall that, given an algebra A with identity, for each $a \in A$, the *spectrum of a* is defined as the set

$$\sigma(A; a) \equiv \{\lambda: a - \lambda \text{ is not invertible}\}$$

THEOREM 1. Let A be a complete unital l.m.c. algebra. Given $(A, \{p_{\alpha}\})$, for each $a \in A$

$$\sigma(A, a) \subseteq V(A, \{p_{\alpha}\}; a) .$$

Proof. For each α let \overline{A}_{α} denote the completion of A_{α} . We have from Corollary 5.3(a), [4, p. 22] that

$$\sigma(A; a) = \bigcup_{\alpha} \sigma(\overline{A}_{\alpha}; a_{\alpha})$$
.

But from Theorem 2.6, [2, p. 19] we have that

$$\sigma(\overline{A}_{\alpha}; a) \subseteq V(\overline{A}_{\alpha}, || \cdot ||_{\alpha}; a_{\alpha})$$

and from Theorem 2.4, [2, p. 16] that

$$V(\overline{A}_{lpha}, || \cdot ||_{lpha}; a_{lpha}) = V(A_{lpha}, || \cdot ||_{lpha}; a_{lpha})$$

so it follows that

$$egin{aligned} \sigma(A;\,a) & \sqsubseteq igcup_lpha \left\{ V(A_lpha,\,|| \,ullet\,\,||_lpha;\,a_lpha)
ight\} \ & = V(A,\,\{p_lpha\};\,a) \;. \end{aligned}$$

THEOREM 2. Let A be a complete unital l.m.c. algebra. For each $a \in A$

$$\operatorname{co} \sigma(A; a) \subseteq \bigcap \{ V(A, \{p_{\alpha}\}; a) \colon \{p_{\alpha}\} \in P(A) \} \subseteq \overline{\operatorname{co}} \sigma(A; a) .$$

Proof. From Theorem 1 we have that

$$\operatorname{co} \sigma(A; a) \subseteq \bigcap \{ V(A, \{p_{\alpha}\}, a); \{p_{\alpha}\} \in P(A) \}$$

If $\operatorname{co} \sigma(A; a)$ is not all the complex plane then, for any $\lambda \notin \overline{\operatorname{co}} \sigma(A; a)$ there exists an open disc D_{λ} center λ such that D_{λ} can be strictly separated from $\overline{\operatorname{co}} \sigma(A; a)$ by a straight line L. Since

$$\sigma(A; a) = \bigcup_{\alpha} \sigma(\overline{A}_{\alpha}; a_{\alpha})$$
,

83

 D_{λ} is strictly separated from $\sigma(\bar{A}_{\alpha}; a_{\alpha})$ for any α , by the straight line L. However, for each α , $\sigma(\bar{A}_{\alpha}; a_{\alpha})$ is a compact set so there exists an open disc $D_{\alpha} \supseteq \sigma(\bar{A}_{\alpha}; a_{\alpha})$ which is strictly separated from D_{λ} by the same straight line L. We have from [2, p. 23] that, for each α , there exists a norm $|| \cdot ||'_{\alpha}$ equivalent to $|| \cdot ||_{\alpha}$ on \bar{A}_{α} such that

$$\sigma(\bar{A}_{\alpha}; a_{\alpha}) \subseteq V(\bar{A}_{\alpha}, || \cdot ||_{\alpha}'; a_{\alpha}) \subseteq D_{\alpha}$$
.

Now for each α ,

$$V(ar{A}_{lpha},\,||ullet\,||_{lpha}';\,a_{lpha})\,=\,V(A,\,||ullet\,||_{lpha}';\,a_{lpha})$$
 .

Defining the semi-norm p'_{α} on A by

$$p'_{lpha}(x)\,=\,||\,x_{lpha}\,||'_{lpha}$$
 ,

it is clear that the family $\{p'_{\alpha}\} \in P(A)$, and

$$V(A, \{p'_{lpha}\}; a) = \bigcup_{lpha} \{V(A_{lpha}, || \cdot ||'_{lpha}; a_{lpha})\}$$
.

So D_{λ} is strictly separated from $V(A, \{p'_{\alpha}\}, a)$ by the straight line L. It follows that D_{λ} is strictly separated from $\bigcap \{V(A, \{p_{\alpha}\}; a): \{p_{\alpha}\} \in P(A)\}$, and this implies that

$$\bigcap \{ V(A, \{p_{\alpha}\}; a) \colon \{p_{\alpha}\} \in P(A) \} \subseteq \overline{\operatorname{co}} \sigma(A; a) .$$

3. Elements with bounded numerical range. We now establish an important set of inequalities which are generalizations of an inequality from the normed algebra theory, and we use them to characterize elements with bounded numerical range.

LEMMA 1. Let A be a unital l.m.c. algebra. Given $(A, \{p_{\alpha}\})$, for $a \in A$ and each α

$$v(A, \{p_{lpha}\}, a) \geqq rac{1}{e} p_{lpha}(a)$$
 .

Proof. From Theorem 4.1, [2, p. 34] we have, for each α

$$v(A_{lpha}, || ullet ||_{lpha}; \, a_{lpha}) \geqq rac{1}{e} || \, a_{lpha} \, ||_{lpha} \; .$$

 \mathbf{S}_{0}

$$egin{aligned} v(A, \{p_lpha\}, a) &= \sup_lpha v(A_lpha, || ullet ||_lpha; a_lpha) \ &\geq rac{1}{e} || \, a_lpha \, ||_lpha &= rac{1}{e} \, p_lpha(a) \; , \end{aligned}$$

for each α .

From the fact that every $\{p_a\} \in P(A)$ is a separating family we can make the following deduction.

COROLLARY 1. If for a given $a \in A$, there exists an $(A, \{p_{\alpha}\})$ such that $V(A, \{p_{\alpha}\}; a) = \{0\}$ then a = 0.

We can also make a statement about elements with bounded numerical range.

COROLLARY 2. Given $(A, \{p_{\alpha}\})$, if for $a \in A$, $V(A, \{p_{\alpha}\}; a)$ is bounded then $\sup_{\alpha} p_{\alpha}(a) < \infty$.

For the characterization of the set of elements with bounded numerical range we also use the following lemma.

LEMMA 2. Let A be a unital l.m.c. algebra. Given $(A, \{p_{\alpha}\})$ we have, for each $a \in A$

$$\sup \operatorname{Re} V(A, \{p_{\alpha}\}; a) = rac{\inf_{\lambda>0}}{\lim_{\lambda\to 0+}} \left\{ rac{1}{\lambda} \left\{ \sup_{\alpha} p_{\alpha}(1 + \lambda a) - 1 \right\} \right\}.$$

Proof. It is clear that for any $f \in D(A, \{p_{\alpha}\}; 1)$ and $\lambda > 0$

$$\operatorname{Re} f(a) \leq \frac{1}{\lambda} \left\{ \sup_{\alpha} p_{\alpha}(1 + \lambda a) - 1 \right\}$$

and therefore,

(1)
$$\sup \operatorname{Re} V(A, \{p_{\alpha}\}, a) \leq \inf_{\lambda > 0} \frac{1}{\lambda} \left\{ \sup_{\alpha} p_{\alpha}(1 + \lambda a) - 1 \right\}.$$

It follows that the result holds when $V(A, \{p_{\alpha}\}; a)$ is unbounded. We consider the case when $V(A, \{p_{\alpha}\}; a)$ is bounded, and write for every α

$$\tau_{\alpha} \equiv \sup \operatorname{Re} V_{\alpha}(A_{\alpha}, || \cdot ||_{\alpha}; a_{\alpha})$$
,

and

 $\tau \equiv \sup \operatorname{Re} V(A, \{p_{\alpha}\}; a)$.

Now by [2, p. 18], for every α

$$rac{1}{\lambda} \{ || \, 1_lpha + \lambda a_lpha \, ||_lpha - 1 \} \leq (1 - \lambda au_lpha)^{-1} \{ au_lpha + \lambda || \, a_lpha^2 \, ||_lpha \} \; ,$$

when $0 < \lambda < ||a_{\alpha}||_{\alpha}^{-1}$. Since $V(A, \{p_{\alpha}\}; \alpha)$ is bounded we have from Corollary 2 to Lemma 1 that there exists an M > 0 such that $M \ge 1$

 $\sup_{\alpha} p_{\alpha}(a)$. Then, for every α

$$\frac{1}{\lambda} \{ p_{\alpha}(1 + \lambda \alpha) - 1 \} \leq (1 - \lambda \tau)^{-1} \{ \tau + \lambda M^2 \}$$

when $0 < \lambda < 1/M$. Therefore,

$$\lim_{\lambda \to 0^+} rac{1}{\lambda} \left\{ \sup_{lpha} p_{lpha} (1 + \lambda a) - 1
ight\} \leq au$$
 .

Together with inequality (1), this completes the proof.

Let A be a unital l.m.c. algebra. Given $(A, \{p_{\alpha}\})$ we can define the subalgebra

$$B \equiv \left\{ x \in A \colon \sup_{\alpha} p_{\alpha}(x) < \infty \right\}$$
.

Now $p(x) \equiv \sup_{\alpha} p_{\alpha}(x)$ is a norm for B since $\{p_{\alpha}\}$ is a separating family and we note that $1 \in B$ and p(1) = 1.

It can be seen from the proof of Theorem 2.3, [1, p. 32], that if A is a complete unital l.m.c. algebra then given $(A, \{p_{\alpha}\})$, the normed subalgebra (B, p) is complete. However, an examination of the sequence $\{x_n\}$ where $x_n = \{1, 2, 3, \dots, n, \dots, n, \dots\}$, in the algebra A of Example 2 below, shows that there exists an incomplete unital l.m.c. algebra A with $\{p_{\alpha}\} \in P(A)$ such that (B, p) is complete.

Given $(A, \{p_{\alpha}\})$, we can characterize elements with bounded numerical range as elements of (B, p).

THEOREM 3. If A is a unital l.m.c. algebra then given $(A, \{p_{\alpha}\})$,

 $B = \{x \in A: V(A, \{p_{\alpha}\}; x) \text{ is bounded}\}$

and when $\{p_{\alpha}\}$ is a directed family, for every $a \in B$

$$\overline{V(A, \{p_{\alpha}\}; a)} = V(B, p; a)$$
.

Proof. If, for a given $a \in A$, $V(A, \{p_{\alpha}\}; a)$ is bounded then Corollary 2 to Lemma 1 implies that $a \in B$. If $a \in B$ then $\sup_{\alpha} p_{\alpha}(1 + \lambda a) = p(1 + \lambda a)$ for all λ , so

$$\begin{split} \sup \operatorname{Re} \, V(A, \{p_{\alpha}\}; a) &= \frac{\inf_{\lambda > 0}}{\lim_{\lambda \to 0+}} \bigg\} \frac{1}{\lambda} \{ p(1 + \lambda a) - 1 \} \\ &= \sup \operatorname{Re} \, V(B, \, p; a) \; . \end{split}$$

Hence, since $V(B, p; \lambda a) = \lambda V(B, p; a)$ and $V(A, \{p_{\alpha}\}; \lambda a) = \lambda V(A, \{p_{\alpha}\}; a)$ for all λ complex, $|\lambda| = 1$, we deduce that every $a \in B$ has bounded numerical range $V(A, \{p_{\alpha}\}; a)$. When $\{p_{\alpha}\}$ is a directed family, both

numerical ranges are convex sets so we deduce from the Krein-Milman Theorem that for every $a \in B$

$$\overline{V(A, \{p_{\alpha}\}; a)} = V(B, p; a)$$
.

The following result relates boundedness of the spectrum to boundedness of the numerical range.

THEOREM 4. Let A be a complete unital l.m.c. algebra. For any $a \in A$, $\sigma(A; a)$ is bounded if and only if there exists an $(A, \{p_{\alpha}\})$ such that $V(A, \{p_{\alpha}\}; a)$ is bounded.

Proof. If for $a \in A$ there exists an $(A, \{p_{\alpha}\})$ such that $V(A, \{p_{\alpha}\}; a)$ is bounded, then it follows from Theorem 1 that $\sigma(A; a)$ is bounded.

Conversely, consider $a \in A$ with $\sigma(A; a)$ bounded. There exists a disc D in the complex plane such that $\sigma(A; a) \subseteq D$. Now $\sigma(A; a) = \bigcup_{\alpha} \sigma(\overline{A}_{\alpha}; a_{\alpha})$. From [2, p. 23], for each α there exists a norm $|| \cdot ||_{\alpha}'$ equivalent to $|| \cdot ||_{\alpha}$ on \overline{A}_{α} such that

$$\sigma(\bar{A}_{lpha}; a_{lpha}) \subseteq V(\bar{A}_{lpha}; || \cdot ||_{lpha}'; a_{lpha}) \subseteq D$$
 .

For each α , defining the semi-norm p'_{α} on A by

$$p'_lpha(x) = ||x_lpha||'_lpha$$
 ,

the family $\{p'_{\alpha}\} \in P(A)$ and

$$\sigma(A; a) \subseteq V(A, \{p'_{\alpha}\}; a) = \bigcup_{\alpha} V(A_{\alpha}, || \cdot ||'_{\alpha}, a_{\alpha}) \subseteq D.$$

Further to the relation between boundedness of the spectrum and the numerical range given in Theorem 4, the following example shows that there exist l.m.c. algebras A where $\sigma(A; a)$ is bounded for a given $a \in A$ but where there exists $(A, \{p_{\alpha}\})$ such that $V(A, \{p_{\alpha}\}; a)$ is unbounded.

EXAMPLE 1. Let A be the algebra of all sequences of complex numbers, $x \equiv \{\lambda_1, \lambda_2, \dots, \lambda_n, \dots\}$ with pointwise definition of addition and multiplication by a scalar, but with convolution multiplication and with unit $1 \equiv \{1, 0, \dots, 0, \dots\}$. A sequence of submultiplicative semi-norms $\{p_n\}$ is defined on A by

$$p_n(x) = \sum_{k=1}^n |\lambda_k|$$
 ,

and the sequence satisfies $p_n(1) = 1$ for all n, and is separating. Consider $a \in A$ such that $\lambda_n \to 0$. Then $p_n(a) \to \infty$, so by Lemma 1 $V(A, \{p_n\}; a)$ is unbounded. But $\sigma(A; a) = \{\lambda_1\}$, which is bounded.

It is worth noting that a complete unital l.m.c. algebra with a

bounded numerical range property has the following property.

THEOREM 5. If A is a complete unital l.m.c. algebra where there exists an $(A, \{p_a\})$ such that $V(A, \{p_a\}; a)$ is bounded for all $a \in A$, then $\sigma(A; a)$ is compact for all $a \in A$.

Proof. We note that B = A and so $\sigma(A; a) = \sigma(B; a)$, for every $a \in A$. Since A is complete it follows that (B, p) is complete and so $\sigma(B; a)$ is compact for every $a \in A$.

However, the algebra A of Example 1 has $\sigma(A; a)$ compact for all $a \in A$ but there exists an $(A, \{p_{\alpha}\})$ such that $V(A, \{p_{\alpha}\}; a)$ is not bounded for all $a \in A$.

It is known that there exist non-normable l.m.c. algebras where $\sigma(A; a)$ is compact for all $a \in A$, [4, p. 80]. The following example gives the further information that there exist non-normable l.m.c. algebras A where, for a certain $(A, \{p_{\alpha}\})$, $V(A, \{p_{\alpha}\}; a)$ is bounded for all $a \in A$.

EXAMPLE 2. Let A be the algebra l^{∞} of all bounded sequences of complex numbers, $x \equiv \{\lambda_1, \lambda_2, \dots, \lambda_n, \dots\}$, with pointwise definition of the algebra operations and with unit $1 \equiv \{1, 1, \dots, 1, \dots\}$. A sequence of submultiplicative semi-norms $\{p_n\}$ is defined on A by

$$p_n(x) = |\lambda_n|$$
,

and the sequence satisfies $p_n(1) = 1$ for all n, and is separating. Now p, defined by

$$p(x) \equiv \sup_{n} \left\{ p_n(x) \right\}$$

is the usual l^{∞} -norm on A, so B = A and from Theorem 3, $V(A, \{p_n\}; a)$ is bounded for all $a \in A$. However, it is clear that $(A, \{p_n\})$ is non-normable.

4. A characterization of b^* -algebras by numerical range. A *l.m.c.* * algebra A is a l.m.c. algebra with a continuous involution *. We let S(A) denote the set $\{x \in A : x = x^*\}$, the selfadjoint elements of A. A b^* -algebra A is a complete l.m.c. * algebra where there exists a family $\{p_{\alpha}\} \in P(A)$ such that $p_{\alpha}(x^*x) = p_{\alpha}(x)^2$ for all $x \in A$ and every α , [1, p. 31].

In a unital normed algebra $(A, || \cdot ||)$, the set of hermitian elements $H(A, || \cdot ||)$ is the set of elements a with real numerical range $V(A, || \cdot ||; a)$. For a unital l.m.c. algebra A, given $(A, \{p_{\alpha}\})$ we define the set of hermitian elements $H(A, \{p_{\alpha}\})$ as the set of elements a with real numerical range $V(A, \{p_{\alpha}\}; a)$. It is clear from the definition of

the numerical range in $(A, \{p_{\alpha}\})$ that $a \in H(A, \{p_{\alpha}\})$ if and only if $a_{\alpha} \in H(A_{\alpha}, || \cdot ||_{\alpha})$ for all α .

One of the outstanding successes of the normed algebra numerical range theory is the Vidav-Palmer Theorem [2, p. 65] which characterizes unital B^* algebras as unital Banach algebras which have an hermitian decomposition. We now consider a generalization of this work to the characterization of unital b^* -algebras amongst the complete unital l.m.c. algebras.

We need the following property of the hermitian elements.

LEMMA 3. Let A be a complete unital l.m.c. algebra. Given $(A, \{p_{\alpha}\})$, the set $H(A, \{p_{\alpha}\})$ is closed.

Proof. Consider h a cluster point of $H(A, \{p_{\alpha}\})$. Then, for each α , h_{α} is a cluster point of $H(A_{\alpha}, || \cdot ||_{\alpha})$. But by Lemma 7, [5, p. 198], $H(A_{\alpha}, || \cdot ||_{\alpha})$ is closed in $(A_{\alpha}, || \cdot ||_{\alpha})$, so $h_{\alpha} \in H(A_{\alpha}, || \cdot ||_{\alpha})$. Since $V(A, \{p_{\alpha}\}; h) = \bigcup_{\alpha} V(A_{\alpha}, || \cdot ||_{\alpha}; h_{\alpha})$ we have that $h \in H(A, \{p_{\alpha}\})$.

THEOREM 6. Let A be a complete unital l.m.c. algebra. Given $(A, \{p_a\})$, the following statements are equivalent.

(i) $A = H(A, \{p_{\alpha}\}) + iH(A, \{p_{\alpha}\}), a \text{ direct sum,}$

(ii) There is an involution * on A such that A is a l.m.c. * algebra where $S(A) = H(A, \{p_{\alpha}\}),$

(iii) There is an involution * on A such that A is * isomorphic to a * subalgebra of a product of B^* -algebras $(\overline{A}_{\alpha}, || \cdot ||_{\alpha})$,

(iv) There is an involution * on A such that A is a b^* -algebra,

(v) There is an involution on B such that (B, p) is a dense B^* -algebra.

Proof. (i) \Rightarrow (ii) Since $A = H(A, \{p_{\alpha}\}) + iH(A, \{p_{\alpha}\})$, we define the involution * on A as follows: for x = h + ik where $h, k \in H(A, \{p_{\alpha}\})$ put $x^* = h - ik$. We need to show that * is continuous on A. Now for every α , $A_{\alpha} = H(A_{\alpha}, || \cdot ||_{\alpha}) + iH(A_{\alpha}, || \cdot ||_{\alpha})$ and * induces an involution * on A_{α} where for $x_{\alpha} = h_{\alpha} + ik$ we have $x_{\alpha}^* = h_{\alpha} - ik_{\alpha}$. But from Lemma 5.8, [2, p. 50], since $A_{\alpha} \subseteq J(\overline{A}_{\alpha})$, we have for every α , that * is continuous on A_{α} and since $p_{\alpha}(x) = ||x_{\alpha}||_{\alpha}$ for all $x \in A$, * is continuous on A. It is clear that with this involution *, $S(A) = H(A, \{p_{\alpha}\})$.

(ii) \Rightarrow (iii) Since $H(A, \{p_{\alpha}\}) = S(A)$ we have $A = H(A, \{p_{\alpha}\}) + iH(A, \{p_{\alpha}\})$ and so, for every α , $A_{\alpha} = H(A_{\alpha}, || \cdot ||_{\alpha}) + iH(A_{\alpha}, || \cdot ||_{\alpha})$. But by Theorem 8.2, [2, p. 74], A_{α} is a pre-*B**-algebra and so \overline{A}_{α} is a *B**-algebra for every α . Our result follows from Michael's characterization of l.m.c. algebras, Proposition 2.7, [4, p. 13].

(iii) \Rightarrow (iv) For every α , since $(\bar{A}_{\alpha}, || \cdot ||_{\alpha})$ is a B^* -algebra and $p_{\alpha}(x) = || x_{\alpha} ||_{\alpha}$ for all $x \in A$, we have $p_{\alpha}(x^*x) = p_{\alpha}(x)^2$ for all $x \in A$; that is, A is a b^* -algebra.

 $(iv) \Rightarrow (v)$ This is proved as Theorem 2.3, [1, p. 32].

 $(\mathbf{v}) \Rightarrow (\mathbf{i})$ Since (B, p) is a unital B^* -algebra, B = H(B, p) + iH(B, p). For any $h, k \in H(B, p)$, we have from Theorem 3, that $h, k \in H(A, \{p_{\alpha}\})$. But then for every $\alpha, h_{\alpha}, k_{\alpha} \in H(A_{\alpha}, || \cdot ||_{\alpha})$ and since $|| x_{\alpha} ||_{\alpha} = p_{\alpha}(x)$ for all $x \in A$, we have from inequality (1), [2, p. 50] that

$$p_{lpha}(h) \leq e p_{lpha}(h+ik)$$
 .

This inequality implies that for any net $\{h_r + ik_r\}$ in *B* convergent to x in *A*, both $\{h_r\}$ and $\{k_r\}$ converge to say *h* and *k*. But by Lemma 3 the set $H(A, \{p_\alpha\})$ is closed in *A* so *h*, $k \in H(A, \{p_\alpha\})$ and x = h + ik. Since *B* is dense in *A*, we have $A = H(A, \{p_\alpha\}) + iH(A, \{p_\alpha\})$.

It should be noted that this theorem gives in $(v) \Rightarrow (iv)$, a converse to Theorem 2.3, [1, p. 32], and in $(iv) \Rightarrow (iii)$, a simpler proof for Theorem 2.4, [1, p. 32], by using numerical range techniques.

The following is an application of Theorem 6 and is a generalization of Theorem 7.6, [2, p. 71].

THEOREM 7. Let Ω be a locally compact Hausdorff space and let $A \equiv \mathscr{C}(\Omega)$ be the algebra of all complex continuous functions on Ω . If A is an F-algebra under the compact-open topology, then any l.m.c. topology generated by a family of semi-norms $\{p_{\alpha}\}$ such that $p_{\alpha}(f) = p_{\alpha}(|f|)$ for all $f \in A$ and $p_{\alpha}(1) = 1$ for all α , under which A is an F-algebra, is the compact-open topology.

Proof. We can introduce the exponential function in A, $\exp a = 1 + \sum_{n=1}^{\infty} (1/n!)a^n$, and it is clear that $(\exp a)_{\alpha} = \exp a_{\alpha}$, for every α . Now if g is a real continuous function on Ω then for real λ and for each α

$$egin{aligned} || \exp\left(i\lambda g_lpha
ight) ||_lpha &= || \left(\exp i\lambda g
ight)_lpha ||_lpha \ &= p_lpha(\exp i\lambda g) \ &= p_lpha(|\exp i\lambda g\,|) \ &= p_lpha(1) = 1 \;. \end{aligned}$$

Therefore, by Lemma 5.2, [2, p. 46], $g_{\alpha} \in H(A_{\alpha}, || \cdot ||_{\alpha})$ for every α , and so $g \in H(A, \{p_{\alpha}\})$. We have then $A = H(A, \{p_{\alpha}\}) + iH(A, \{p_{\alpha}\})$ and by Theorem 6 we conclude that A is a b*-algebra. Now, by Theorem 4.2, [1, p. 36], A with the compact-open topology is also a b*-algebra. So by Theorem 3.7, [1, p. 35], the l.m.c. topology generated by $\{p_{\alpha}\}$ is the compact-open topology.

In the above theorem we note that A with the compact-open topology and A with the l.m.c. topology generated by the semi-norms, must both be F-algebras; one of these being an F-algebra is not sufficient. We are indebted to the referee for the following examples which illustrate this point.

EXAMPLE 3. Let $\Omega \equiv [0, 1]$ with the usual topology and let $\{K_{\alpha}\}$ be the set of compact countable subsets of Ω . A family of semi-norms $\{p_{\alpha}\}$ defined on A by

$$p_{lpha}(x) = \sup_{x \in K_{lpha}} \left\{ \left| f(x) \right|
ight\}$$
 ,

satisfies the conditions of the theorem except that A with this topology is not an *F*-algebra, [4, Example 3.8, p. 19]. However, A with the compact-open topology is a Banach algebra, so it is clear that the topology generated by the family of semi-norms is not the compact-open topology.

EXAMPLE 4. Let Ω be the set of ordinal numbers smaller than the first uncountable ordinal, with the order topology. With norm pdefined on A by

$$p(x) = \sup_{x \in \mathcal{Q}} \left\{ \left| f(x) \right|
ight\}$$
 ,

A is a Banach algebra. However, A with the compact-open topology is not an F-algebra, [4, Example 3.7, p. 19], so it is clear that the norm topology on A is not the compact-open topology.

Note added in proof. We are indebted to Dr. R. T. Moore for pointing out, in connection with Example 2, that the following result can be deduced from Theorem 3 by the Open Mapping Theorem.

THEOREM. A unital F-algebra A is normable if and only if there exists an $(A, \{p_{\alpha}\})$ such that $V(A, \{p_{\alpha}\}; a)$ is bounded for all $a \in A$.

References

1. C. Apostol, b*-algebras and their representation, J. London Math. Soc., 2nd Series, 3 (1971), 30-38.

2. F. F. Bonsall and J. Duncan, Numerical Range of Operators on Normed Spaces and of Elements of Normed Algebras, London Math. Soc. Lecture Note Series 2, Cambridge, 1971.

3. R. M. Brooks, On locally m-convex *-algebras, Pacific J. Math, 23 (1967), 5-23.

4. E. A. Michael, Locally multiplicatively convex topological algebras, Amer. Math. Soc. Mem., **11** (1952).

5. B. Sims, A characterization of Banach-star-algebras by numerical range, Bull. Austral. Math. Soc., 4 (1971), 193-200.

ON NUMERICAL RANGES OF ELEMENTS OF LOCALLY *m*-CONVEX 91

Received August 21, 1972 and in revised form September 28, 1972. The work of the first author was partially supported under a visiting postdoctoral fellowship at McMaster University, Ontario. The work of the second author was supported under a visiting senior lectureship at the University of Newcastle, N.S.W.

UNIVERSITY OF NEWCASTLE AND MIAMI UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor) University of California Los Angeles, California 90024

R. A. BEAUMONT University of Washington Seattle, Washington 98105 J. DUGUNDJI* Department of Mathematics University of Southern California Los Angeles, California 90007

D. GILBARG AND J. MILGRAM Stanford University Stanford, California 94305

ASSOCIATE EDITORS

B. H. NEUMANN

E. F. BECKENBACH

F. Wolf

K. Yoshida

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON * * * * AMERICAN MATHEMATICAL SOCIETY

AMERICAN MATHEMATICAL SOCIETY NAVAL WEAPONS CENTER

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Pacific Journal of Mathematics May, 1973

Vol. 49, No. 1

A. Bigard, Free lattice-ordered modules	1
Richard Bolstein and Warren R. Wogen, <i>Subnormal operators in strictly cyclic</i> operator algebras	7
Herbert Busemann and Donald E. Glassco, II, <i>Irreducible sums of simple</i>	,
multivectors	13
W. Wistar (William) Comfort and Victor Harold Saks, <i>Countably compact groups</i> <i>and finest totally bounded topologies</i>	33
Mary Rodriguez Embry, Maximal invariant subspaces of strictly cyclic operator	
algebras	45
Ralph S. Freese and James Bryant Nation, <i>Congruence lattices of semilattices</i>	51
Ervin Fried and George Grätzer, A nonassociative extension of the class of	50
Laba D. Ciles and Denald Otta Kashlar. On summing language of denaute of the alle	39
John R. Giles and Donald Otto Koenier, On numerical ranges of elements of locally	70
Devid A Hill On dominant and codominant dimension of OE – 3 rings	03
John Sollion Heis and Pohert Poul Johnson Pound and Pfieter forms over $P(t)$	101
John Sonion Tisia and Robert Faul Johnson, <i>Round and Tisier Johns over</i> $R(t) \dots$	101
A then assiss G. Kortsetes and Edward Parry Soff. Hungmahungmial approximation	109
Anianassios O. Kansatos and Edward Barry San, Hyperpolynomial approximation	117
Shin'ichi Kinoshita. On alamantary ideals of θ curves in the 3 sphere and 2 links in	11/
the 4-sphere	127
Ronald Brian Kirk Convergence of Baire measures	135
R I Knill The Seifert and Van Kampen theorem via regular covering spaces	149
Amos A Kovacs. Homomorphisms of matrix rings into matrix rings	161
Voung K Kwon, HD-minimal but no HD-minimal	171
Makoto Maejima. On the renewal function when some of the mean renewal lifetimes	1/1
are infinite	177
Juan José Martínez, Cohomological dimension of discrete modules over profinite	
groups	185
W. K. Nicholson, Semiperfect rings with abelian group of units.	191
Louis Jackson Ratliff, Jr., Three theorems on imbedded prime divisors of principal	
ideals	199
Billy E. Rhoades and Albert Wilansky, Some commutants in $B(c)$ which are almost	
matrices	211
John Philip Riley Jr., Cross-sections of decompositions	219
Keith Duncan Stroyan, A characterization of the Mackey uniformity $m(L^{\infty}, L^{1})$ for finite measures	223
Edward G. Thurber. The Scholz-Brauer problem on addition chains	220
Ioze Vrahec, Submanifolds of acyclic 3-manifolds	243
Philip William Walker, Adjoint boundary value problems for compactified singular	213
differential operators	265
Roger P. Ware. When are Witt rings group rings	279
James D. Wine, <i>Paracompactifications using filter bases</i>	285