

Pacific Journal of Mathematics

SEMIPERFECT RINGS WITH ABELIAN GROUP OF UNITS

W. K. NICHOLSON

SEMI PERFECT RINGS WITH ABELIAN GROUP OF UNITS

W. K. NICHOLSON

In 1963, Gilmer characterized all finite commutative rings with a cyclic group of units and, in 1967, Eldridge and Fischer generalized these results to rings with minimum condition. In the present paper these results are extended to semiperfect rings and generalizations of the three theorems are obtained. It is shown that a semiperfect ring with cyclic group of units is finite and is either commutative or is the direct sum of a commutative ring and the 2×2 upper triangular matrix ring over the field of two elements. Let R be semiperfect with an abelian group of units. It is shown that R is finite if either the group of units is finite or the group of units is finitely generated and the Jacobson radical is nil.

The proofs of all these results depend on our main theorem: The structure of a semiperfect ring R with an abelian group of units is described completely up to the structure of commutative local rings. (That is commutative rings with a unique maximal ideal.) The groups of units of these local rings are shown to be direct factors of the group of units of R .

1. Preliminaries. Throughout this paper we assume that all rings are associative and have an identity and that all modules are unital. If R is a ring we denote its group of units by R^* and its Jacobson radical by $J(R)$. The ring of residues of the integers modulo n will be denoted by \mathcal{R}_n . The following notions will be referred to several times below.

DEFINITION 1. Let R_1, R_2, \dots, R_n be rings and, if $i \neq j$, let X_{ij} be an $R_i - R_j$ bimodule. We define the *semidirect sum* $[R_i, X_{ij}]$ to be the ring of all $n \times n$ "matrices" (x_{ij}) where $x_{ii} \in R_i$ for each i and $x_{ij} \in X_{ij}$ for all $i \neq j$. These are added componentwise and we define the product $(x_{ij})(y_{ij}) = (z_{ij})$ as follows:

$$\begin{aligned} z_{ii} &= x_{ii}y_{ii} && \text{for all } i = 1, 2, \dots, n, \\ z_{ij} &= x_{ii}y_{ij} + x_{ij}y_{jj} && \text{for all } i \neq j. \end{aligned}$$

It is easy to verify that $[R_i, X_{ij}]$ is an associative ring. If the bimodules X_{ij} are all zero the semidirect sum $[R_i, X_{ij}]$ reduces to the usual direct sum $R_1 \oplus R_2 \oplus \dots \oplus R_n$. More generally, we have that $[0, X_{ij}]$ is an ideal of $[R_i, X_{ij}]$ which squares to zero, and the quotient ring is isomorphic to $R_1 \oplus R_2 \oplus \dots \oplus R_n$. Clearly the direct sum of two of these semidirect sums is again a semidirect sum.

DEFINITION 2. Let R be a ring. A left R -module X is said to be G -unital if $ux = x$ for all $u \in R^*$ and all $x \in X$. A bimodule is called G -unital if it is G -unital as a left and as a right module.

Clearly 0 is a G -unital module and submodules and quotient modules of such modules are again of the same type. If X is a G -unital R -module we have $x + x = 0$ for every $x \in X$ since $-1 \in R^*$. In other words X is an elementary abelian 2-group.

PROPOSITION 1. Let R_1, R_2, \dots, R_n be rings and let X_{ij} be an $R_i - R_j$ bimodule for all $i \neq j$. The semidirect sum $[R_i, X_{ij}]$ has the following properties:

- (1) $[R_i, X_{ij}]^* = \{(x_{ij}) \mid x_{ii} \in R_i^* \text{ for each } i\}$.
- (2) $[R_i, X_{ij}]^*$ is abelian if and only if each R_i^* is abelian and each X_{ij} is G -unital.
- (3) If $[R_i, X_{ij}]^*$ is abelian it is isomorphic to the direct product of all the multiplicative groups R_i^* and all the additive groups X_{ij} .

Proof. (1) If (x_{ij}) is given and $x_{ii} \in R_i^*$ for each i it is easy to verify that $(x_{ij})^{-1} = (y_{ij})$ where $y_{ii} = x_{ii}^{-1}$ for each i and $y_{ij} = -x_{ii}^{-1}x_{ij}x_{jj}^{-1}$ for all $i \neq j$. The converse is clear.

(2) Suppose $[R_i, X_{ij}]^*$ is abelian. Then (1) and the definition of multiplication in $[R_i, X_{ij}]$ imply that each R_i^* is abelian. Now choose arbitrary elements $x_{ii}, y_{ii} \in R_i^*$ for each i and $x_{ij}, y_{ij} \in X_{ij}$ for all $i \neq j$. The units (x_{ij}) and (y_{ij}) commute so, for all $i \neq j$:

$$x_{ii}y_{ij} + x_{ij}y_{jj} = y_{ii}x_{ij} + y_{ij}x_{jj}.$$

If $x_{ij} = 0$ and $x_{jj} = 1$ this shows $x_{ii}y_{ij} = y_{ij}$. Similarly $x_{ij}y_{jj} = x_{ij}$ and it follows that each X_{ij} is G -unital.

Conversely: If $(x_{ij}), (y_{ij}) \in [R_i, X_{ij}]^*$ then, using (1), $x_{ii}y_{ii} = y_{ii}x_{ii}$. Furthermore, since the X_{ij} are G -unital, we have

$$x_{ii}y_{ij} + x_{ij}y_{jj} = y_{ij} + x_{ij} = x_{ij} + y_{ij} = y_{ii}x_{ij} + y_{ij}x_{jj}$$

for all $i \neq j$ and it follows that $[R_i, X_{ij}]^*$ is abelian.

(3) This follows easily from the definition of multiplication in $[R_i, X_{ij}]$ and the fact that each X_{ij} is G -unital.

PROPOSITION 2. Let R be a local ring. ($R/J(R)$ a divisor ring.)

- (1) If the group of units of R is abelian then R is commutative.
- (2) If R possesses a nonzero G -unital module then $R/J(R) \cong \mathcal{K}_2$.
- (3) If $R/J(R) \cong \mathcal{K}_2$ the G -unital R -modules are precisely the (additive) elementary abelian 2-groups.

Proof. (1) If $a, b \in J(R)$ then $1 + a, 1 + b \in R^*$ so $(1 + a)(1 + b) =$

$(1 + b)(1 + a)$. This implies $ab = ba$. If $a \in J(R)$ and $u \in R^*$ then $(1 + a)u = u(1 + a)$ so $au = ua$.

(2) Let $X \neq 0$ be a G -unital left R -module. If $x \in X$ and $a \in J(R)$ then $(1 + a)x = x$ so $ax = 0$. Hence X is a G -unital $R/J(R)$ module so we can assume R is a division ring. But then if $0 \neq r \in R$ we have that $(1 - r)x = 0$ for every $x \in X$. Since $X \neq 0$ this means $1 - r$ is not a unit so $r = 1$. Hence $R \cong \mathcal{K}_2$.

(3) If X is an elementary abelian 2-group then it is a vector space over \mathcal{K}_2 . Since $R/J(R) \cong \mathcal{K}_2$, R acts on X as follows: If $x \in X$ and $r \in R$ we have $rx = x$ if $r \in R^*$ and $rx = 0$ if $r \in J(R)$. Clearly, X is G -unital. Conversely: Every G -unital module is an elementary 2-group since $-1 \in R^*$ and the action is as described.

In the next section we shall use these results to characterize the semiperfect rings R where R^* is abelian.

2. The main theorem. Throughout this section R will denote a semiperfect ring with R^* abelian. It is well known [2, Th. 20, p. 159] that R is semiperfect if and only if we can write $1 = e_1 + e_2 + \dots + e_n$ where the e_i are orthogonal local idempotents. Hence each of the rings e_iRe_i is local and, if $i \neq j$, the ring e_iRe_j is an $e_iRe_i - e_jRe_j$ bimodule. For the moment let $[e_iRe_j]$ denote the set of all $n \times n$ "matrices" (x_{ij}) with the (i, j) entry x_{ij} drawn from e_iRe_j . This is a ring if ordinary matrix operations are used.

Define a map $\phi: R \rightarrow [e_iRe_j]$ by $\phi(r) = (e_i r e_j)$ for each $r \in R$. Then ϕ is clearly a homomorphism of additive groups and it is a ring homomorphism since the (i, j) entry of $\phi(r)\phi(s)$ is

$$\sum_k (e_i r e_k)(e_k s e_j) = e_i r (e_1 + e_2 + \dots + e_n) s e_j = e_i r s e_j.$$

Moreover ϕ is one-to-one. Indeed, if $\phi(r) = 0$ then $e_i r e_j = 0$ for all i, j and so $r = \sum_{i,j} e_i r e_j = 0$. Finally ϕ is onto. For if $(e_i r_{ij} e_j) \in [e_iRe_j]$ is given let $r = \sum_{i,j} e_i r_{ij} e_j$. It is easy to check that $\phi(r) = (e_i r_{ij} e_j)$. Hence ϕ is a ring isomorphism and so we have represented R as a generalized matrix ring. Our aim is to show that it is a semidirect sum.

LEMMA 1. If $e_i \neq e_j$ and $e_j \neq e_k$ then $e_iRe_jRe_k = 0$.

Proof. Let $x \in e_iRe_j$ and $y \in e_jRe_k$. Then $x^2 = 0$ (since $e_i e_j = 0$) so $1 + x$ is a unit. Similarly $1 + y$ is a unit and so, since R^* is abelian, $xy = yx$. But $x = e_i x$ and $y = e_j y$ so that $xy = e_i xy = e_i yx = e_i e_j yx = 0$.

It now follows easily that the multiplication in $[e_iRe_j]$ is that of

the semidirect sum. Indeed if $(x_{ij}), (y_{ij}) \in [e_i Re_j]$ and we write $(x_{ij})(y_{ij}) = (z_{ij})$, then using the lemma:

$$z_{ii} = \sum_k x_{ik} y_{ki} = x_{ii} y_{ii} \quad \text{for all } i = 1, 2, \dots, n,$$

$$z_{ij} = \sum_k x_{ik} y_{kj} = x_{ii} y_{ij} + x_{ij} y_{jj} \quad \text{for all } i \neq j.$$

Hence, in the notation of § 1, $R \cong [e_i Re_i, e_i Re_j]$. But then Proposition 1 shows that each $(e_i Re_i)^*$ is abelian and each $e_i Re_j$ is a G -unital bimodule. Since each $e_i Re_i$ is local, it is commutative by Proposition 2. Furthermore, e_i is central if and only if $e_i Re_j = 0 = e_j Re_i$ for all $j \neq i$. It follows that either e_i is central or $e_i Re_i$ possesses a nonzero G -unital module. In the latter case $e_i Re_i / J(e_i Re_i) \cong \mathcal{R}_2$ by Proposition 2. This proves the "only if" part of the following theorem; the rest follows from Propositions 1 and 2.

THEOREM 1. *Let R be a semiperfect ring. The group of units of R is abelian if and only if $R \cong T \oplus S$ where T is zero or a direct sum of commutative local rings R_i and S is zero or $S \cong [L_i, X_{ij}]$. Here each L_i is a commutative local ring with $L_i / J(L_i) \cong \mathcal{R}_2$ and each X_{ij} is a G -unital $L_i - L_j$ bimodule. Moreover:*

(1) *The bimodules X_{ij} can be chosen to be arbitrary elementary abelian 2-group where the action of L_i is defined as follows: If $r \in L_i$ and $x \in X_{ij}$ set $rx = x$ if $r \in L_i^*$ and $rx = 0$ if $r \in J(L_i)$.*

(2) *The group of units of R is isomorphic to the direct product of all the groups R_i^* , all the groups L_i^* and all the (additive) groups X_{ij} .*

This characterizes R completely up to the structure of the commutative local rings involved. The groups of units of these local rings inherit many properties from R^* by (2) and often this leads to a complete characterization. This will be exemplified in § 3 below in the case where R^* is assumed to be cyclic. Also, each of these local rings is a homomorphic image of R so they inherit many ring-theoretic conditions which could be imposed on R , for example the descending chain condition.

An immediate consequence of Theorem 1 is that if R is semiperfect and R^* is abelian then $R/J(R)$ is a finite direct sum of fields. Of course this result follows from structure theory.

The next result is a generalization (in the case where R has an identity) of a theorem of Eldridge and Fischer ([3], Th. 1, p. 244).

COROLLARY 1. *Let R be a semiperfect ring with abelian group of units. Then R is commutative if either of the following conditions is satisfied:*

- (1) $2x = 0$ in R implies $x = 0$
- (2) R^* has no direct factor each element of which has order 2.

Proof. Let $[L_i, X_{ij}]$ be the semidirect sum appearing in the decomposition of R . If $x \in X_{ij}$ then $2x = 0$ since X_{ij} is G -unital so each $X_{ij} = 0$ if condition (1) holds. If condition (2) holds each $X_{ij} = 0$ by (2) of the theorem. The result follows.

COROLLARY 2. *Let R be a semiperfect ring. If the group of units of R is abelian and finite then R is finite.*

Proof. By (2) of Theorem 1 each of the local rings appearing in the decomposition of R has a finite group of units and each of the bimodules appearing in the semidirect sum is finite. But if L is a local ring and L^* is finite then $J(L)$ is finite since $1 + J(L) \subseteq L^*$ and $L/J(L)$ is finite since $[L/J(L)]^* \cong L^*/(1 + J(L))$. This implies L is finite and the result follows.

A natural question is whether a semiperfect ring R must be finite when R^* is assumed to be abelian and finitely generated. The answer is yes if R^* is cyclic (see Theorem 2 below) or if $J(R)$ is nil. The next result will be useful in both cases.

LEMMA 2. *Let R be a commutative local ring. If R^* is finitely generated then R is noetherian, $R/J(R)$ is a finite field and $J(R)^n/J(R)^{n+1}$ is a finite ring for each n .*

Proof. If $A_1 \subseteq A_2 \subseteq \dots$ is a chain of (proper) ideals of R we have the chain $1 + A_1 \subseteq 1 + A_2 \subseteq \dots$ of subgroups of R^* . It follows that R is noetherian. We have $[R/J(R)]^* \cong R^*/1 + J(R)$ so the field $R/J(R)$ has a finitely generated group of units. Hence it is finite. Finally, $J(R)^n/J(R)^{n+1}$ is a vector space over $R/J(R)$ and is finite dimensional since R is noetherian.

We can now prove the following result which generalizes another result of Eldridge and Fischer ([3], Th. 2, p. 245).

PROPOSITION 3. *Let R be a semiperfect ring with R^* abelian. If R^* is finitely generated and $J(R)$ is nil then R is finite.*

Proof. Decompose R as in Theorem 1 and let L be one of the commutative local rings which appear. Then L^* is finitely generated by (2) of Theorem 1 and $J(L)$ is nil ($L = eRe$ for some $e^2 = e \in R$). Since L is noetherian by Lemma 2, write $J(L) = La_1 + La_2 + \dots + La_n$ where the $a_i \in J(L)$. Hence, if $m \geq 1$ we have $J(L)^m = \sum Lx_1^{k_1} x_2^{k_2} \dots x_n^{k_n}$

where the sum is taken over all $k_i \geq 0$ satisfying $k_1 + k_2 + \cdots + k_n = m$. Since $J(L)$ is nil this implies that $J(L)$ is nilpotent. But then Lemma 2 implies that $J(L)$, and hence L , is finite.

It remains to show that if $[L_i, X_{ij}]$ is the semidirect sum appearing in the decomposition of R then each X_{ij} is finite. But each X_{ij} is finitely generated as an additive group and so, since it is a vector space over \mathcal{K}_2 , it is finite. This completes the proof.

We remark that the hypothesis that $J(R)$ is nil was used only to show that $J(L)$ is nilpotent.

3. Cyclic groups of units. Gilmer [4] has characterized all finite commutative rings with a cyclic group of units and Eldridge and Fischer [3] have extended these results to artinian rings. In order to cover the semiperfect case we need the following negative result.

PROPOSITION 4. *If R is a commutative local ring the group of units of R is not infinite cyclic.*

Proof. Assume, on the contrary, that R^* is infinite cyclic. Then the characteristic of R is two since $(-1)^2 = 1$. By Lemma 2 R is noetherian and $R/J(R)$ is finite. Hence, $J(R)^2 \neq J(R)$. But the additive group $J(R)/J(R)^2$ is cyclic since it is naturally isomorphic to the multiplicative group $1 + J(R)/1 + J(R)^2$. Since the characteristic is two, it follows that $J(R)/J(R)^2$ has two elements. But $J(R)/J(R)^2$ is a vector space over the field $R/J(R)$ so $R/J(R) \cong \mathcal{K}_2$.

We now claim that R is an integral domain. If not let P be any prime ideal of R . Then $P \neq 0$ so $(R/P)^* \cong R^*/1 + P$ is finite cyclic. Since R/P is local it follows that R/P is a finite integral domain and hence that P is maximal. Hence every prime ideal is maximal and so ([6], p. 203) R is artinian. But then R is finite by Proposition 3, a contradiction. Hence R is an integral domain.

Now let u be a generator of $R^* = 1 + J(R)$. Write $u = 1 + a$ and $u^{-1} = 1 + b$ where $a, b \in J(R)$. Then $1 + a + a^2$ is a unit so $1 + a + a^2 = (1 + a)^k$ for some $k \in \mathcal{K}$. It is easy to check that $k = 0, 1, 2$ are impossible. Suppose $k \geq 3$. Then we have

$$1 + a + a^2 = 1 + \gamma a + \delta a^2 + a^t u$$

where $3 \leq t \leq k$, u is a unit and γ and δ are each either 0 or 1. Since R is a domain it is easy to check that each of these possibilities for γ and δ lead to a contradiction. Hence we must have $1 + a + a^2 = u^{-l}$ for some $l \geq 1$. But $1 + a + a^2 = 1 + u + u^2$ and $1 + b + b^2 = 1 + u^{-1} + u^{-2}$ as is easily verified. Hence

$$1 + b + b^2 = u^{-2}(1 + a + a^2) = u^{-2}u^{-l} = (1 + b)^{l+2}.$$

This leads to a contradiction just as before and so completes the proof.

We can now obtain a generalization of another result of Eldridge and Fischer ([3], Th. 3, p. 248) and, in so doing, obtain a much easier proof of that result.

THEOREM 2. *Let R be a semiperfect ring with cyclic group of units. Then R is finite and is either commutative or is isomorphic to the direct sum of a commutative ring and the ring of 2×2 upper triangular matrices over \mathcal{K}_2 .*

Proof. Decompose R as in Theorem 1. By (2) of Theorem 1, each of the local rings appearing has a cyclic group of units and, by Proposition 4, it is finite. It follows that each of these local rings is finite. Now let $[L_i, X_{ij}]$ be the semidirect sum appearing. Each X_{ij} is cyclic as an additive group and so, since it is an elementary abelian 2-group, it has two elements. Hence R is finite. Furthermore, the fact that all X_{ij} are direct factors of R^* means that at most one is nonzero. If all are zero then $[L_i, X_{ij}]$ is commutative so R is commutative. If, without loss of generality, $X_{12} \neq 0$, we have

$$[L_i, X_{ij}] = \begin{pmatrix} L_1 X_{12} \\ 0 & L_2 \end{pmatrix} \oplus L_3 \oplus \cdots \oplus L_n$$

where $\begin{pmatrix} L_1 X_{12} \\ 0 & L_2 \end{pmatrix}$ is a semidirect sum.

Moreover, each L_i^* has odd order so L_i has characteristic two. But then, if $a \in J(L_i)$, there exists an odd integer n such that $1 = (1 + a)^n = 1 + a + a^2 r$ where $r \in L_i$. Hence $a(1 + ar) = 0$ so $a = 0$. This means $J(L_i) = 0$ so $L_i \cong \mathcal{K}_2$. In particular, $\begin{pmatrix} L_1 X_{12} \\ 0 & L_2 \end{pmatrix}$ is isomorphic to the ring of 2×2 upper triangular matrices over \mathcal{K}_2 . This completes the proof.

This theorem completely characterizes the semiperfect rings with a cyclic group of units since the finite commutative local rings of this type have been characterized by Gilmer [4] and later by Ayoub [1] and Pearson and Schneider [5]. Gilmer cited the ring of 2×2 upper triangular matrices over \mathcal{K}_2 as an example of a finite noncommutative ring with cyclic group of units. Theorem 2 shows that this is essentially the only such semiperfect ring.

REFERENCES

1. C. W. Ayoub, *On finite primary rings and their groups of units*, *Compositio Math.*, **21** (1969), 247-252.
2. E. A. Behrens, *Ring Theory*, Academic Press, New York, 1972.
3. K. E. Eldridge and I. Fischer, *D. C. C. rings with a cyclic group of units*, *Duke Math. J.*, **34** (1967), 243-248.
4. R. W. Gilmer, *Finite rings with a cyclic multiplicative group of units*, *Amer. J. Math.*, **85** (1963), 447-452.
5. K. R. Pearson and J. E. Schneider, *Rings with a cyclic group of units*, *J. of Algebra*, **16** (1970), 243-251.
6. D. Zariski and P. Samuel, *Commutative Algebra*, Vol. I, Van Nostrand, Princeton, N. J., 1958.

Received August 4, 1972.

THE UNIVERSITY OF CALGARY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

A. Bigard, <i>Free lattice-ordered modules</i>	1
Richard Bolstein and Warren R. Wogen, <i>Subnormal operators in strictly cyclic operator algebras</i>	7
Herbert Busemann and Donald E. Glassco, II, <i>Irreducible sums of simple multivectors</i>	13
W. Wistar (William) Comfort and Victor Harold Saks, <i>Countably compact groups and finest totally bounded topologies</i>	33
Mary Rodriguez Embry, <i>Maximal invariant subspaces of strictly cyclic operator algebras</i>	45
Ralph S. Freese and James Bryant Nation, <i>Congruence lattices of semilattices</i>	51
Ervin Fried and George Grätzer, <i>A nonassociative extension of the class of distributive lattices</i>	59
John R. Giles and Donald Otto Koehler, <i>On numerical ranges of elements of locally m-convex algebras</i>	79
David A. Hill, <i>On dominant and codominant dimension of $\mathbf{QF} - 3$ rings</i>	93
John Sollion Hsia and Robert Paul Johnson, <i>Round and Pfister forms over $R(t)$</i>	101
I. Martin (Irving) Isaacs, <i>Equally partitioned groups</i>	109
Athanassios G. Kartsatos and Edward Barry Saff, <i>Hyperpolynomial approximation of solutions of nonlinear integro-differential equations</i>	117
Shin'ichi Kinoshita, <i>On elementary ideals of θ-curves in the 3-sphere and 2-links in the 4-sphere</i>	127
Ronald Brian Kirk, <i>Convergence of Baire measures</i>	135
R. J. Knill, <i>The Seifert and Van Kampen theorem via regular covering spaces</i>	149
Amos A. Kovacs, <i>Homomorphisms of matrix rings into matrix rings</i>	161
Young K. Kwon, <i>HD-minimal but no HD-minimal</i>	171
Makoto Maejima, <i>On the renewal function when some of the mean renewal lifetimes are infinite</i>	177
Juan José Martínez, <i>Cohomological dimension of discrete modules over profinite groups</i>	185
W. K. Nicholson, <i>Semiperfect rings with abelian group of units</i>	191
Louis Jackson Ratliff, Jr., <i>Three theorems on imbedded prime divisors of principal ideals</i>	199
Billy E. Rhoades and Albert Wilansky, <i>Some commutants in $B(c)$ which are almost matrices</i>	211
John Philip Riley Jr., <i>Cross-sections of decompositions</i>	219
Keith Duncan Stroyan, <i>A characterization of the Mackey uniformity $m(L^\infty, L^1)$ for finite measures</i>	223
Edward G. Thurber, <i>The Scholz-Brauer problem on addition chains</i>	229
Joze Vrabec, <i>Submanifolds of acyclic 3-manifolds</i>	243
Philip William Walker, <i>Adjoint boundary value problems for compactified singular differential operators</i>	265
Roger P. Ware, <i>When are Witt rings group rings</i>	279
James D. Wine, <i>Paracompactifications using filter bases</i>	285