

Pacific Journal of Mathematics

**DEFINITE AND SEMIDEFINITE MATRICES IN A REAL
SYMMETRIC MATRIX PENCIL**

FRANK UHLIG

DEFINITE AND SEMIDEFINITE MATRICES IN A REAL SYMMETRIC MATRIX PENCIL

FRANK UHLIG

Pencils that contain a definite matrix (*d*-pencils) have been characterized in several ways. Here *d*-pencils will be characterized by the property of the set $L = \{(a_i, b_i)\} \subseteq R^2$ if S and T are simultaneously congruent to $\text{diag}(a_i)$ and $\text{diag}(b_i)$, respectively. This way one can describe all definite and semidefinite matrices in a *d*-pencil. Similarly one can characterize all pencils that contain semidefinite but no definite matrices (*s.d.* pencils). The explicit condition on L for *d*-pencils is then used to reprove the theorem that two real symmetric matrices generate a *d*-pencil iff their associated quadratic forms do not vanish simultaneously.

DEFINITION 1. If S is symmetric we define $Q_S = \{x \in R^n \mid x'Sx = 0\}$.

DEFINITION 2. For real symmetric (r.s.) matrices S and T one defines the pencil $P(S, T) = \{aS + bT \mid a, b \in R\}$.

DEFINITION 3. (a) $P(S, T)$ is called *d*-pencil if $P(S, T)$ contains a definite matrix.

(b) $P(S, T)$ is called *s.d. pencil* if $P(S, T)$ contains a nonzero semidefinite, but no definite matrix.

The following theorem is classical:

THEOREM 0.1. If $P(S, T)$ is a *d*-pencil then S and T can be diagonalized simultaneously by a real congruence transformation.

The question whether a given pencil of r.s. matrices contains a positive definite matrix was treated in chronological order by Finsler [6], Albert [1], Reid [12], Hestenes and McShane [9], Dines [5], Calabi [4], Taussky [13], Hestenes [8], Theorem 3, and Berman [3].

Their main results are the following:

THEOREM 0.2. Let S and T be r.s. $n \times n$ matrices. If $n \geq 3$, then the following are equivalent:

- (i) $P(S, T)$ is a *d*-pencil,
- (ii) $Q_S \cap Q_T = \{0\}$, and
- (iii) $\text{trace } YS = \text{trace } YT = 0$ for Y positive semidefinite implies $Y = 0$.

The equivalence of (i) and (ii) was proved by Calabi [4], while Berman [3] showed the equivalence of (i) and (iii).

Before Calabi [4] only condition

$$(ii') \quad x'Sx = 0 \quad \text{implies} \quad x'Tx > 0$$

was generally used instead of (ii). And thus only the fact that (ii') implies (i) was proved.

In view of Theorem 0.1, to characterize d -pencils means to characterize the sets $L = \{(a_i, b_i) \mid i = 1, \dots, n\} \subseteq \mathbf{R}^2$ for which there exist $\lambda, \mu \in \mathbf{R}$ with $\lambda a_i + \mu b_i > 0$ for all i .

THEOREM 1.1. *$P(S, T)$ is a d -pencil iff S and T can be simultaneously diagonalized by a real congruence transformation X : $X'SX = \text{diag}(a_i)$, $X'TX = \text{diag}(b_i)$, and in case that there are indices i, j with $a_i a_j < 0$ we have*

$$(a) \quad \max_{a_i > 0} \frac{b_i}{a_i} < \max_{a_i < 0} \frac{b_i}{a_i} \quad \text{and} \quad b_i < 0 \quad \text{whenever} \quad a_i = 0,$$

or

$$(b) \quad \min_{a_i > 0} \frac{b_i}{a_i} > \min_{a_i < 0} \frac{b_i}{a_i} \quad \text{and} \quad b_i > 0 \quad \text{whenever} \quad a_i = 0,$$

while in case that all a_i have one sign we have

$$(c) \quad \text{either } b_i < 0 \text{ whenever } a_i = 0 \text{ or } b_i > 0 \text{ whenever } a_i = 0.$$

Proof. If $P(S, T)$ is a d -pencil, then by Theorem 0.1 the matrices S and T can be diagonalized simultaneously by a real congruence transformation: $X'SX = \text{diag}(a_i)$, $X'TX = \text{diag}(b_i)$. Furthermore, there exist $\lambda, \mu \in \mathbf{R}$ s.t. $\lambda S + \mu T = X' \text{diag}(\lambda a_i + \mu b_i) X$ is positive definite.

Hence the set $L = \{(a_i, b_i) \mid i = 1, \dots, n\} \subseteq \mathbf{R}^2$ must lie in an open half plane of \mathbf{R}^2 and it follows that either (a) or (b) or (c) must hold for L .

Conversely if S and T are simultaneously congruent to $\text{diag}(a_i)$ and $\text{diag}(b_i)$ then in case of (a) all points $(a_i, b_i) \in \mathbf{R}^2$ lie "below" any line L_α thru zero that has slope $\max_{a_i > 0} b_i/a_i < \alpha < \max_{a_i < 0} b_i/a_i$. In case of (b), they all lie "above" any such line L_α with $\min_{a_i > 0} b_i/a_i > \alpha > \min_{a_i < 0} b_i/a_i$ and in case of (c) they all lie to the "right" or to the "left" of any line L_α where either

$$\max_{a_i > 0} \frac{b_i}{a_i} < \alpha < \infty \quad \text{or} \quad -\infty < \alpha < \min_{a_i > 0} \frac{b_i}{a_i}.$$

Thus the set L lies in an open half plane and hence there exist $\lambda, \mu \in \mathbf{R}$ s.t. $\lambda a_i + \mu b_i > 0$ for all i . Thus $P(S, T)$ is a d -pencil.

Next, if $P(S, T)$ is a d -pencil we will explicitly describe the set $M^+ = \{(\lambda, \mu) \mid \lambda S + \mu T \text{ pos. def.}\} \subseteq R^2$ and determine the positive semidefinite matrices in $P(S, T)$. The sets M^+ and its closure $M = \{(\lambda, \mu) \mid \lambda S + \mu T \text{ pos. semidef.}\}$ have been previously characterized as convex cones and this fact together with properties of quadratic forms have been used to prove the equivalence of (i) and (ii) in Theorem 0.2 (Hestenes [8]). Hestenes [8] moreover treats related questions for positively elliptic pairs of quadratic forms in Hilbert space.

THEOREM 1.2. *Let $P(S, T)$ be a d -pencil and let S, T be simultaneously congruent to $\text{diag}(a_i)$ and $\text{diag}(b_i)$.*

Then with (a), (b), and (c) from Theorem 1.1 we have

(1) $\lambda S + \mu T$ is positive definite iff

in case of (a) $-\left(\max_{a_i > 0} \frac{b_i}{a_i}\right)^{-1} < \frac{\lambda}{\mu} < -\left(\max_{a_i < 0} \frac{b_i}{a_i}\right)^{-1}$ and

in case of (b) $-\left(\min_{a_i > 0} \frac{b_i}{a_i}\right)^{-1} > \frac{\lambda}{\mu} > -\left(\min_{a_i < 0} \frac{b_i}{a_i}\right)^{-1}$ while

in case of (c) if all a_i are nonnegative

$$-\left(\max_{a_i > 0} \frac{b_i}{a_i}\right)^{-1} < \frac{\lambda}{\mu} < 0 \quad \text{if } b_i < 0 \text{ whenever } a_i = 0$$

and

$$0 < \frac{\lambda}{\mu} < -\left(\min_{a_i > 0} \frac{b_i}{a_i}\right)^{-1} \quad \text{if } b_i > 0 \text{ whenever } a_i = 0$$

and, if all $a_i \leq 0$, we have

$$-\left(\max_{a_i < 0} \frac{b_i}{a_i}\right)^{-1} > \frac{\lambda}{\mu} > 0 \quad \text{if } b_i < 0 \text{ whenever } a_i = 0$$

while

$$0 > \frac{\lambda}{\mu} > -\left(\min_{a_i < 0} \frac{b_i}{a_i}\right)^{-1} \quad \text{if } b_i > 0 \text{ whenever } a_i = 0.$$

(2) $\lambda S + \mu T$ is positive semidefinite and not definite iff in each case respectively, λ/μ is equal to either one of the endpoints of the intervals under (1).

Here we set

$$(0/a)^{-1} = \begin{cases} \infty & \text{if } a > 0 \\ -\infty & \text{if } a < 0. \end{cases}$$

Proof. The proof follows from Theorem 1.1 and by the geometry

of \mathbf{R}^2 . Concerning (2) one can say that if λ/μ is an endpoint of an appropriate interval in (1), then $\lambda S + \mu T$ is positive semidefinite of rank $n - r \leq n$ iff $r = |\{j \mid b_j/\alpha_j = \lambda/\mu\}|$.

As corollaries we observe that

COROLLARY 1. *If in Theorem 1.1 the strict reverse inequalities hold in (a) and (b), or (c), respectively, then $\mu S + \lambda T$ is indefinite for any $(\lambda, \mu) \neq (0, 0)$. And then $\mu S + \lambda T$ has at least rank 2 for any $(\lambda, \mu) \neq (0, 0)$.*

Before characterizing *s.d.* pencil we will quote a simplified version of the canonical pair form theorem for nonsingular pairs of r.s. matrices.

DEFINITION 4. A pair of r.s. matrices S, T is called *nonsingular*, if S is nonsingular.

THEOREM 1.3. *Let S and T be a nonsingular pair of r.s. matrices. Let $S^{-1}T$ have real Jordan normal form $\text{diag}(J_1, \dots, J_m)$. Then S and T are simultaneously congruent by a real congruence transformation to $\text{diag}(\varepsilon_i E_i)$ and $\text{diag}(\varepsilon_i E_i J_i)$ respectively, where $\varepsilon_i = \pm 1$ and E_i denotes the square matrix $\begin{pmatrix} 0 & & & 1 \\ & \cdot & \cdot & \\ & & \cdot & \\ 1 & & & 0 \end{pmatrix}$ of the same size as J_i for $i = 1, \dots, m$.*

The relevant notation has recently been described in Uhlig [15]. There it was shown that the canonical pair form just described is a finest simultaneous block diagonalization.

Note that if $J = \begin{pmatrix} \lambda & 1 & & 0 \\ & \cdot & \cdot & \\ & & \cdot & \\ 0 & & & \lambda & 1 \end{pmatrix}$ is a Jordan block for $\lambda \in \mathbf{R}$ then for $E = \begin{pmatrix} 0 & & & 1 \\ & \cdot & \cdot & \\ & & \cdot & \\ 1 & & & 0 \end{pmatrix}$ with $\dim E = \dim J \geq 3$ we have

$$aE + bEJ = \begin{pmatrix} 0 & & & a + b\lambda \\ & \cdot & \cdot & b \\ & & \cdot & \\ a + b\lambda & b & & 0 \end{pmatrix}.$$

Hence $aE + bEJ$ is indefinite for all $(a, b) \neq (0, 0)$. The same holds for any J corresponding to an eigenvalue $\lambda = \alpha + \beta i \notin \mathbf{R}$. With this in mind we can characterize *s.d.* pencils:

THEOREM 1.4. *Let S, T be a nonsingular pair of r.s. matrices of dimension greater than 2.*

Then $P(S, T)$ is a *s.d.* pencil iff either S and T are simultaneously congruent to $\text{diag}(a_i)$ and $\text{diag}(b_i)$ with $b_i/a_i \neq b_j/a_j$ for at least one pair of indices (i, j) and we have

$$(a') \max_{a_i > 0} \frac{b_i}{a_i} = \max_{a_i < 0} \frac{b_i}{a_i} \quad \text{or} \quad (b') \min_{a_i > 0} \frac{b_i}{a_i} = \min_{a_i < 0} \frac{b_i}{a_i},$$

or S and T are simultaneously congruent to

$$\text{diag}(\varepsilon E, \dots, \varepsilon E, \varepsilon_{k+1}, \dots, \varepsilon_j, a_{j+1}, \dots, a_n)$$

and

$$\text{diag}(\varepsilon EJ, \dots, \varepsilon EJ, \varepsilon_{k+1}\alpha, \dots, \varepsilon_j\alpha, b_{j+1}, \dots, b_n),$$

where $J = \begin{pmatrix} \alpha & 1 \\ 0 & \alpha \end{pmatrix}$, $E = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$; $\varepsilon, \varepsilon_i = \pm 1$, $\alpha \in \mathbf{R}$, $b_l \neq \alpha a_l$ and $\varepsilon, b_l - \alpha a_l$ all have the same sign for $l = j + 1, \dots, n$.

Proof. We recall that $P(S, T)$ is a *s.d.* pencil if there is a semi-definite but no definite matrix in $P(S, T)$.

If S and T can be diagonalized simultaneously to yield $\text{diag}(a_i)$ and $\text{diag}(b_i)$, then $P(S, T)$ is a *s.d.* pencil iff $L = \{(a_i, b_i)\} \subseteq \mathbf{R}^2$ lies in a closed half plane, but in no open half plane of \mathbf{R}^2 , nor on a line thru 0; which is the case iff (a') or (b') holds. (Note that since S is nonsingular we have $a_i \neq 0$ for all i .)

If $P(S, T)$ is a *s.d.* pencil and S and T cannot be simultaneously diagonalized, then the real Jordan normal form J_0 of $S^{-1}T$ cannot contain a Jordan block of dimension greater than two for a real eigenvalue, nor can $S^{-1}T$ have complex roots as we have observed in the sequel of Theorem 1.3. But for $J = \begin{pmatrix} \alpha & 1 \\ 0 & \alpha \end{pmatrix}$ and $E = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ we have $\lambda E + \mu EJ = \begin{pmatrix} 0 & \lambda + \mu\alpha \\ \lambda + \mu\alpha & \mu \end{pmatrix}$ is semidefinite iff $\lambda + \mu\alpha = 0$. Hence if there are several 2-dimensional blocks in J_0 they must all correspond to the same eigenvalue $\alpha \in \mathbf{R}$ and they all must carry the same sign ε . The signs of the one-dimensional blocks corresponding to the same eigenvalue α cannot be specified since $\varepsilon_i(\lambda + \mu\alpha) = 0$ independent of $\varepsilon_i = \pm 1$, but for one-dimensional blocks in J_0 not corresponding to α , i.e., if $b_l \neq \alpha a_l$, then we must have that $\varepsilon\mu$ and $\lambda a_l + \mu b_l = \mu(b_l - \alpha a_l)$ all have the same sign for $l = j + 1, \dots, n$. If $\mu = 0$ we conclude that if $\lambda E + \mu EJ = \lambda E$ is semidefinite then $\lambda = 0$. Thus μ cannot be zero. And the theorem is proved, since the converse is obvious in this case.

Finally we will apply Theorem 1.1 to give a new elementary proof of the equivalence of (i) and (ii) in Theorem 0.2.

For this we need a theorem of Greub and Milnor [7, p. 256]:

THEOREM 2.1. *Let S, T be r.s. matrices of dimension greater than 2. If $Q_S \cap Q_T = \{0\}$, then S and T can be diagonalized simultaneously by a real congruence transformation.*

This Theorem was also proved by Majindar [11], Kraljevic [10], Wonenburger [16], Au-Yeung [1], and the author [14].

Proof. [(i) and (ii) of Theorem 0.2 are equivalent.] That (i) implies (ii) is obvious since for definite matrices U we have $Q_U = \{0\}$ and clearly for any $U \in P(S, T)$ we have $Q_U \supseteq Q_S \cap Q_T$.

If (ii) holds, then by Theorem 2.1 the matrices S and T are simultaneously congruent to $D_1 = \text{diag}(a_i)$ and $D_2 = \text{diag}(b_i)$. We are going to show that

$$\sum a_i x_i^2 = \sum b_i x_i^2 = 0 \quad \text{implies } x_i = 0 \quad \text{for all } i$$

cannot hold unless either (a) or (b) or (c) of Theorem 1.1 is the case.

In the case that, say, all $a_i \geq 0$ and (c) is violated, then we have $a_i = a_j = 0$ for $i \neq j$, while $b_i \geq 0$ and $b_j \leq 0$. With e_i denoting the i th unit vector we then have

$$0 \neq x = e_i + \sqrt{\frac{b_i}{-b_j}} \cdot e_j \in Q_{D_1} \cap Q_{D_2} \quad \text{if } b_j \neq 0,$$

while else $e_j \in Q_{D_1} \cap Q_{D_2}$, hence $Q_S \cap Q_T \neq \{0\}$ as well.

In the case that there are positive as well as negative a_i and if both

$$\left[\max_{a_i > 0} \frac{b_i}{a_i} \geq \max_{a_i < 0} \frac{b_i}{a_i} \text{ or } b_i \geq 0 \text{ for some } a_i = 0 \right]$$

and

$$\left[\min_{a_i > 0} \frac{b_i}{a_i} \leq \min_{a_i < 0} \frac{b_i}{a_i} \text{ or } b_i < 0 \text{ for some } a_i = 0 \right]$$

hold, i.e., if both (a) and (b) of Theorem 1.1 are violated, then we will show that $Q = Q_{D_1} \cap Q_{D_2} \neq \{0\}$ and then the equivalence of (i) and (ii) will be proved.

We now go into subcases:

If both $b_i \geq 0$ and $b_j \leq 0$ for $a_i = a_j = 0$, then $Q_{D_1} \cap Q_{D_2} \neq \{0\}$ as we have just seen.

If both $\max_{a_i > 0} b_i/a_i \geq \max_{a_i < 0} b_i/a_i$ and $b_i \leq 0$ for some $a_i = 0$ hold, then we assume without loss of generality that the indices in question are 1, 2, 3, respectively, i.e., $b_1/a_1 \geq b_2/a_2$ where $a_1 > 0$, $a_2 < 0$, and $a_3 = 0$, while $b_3 \leq 0$.

If $b_3 = 0$, then $e_3 \in Q$. If $b_1/a_1 = b_2/a_2$, then $e_1 + (\sqrt{-a_1/a_2}) e_2 \in Q$. Otherwise with $x = (1, \sqrt{-a_1/a_2}, \alpha, 0, \dots, 0)$ we get

$$\sum a_i x_i^2 = a_1 - a_1 \frac{a_2}{a_2} = 0 \text{ independent of } \alpha,$$

and

$$\sum b_i x_i^2 = b_1 - \frac{b_2 a_1}{a_2} + b_3 \alpha^2.$$

Now by assumption $b_1 - (a_1/a_2) b_2 > 0$ and $b_3 < 0$, hence there is a real α s.t. $\sum b_i x_i^2 = 0$ as well, and consequently $Q \neq \{0\}$. If both $\min_{a_i > 0} b_i/a_i \leq \min_{a_i < 0} b_i/a_i$ and $b_i \geq 0$ for some $a_i = 0$ hold, then a similar argument applies.

Finally assume both

$$(*) \quad \max_{a_i > 0} \frac{b_i}{a_i} \geq \max_{a_i < 0} \frac{b_i}{a_i} \quad \text{and} \quad \min_{a_i > 0} \frac{b_i}{a_i} \leq \min_{a_i < 0} \frac{b_i}{a_i} \text{ hold.}$$

If in (*) there is an equal sign, say $b_1/a_1 = b_2/a_2$ with $a_1 > 0, a_2 < 0$, then as above $e_1 + (\sqrt{-a_1/a_2}) e_2 \in Q$. Thus the only case to remain to be proved is if both inequalities in (*) are strict. For $n \geq 4$ assume that the indices in question are 1, 2, 3, 4; i.e., $b_1/a_1 > b_2/a_2$ with $a_1 > 0, a_2 < 0$ and $b_3/a_3 < b_4/a_4$ with $a_3 > 0, a_4 < 0$. Then with $x = (1, (\sqrt{-a_1/a_2}), \alpha, (\sqrt{-a_3/a_4}) \alpha, 0, \dots, 0)$ we have

$$\sum a_i x_i^2 = a_1 - a_1 + \alpha^2(a_3 - a_3) = 0 \text{ independently of } \alpha,$$

and

$$\sum b_i x_i^2 = b_1 - \frac{a_1}{a_2} b_2 + \alpha^2 \left(b_3 - b_4 \frac{a_3}{a_4} \right).$$

By assumption $b_1 - (a_1/a_2) b_2 > 0$ and $b_3 - b_4 (a_3/a_4) < 0$. Thus $\alpha \in \mathbf{R}$ can be chosen s.t. $\sum b_i x_i^2 = 0$.

For $n = 3$ we may WLOG assume that $\min_{a_i > 0} b_i/a_i = \max_{a_i > 0} b_i/a_i = b_1/a_1$. Then (*) reads like $b_1/a_1 > b_2/a_2$ and $b_1/a_1 < b_3/a_3$ with $a_1 > 0; a_2, a_3 < 0$. Then with $x = (1, \sqrt{-a_1 \alpha/a_2}, \sqrt{-a_1 \beta/a_3})$ we have

$$\sum a_i x_i^2 = a_1 - a_1(\alpha + \beta) \text{ and } \sum b_i x_i^2 = b_1 - \frac{b_2 a_1}{a_2} \alpha - \frac{b_3 a_1}{a_3} \beta.$$

Now the linear system in α and β :

$$\alpha + \beta = 1 \quad \frac{b_2 a_1}{a_2} \alpha + \frac{b_3 a_1}{a_3} \beta = b_1$$

can be solved for arbitrary b_1 , since its determinant

$$a_1 \left(\frac{b_3}{a_3} - \frac{b_2}{a_2} \right) \neq 0$$

by assumption.

This completes the proof.

REFERENCES

1. A. A. Albert, *A quadratic form problem in the calculus of variations*, Bull. Amer. Math. Soc., **44** (1938), 250-253.
2. Y.-H. Au-Yeung, *Some theorems on the real pencil and simultaneous diagonalization of two hermitian bilinear forms*, Proc. Amer. Math. Soc., **23** (1969), 246-253.
3. A. Berman, *Thesis*, Northwestern University, Evanston, Ill., 1970.
4. E. Calabi, *Linear systems of real quadratic forms*, Proc. Amer. Math. Soc., **15** (1964), 844-846.
5. L. Dines, *On the mapping of quadratic forms*, Bull. Amer. Math. Soc., **47** (1941), 494-498.
6. P. Finsler, *Über das Vorkommen definitiver und semidefiniter Formen in Scharen quadratischer Formen*, Commentarii Math. Helvetici, **9** (1937), 188-192.
7. W. Greub, *Linear Algebra*, Springer, Berlin, 3rd edition, 1967.
8. M. Hestenes, *Pairs of quadratic forms*, Linear Algebra and Appl. **1** (1968), 397-407.
9. M. Hestenes and McShane, *Quadratic forms and the calculus of variations*, Trans. Amer. Math. Soc., **38** (1935), 145.
10. H. Kraljevic, *Simultaneous diagonalization of two symmetric bilinear symmetric functionals*, Glasnik Matematički, **1** (1966), 57-63.
11. K. Majindar, *On simultaneous hermitian congruence transformations of matrices*, Amer. Math. Monthly, **70** (1963), 842-844.
12. W. Reid, *A theorem on quadratic forms*, Bull. Amer. Math. Soc., **44** (1938), 437.
13. O. Taussky, *Positive Definite Matrices*, in "Inequalities", O. Shisha, Ed., Academic Press, N.Y., 1967.
14. F. Uhlig, *A study of the canonical form of a pair of real symmetric matrices and applications to pencils and to pairs of quadratic forms*, Thesis, California Institute of Technology, 1971-72.
15. ———, *Simultaneous block diagonalization of two real symmetric matrices*, Linear Algebra and Appl., **7** (1973), 281-289.
16. M. Wonenburger, *Simultaneous diagonalization of symmetric bilinear forms*, J. of Math. and Mech., **15** (1966), 617-622.

Received July 12, 1972.

UNIVERSITÄT WÜRZBURG

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI*

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$48.00 a year (6 Vols., 12 issues). Special rate: \$24.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by
Pacific Journal of Mathematics
All Rights Reserved

Pacific Journal of Mathematics

Vol. 49, No. 2

June, 1973

Wm. R. Allaway, <i>On finding the distribution function for an orthogonal polynomial set</i>	305
Eric Amar, <i>Sur un théorème de Mooney relatif aux fonctions analytiques bornées</i>	311
Robert Morgan Brooks, <i>Analytic structure in the spectrum of a natural system</i>	315
Bahattin Cengiz, <i>On extremely regular function spaces</i>	335
Kwang-nan Chow and Moses Glasner, <i>Atoms on the Royden boundary</i>	339
Paul Frazier Duvall, Jr. and Jim Maxwell, <i>Tame Z^2-actions on E^n</i>	349
Allen Roy Freedman, <i>On the additivity theorem for n-dimensional asymptotic density</i>	357
John Griffin and Kelly Denis McKennon, <i>Multipliers and the group L_p-algebras</i>	365
Charles Lemuel Hagopian, <i>Characterizations of λ connected plane continua</i>	371
Jon Craig Helton, <i>Bounds for products of interval functions</i>	377
Ikuko Kayashima, <i>On relations between Nörlund and Riesz means</i>	391
Everett Lee Lady, <i>Slender rings and modules</i>	397
Shozo Matsuura, <i>On the Lu Qi-Keng conjecture and the Bergman representative domains</i>	407
Stephen H. McCleary, <i>The lattice-ordered group of automorphisms of an α-set</i>	417
Stephen H. McCleary, <i>o-2-transitive ordered permutation groups</i>	425
Stephen H. McCleary, <i>o-primitive ordered permutation groups. II</i>	431
Richard Rochberg, <i>Almost isometries of Banach spaces and moduli of planar domains</i>	445
R. F. Rossa, <i>Radical properties involving one-sided ideals</i>	467
Robert A. Rubin, <i>On exact localization</i>	473
S. Sribala, <i>On Σ-inverse semigroups</i>	483
H. M. (Hari Mohan) Srivastava, <i>On the Konhauser sets of biorthogonal polynomials suggested by the Laguerre polynomials</i>	489
Stuart A. Steinberg, <i>Rings of quotients of rings without nilpotent elements</i>	493
Daniel Mullane Sunday, <i>The self-equivalences of an H-space</i>	507
W. J. Thron and Richard Hawks Warren, <i>On the lattice of proximities of Čech compatible with a given closure space</i>	519
Frank Uhlig, <i>The number of vectors jointly annihilated by two real quadratic forms determines the inertia of matrices in the associated pencil</i>	537
Frank Uhlig, <i>On the maximal number of linearly independent real vectors annihilated simultaneously by two real quadratic forms</i>	543
Frank Uhlig, <i>Definite and semidefinite matrices in a real symmetric matrix pencil</i> ...	561
Arnold Lewis Villone, <i>Self-adjoint extensions of symmetric differential operators</i>	569
Cary Webb, <i>Tensor and direct products</i>	579
James Victor Whittaker, <i>On normal subgroups of differentiable homeomorphisms</i>	595
Jerome L. Paul, <i>Addendum to: "Sequences of homeomorphisms which converge to homeomorphisms"</i>	615
David E. Fields, <i>Correction to: "Dimension theory in power series rings"</i>	616
Peter Michael Curran, <i>Correction to: "Cohomology of finitely presented groups"</i>	617
Billy E. Rhoades, <i>Correction to: "Commutants of some Hausdorff matrices"</i>	617
Charles W. Trigg, <i>Corrections to: "Versum sequences in the binary system"</i>	619