

Pacific Journal of Mathematics

SELF-ADJOINT EXTENSIONS OF SYMMETRIC DIFFERENTIAL OPERATORS

ARNOLD LEWIS VILLONE

SELF-ADJOINT EXTENSIONS OF SYMMETRIC DIFFERENTIAL OPERATORS

ARNOLD VILLONE

Let \mathcal{H} denote the Hilbert space of square summable analytic function on the unit disk, and consider those formal differential operators

$$L = \sum_{i=0}^n p_i D^i$$

which give rise to symmetric operators in \mathcal{H} . This paper is devoted to a study of when these operators are actually self-adjoint or admit of self-adjoint extensions in \mathcal{H} . It is shown that in the first order case the operator is always self-adjoint. For $n > 1$ sufficient conditions on the p_i are obtained for the existence of self-adjoint extensions. In particular a condition on the coefficients is obtained which insures that the operator has defect indices equal to the order of L .

Let \mathcal{A} denote the space of functions analytic on the unit disk and \mathcal{H} the subspace of square summable functions in \mathcal{A} with inner product

$$(f, g) = \int_{|z| < 1} \int f(z) \overline{g(z)} dx dy .$$

A complete orthonormal set for \mathcal{H} is provided by the normalized powers of z ,

$$e_n(z) = [(n + 1)/\pi]^{1/2} z^n , \quad n = 0, 1, \dots .$$

From this it follows that \mathcal{H} is identical with the space of power series $\sum_{n=0}^{\infty} a_n z^n$ which satisfy

$$(1.1) \quad \sum_{n=0}^{\infty} |a_n|^2 / (n + 1) < \infty .$$

Consider the formal differential operator

$$L = p_n D^n + \dots + p_1 D + p_0 ,$$

where $D = d/dz$ and the p_i are in \mathcal{H} . We now associate two operators as follows. Let \mathcal{D}_0 denote the span of the e_n and \mathcal{D} the set of all f in \mathcal{H} for which Lf is in \mathcal{H} , and define T_0 and T as

$$\begin{aligned} T_0 f &= Lf & f \in \mathcal{D}_0 \\ T f &= Lf & f \in \mathcal{D} . \end{aligned}$$

It is shown in [2] that T_0 and T are both densely defined operators

in \mathcal{H} , $T_0 \subseteq T$ and T is closed. Moreover, T_0 is symmetric if and only if

$$(1.2) \quad (Le_n, e_m) = (e_n, Le_m), \quad n, m = 0, 1, \dots$$

Such a formal operator is said to be formally symmetric. Regarding symmetric T_0 we have the following result.

THEOREM 1.1. *If T_0 is symmetric, $T_0^* = T$ and $T^* \subseteq T$. The closure of T_0 , $S = T_0^{**} = T^*$, is self-adjoint if and only if $S = T$.*

Proof. See [2].

For f and g in \mathcal{D} consider the bilinear form

$$(1.3) \quad \langle f, g \rangle = (Lf, g) - (f, Lg),$$

and let $\tilde{\mathcal{D}}$ be the set of those f in \mathcal{D} for which $\langle f, g \rangle = 0$ for all g in \mathcal{D} . Since $S = T^*$ and $\mathcal{D}(T^*) = \tilde{\mathcal{D}}$, S has domain $\tilde{\mathcal{D}}$.

Let \mathcal{D}^+ and \mathcal{D}^- denote the set of all solutions of the equation $Lu = iu$ and $Lu = -iu$ respectively, which are in \mathcal{H} . It is known from the general theory of Hilbert space [1, p. 1227-1230] that $\tilde{\mathcal{D}} = \tilde{\mathcal{D}} + \mathcal{D}^+ + \mathcal{D}^-$, and every $f \in \tilde{\mathcal{D}}$ has a unique such representation. Let the dimensions of \mathcal{D}^+ and \mathcal{D}^- be m^+ and m^- respectively. Clearly, m^+ and m^- cannot exceed the order of L . These integers are referred to as the deficiency indices of S , and S has self-adjoint extensions if and only if $m^+ = m^-$. Moreover, S is self-adjoint if and only if $m^+ = m^- = 0$.

2. In [2] it is shown that the general formally symmetric first order operator is given by

$$(2.1) \quad L = (cz^2 + az + \bar{c})D + (2cz + b)$$

where a and b are real. In this case it is possible to compute the solutions of $Lu = \pm iu$ explicitly and show that the solutions so obtained are not in \mathcal{H} . Proceeding in this manner we obtain the following result.

THEOREM 2.1. *If L is a first order formally symmetric operator, the associated operator T is self-adjoint.*

Proof. We shall show that m^+ and m^- are both zero. When $c = 0$ L is just the first order Euler operator, and hence T is self-adjoint by the corollary to Theorem 1.3 of [2]. When $c \neq 0$ we have

$$(2.2) \quad (z^2 + (a/c)z + \bar{c}/c)u' + (2z + b/c - i/c)u = 0$$

$$(2.3) \quad (z^2 + (a/c)z + \bar{c}/c)u' + (2z + b/c + i/c) = 0 .$$

The coefficient of u' has zeros at

$$\begin{aligned} \alpha &= -a/2c + (a^2 - 4|c|^2)^{1/2}/2c . \\ \beta &= -a/2c - (a^2 - 4|c|^2)^{1/2}/2c . \end{aligned}$$

There are three cases to consider:

1. $a^2 < 4|c|^2$
2. $a^2 = 4|c|^2$
3. $a^2 > 4|c|^2$.

In case 1 we have $\alpha = -a/2c + iR/2c, \beta = -a/2c - iR/2c$ where $R = (4|c|^2 - a^2)^{1/2}$, moreover $|\alpha| = |\beta| = 1$. Every solution of (2.2) is a multiple of the fundamental solution $\phi(z) = (z - \alpha)^{-r}(z - \beta)^{-s}$ where $r = (R - 1)/R - i(b - a)/R$ and $s = (R + 1)/R + i(b - a)/R$. Hence every (nontrivial) solution of (2.2) is analytic in the open unit disc D with at least one singularity on the boundary at $z = \beta$. We now show that ϕ is not in \mathcal{H} , i.e., the integral $\int_D \int |\phi(z)|^2 dx dy$ diverges. Introduce polar coordinates at β so $z - \beta = \rho e^{i\theta}$. Let δ be less than $|\beta - \alpha|$, then there exist suitable θ_1 and θ_2 such that for $0 < \varepsilon < \delta$, the regions $W_\varepsilon = \{z | \varepsilon \leq \rho \leq \delta, \theta_1 \leq \theta \leq \theta_2\}$ lie within D and $\alpha \notin W_\varepsilon$. Now

$$(2.4) \quad \int_D \int |\phi(z)|^2 dx dy \geq \lim_{\varepsilon \rightarrow 0} \int_{W_\varepsilon} \int |(z - \alpha)^{-r}|^2 |(z - \beta)^{-s}|^2 dx dy .$$

Since $\alpha \notin W_\varepsilon$ it follows from continuity that $|(z - \alpha)^{-r}|^2 \geq m > 0$ for z in W_ε , all $0 < \varepsilon < \delta$. Using this and the fact that $|(z - \beta)^{-s}| = \rho^{-u} e^{v\theta}$, where $s = u + iv$, the inequality of (2.4) becomes

$$\begin{aligned} \int_D \int |\phi(z)|^2 dx dy &\geq \lim_{\varepsilon \rightarrow 0} m \int_{\theta_1}^{\theta_2} \int_{\varepsilon}^{\delta} \rho^{-2u+1} e^{2v\theta} d\rho d\theta \\ &\geq \lim_{\varepsilon \rightarrow 0} mk(\theta_2 - \theta_1) \int_{\varepsilon}^{\delta} \rho^{-2u+1} d\rho , \end{aligned}$$

where $k = \text{infimum of } e^{2v\theta} \text{ on } \theta_1 \leq \theta \leq \theta_2$ which is greater than zero. But $-2u + 1 = -2(R + 1)/R + 1 = -1 - 2/R < -1$, hence the integral on the left diverges and ϕ is not square summable.

The fundamental solution for (2.3) is given by $\phi(z) = (z - \alpha)^{-r}(z - \beta)^{-s}$, where $r = (R + 1)/R - i(b - a)/R$ and $s = (R - 1)/R + i(b - a)/R$. Hence $\phi(z)$ is analytic in the open unit disc D with a singularity on the boundary at α . Let $z - \alpha = \rho e^{i\theta}$, then there exist suitable θ_1 and θ_2 such that for $0 < \varepsilon < \delta < |\alpha - \beta|$, the regions $W_\varepsilon = \{z | \varepsilon \leq \rho \leq \delta, \theta_1 \leq \theta \leq \theta_2\}$ lie within D and $\beta \notin W_\varepsilon$. As before, we obtain

$$\int_D \int |\phi(z)|^2 dx dy \geq \lim_{\varepsilon \rightarrow 0} mk(\theta_2 - \theta_1) \int_{\varepsilon}^{\delta} \rho^{-2u+1} d\rho$$

where $|(z - \beta)^{-s}|^2 \geq m > 0$ for all z in W_ε and $0 < \varepsilon < \delta$, k is the infimum of $e^{2v\theta}$ on $\theta_1 \leq \theta \leq \theta_2$ and $r = u + iv$. But $-2u + 1 = -(R + 2)/R < -1$, hence the integral on the left diverges and ϕ is not square summable.

In case 2 the coefficient of u' has a double zero at $\alpha = -a/2c$ where $|\alpha|^2 = a^2/4|c|^2 = 1$. The functions $\phi_+(z) = (z - \alpha)^{-2}e^{r(z-\alpha)^{-1}}$, $r = (b - a - i)/c$ and $\phi_-(z) = (z - \alpha)^{-2}e^{r(z-\alpha)^{-1}}$, $r = (b - a + i)/c$ are fundamental solutions for (2.2) and (2.3) respectively. Let us introduce polar coordinates at $z = \alpha$ so that $z - \alpha = \rho e^{i\theta}$ and let us agree to set $\theta = 0$ so that for $|z| < 1$, the argument of $z - \alpha$ is restricted to the intervals $0 \leq \theta < \pi/2$ and $3\pi/2 < \theta < 2\pi$. Let $r = u + iv$, then

$$\begin{aligned} |\phi_{\pm}(z)| &= |\rho^{-2}e^{-i2\theta}e^{(u+iv)(\cos\theta-i\sin\theta)/\rho}| \\ &= \rho^{-2}e^{(u\cos\theta+v\sin\theta)/\rho} . \end{aligned}$$

We note that u and v are not both zero, for then $b - a \pm i = 0$ where a and b are real. Now consider the function $F(\theta) = u \cos \theta + v \sin \theta$. If $u > 0$, $F(0) = u > 0$ and by continuity there exist θ_1 and θ_2 such that $F(\theta) \geq u/2 > 0$ for $\theta_1 \leq \theta \leq \theta_2 < \pi/2$, similarly if $v > 0$, $F(\pi/2) = v$ and there exist θ_1 and θ_2 such that $F(\theta) \geq v/2 > 0$ for $\theta_1 \leq \theta \leq \theta_2 \leq \pi/2$. If $v < 0$, $F(3\pi/2) = -v > 0$ and there exist θ_1 and θ_2 such that $F(\theta) \geq -v/2 > 0$ for $3\pi/2 < \theta_1 \leq \theta \leq \theta_2$. Hence for all $r = u + iv$, except for the case $u < 0, v = 0$, there exists a $M > 0$ and suitable θ_1 and θ_2 for which $F(\theta) \geq M, \theta_1 \leq \theta \leq \theta_2$. This case requires only a minor modification which will be provided shortly. It is easy to see that for given θ_1 and θ_2 we can find $\delta > 0$ for which the regions $W_\varepsilon = \{z | \varepsilon \leq \rho < \delta, \theta_1 \leq \theta \leq \theta_2\}$ lie entirely within the disc for $0 < \varepsilon < \delta$.

Now consider $\|\phi_{\pm}\|^2$:

$$\begin{aligned} \int_D \int |\phi_{\pm}(z)|^2 dx dy &\geq \lim_{\varepsilon \rightarrow 0} \int_{W_\varepsilon} \int |\phi_{\pm}(z)|^2 dx dy \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\theta_1}^{\theta_2} \int_{\varepsilon}^{\delta} \rho^{-3} e^{2F(\theta)/\rho} d\rho d\theta \\ &\geq \lim_{\varepsilon \rightarrow 0} (\theta_2 - \theta_1) \int_{\varepsilon}^{\delta} e^{2M/\rho} \rho^{-3} d\rho . \end{aligned}$$

Since $\int_0^\delta e^{2M/\rho} \rho^{-3} d\rho$ diverges it follows that the ϕ_{\pm} are not square summable, provided r is not a negative number. When $r = u + iv = u < 0$ we merely agree to set $\theta = 0$ so that for $|z| < 1$ the argument of $z - \alpha$ is restricted to the interval $\pi/2 < \theta < 3\pi/2$. Then $F(\pi) = -u > 0$ and the argument is the same as before.

In case 3, $a^2 > 4|c|^2$, the coefficient of u' has distinct zeros at $\alpha = (-a + R)/2c$ and $\beta = (-a - R)/2c$ where $R = (a^2 - 4|c|^2)^{1/2} > 0$. For $a > 0$,

$$|\beta| = \frac{R + a}{2|c|} > \frac{a}{2|c|} > 1,$$

and therefore $|\alpha| < 1$. For $a < 0$,

$$|\alpha| = \frac{R - a}{2|c|} > \frac{|a|}{2|c|} > 1,$$

and therefore $|\beta| < 1$. Without loss of generality we assume $|\alpha| < 1$, and $|\beta| > 1$. For $|z| < |\alpha| < 1$, the functions ϕ_+ and ϕ_- given by

$$\begin{aligned} \phi_+(z) &= (z - \alpha)^{-r}(z - \beta)^{-t} \\ \phi_-(z) &= (z - \beta)^{-s}(z - \alpha)^{-u} \end{aligned}$$

where $r = (R + b - a)/R - i/R$ and $s = (R + b - a)/R + i/R$, are fundamental solutions for $Lu = iu$ and $Lu = -iu$ respectively. Now suppose ψ is any nontrivial element of \mathcal{H} which satisfies $Lu = \pm iu$. In particular ψ is analytic for $|z| < |\alpha| < 1$. From uniqueness results this implies that $\psi(z) = c\phi_{\pm}(z)$ for $|z| < |\alpha|$, where $c \neq 0$. By the identity theorem for analytic functions this implies $\psi(z) = c\phi_{\pm}(z)$ for $|z| < 1$, hence $\phi_{\pm}(z)$ is analytic in $|z| < 1$. But $\phi_{\pm}(z)$ has a singularity at $|\alpha| < 1$, therefore, the equations $Lu = \pm iu$ have no nontrivial solutions in \mathcal{H} .

3. In this section we obtain conditions on the coefficients of L which insure that for all λ every solution of $L\phi = \lambda\phi$ is in \mathcal{H} . If L is a formally symmetric operator satisfying these conditions the defect indices of the operator T_0 are equal to the order of L and T_0 has a self-adjoint extension in \mathcal{H} .

In [2] it was shown that if $L = \sum_{k=0}^n p_k D^k$ is formally symmetric then the p_i are polynomials of degree at most $n + i$. Regarding such L with polynomial coefficients we have

THEOREM 3.1. *Let $L = \sum_{k=0}^n p_k D^k$ where $n \geq 2$, $p_n(0) \neq 0$, and $p_k = \sum_{i=0}^{n+k} a_i(k)z^k$, and*

$$\begin{aligned} (3.1) \quad A &= |a_0(n)|^{-1} \sum_{i=1}^{2n} |a_i(n)|, \\ \hat{B} &= n(n + 1)/2, \quad \text{and} \\ B &= |a_0(n)|^{-1} \sum_{i=1}^{2n} |a_i(n)n[(n + 1)/2 - i] + a_{i-1}(n - 1)|. \end{aligned}$$

If $A < 1$ or $A = 1$ and $B < \hat{B}$ then every solution of $L\phi = 0$ is in \mathcal{H} .

Proof. Since $p_n(0) = a_0(n) \neq 0$, every solution of $Lu = 0$ at the origin is analytic in some neighborhood of the origin. Let $\phi(z) = \sum_{j=0}^{\infty} b_j z^j$ be any such solution, we will show that there exists a positive constant K and positive integer p such that $|b_j| \leq K j^{-1/p}$ for j sufficiently large. Consequently the series $\sum_{j=0}^{\infty} |b_j|^2 / (j + 1)$ converges and ϕ belongs to \mathcal{H} .

We begin by obtaining a recursion formula for the b_j . Substituting $\phi(z) = \sum_{j=0}^{\infty} b_j z^j$ into the equation $L\phi(z) = 0$ we obtain

$$L\phi(z) = \sum_{j=0}^{\infty} \sum_{k=0}^n \sum_{i=0}^{n+k} a_i(k) \pi_k(j - i + k) b_{j-i+k} z^j,$$

where

$$\begin{aligned} \pi_k(\lambda) &= \lambda(\lambda - 1) \cdots (\lambda - k + 1) & k \leq \lambda \\ &= 0 & k > \lambda. \end{aligned}$$

Hence $L\phi = 0$ if and only if the following relationship holds for all j .

$$(3.2) \quad \sum_{k=0}^n \sum_{i=0}^{n+k} a_i(k) \pi_k(j - i + k) b_{j-i+k} = 0.$$

Hence,

$$\begin{aligned} &\sum_{k=0}^{n-1} \sum_{i=0}^{n+k} a_i(k) \pi_k(j - i + k) b_{j-i+k} \\ &+ \sum_{i=1}^{2n} a_i(n) \pi_n(j - i + n) b_{j-i+n} + a_0(n) \pi_n(j + n) b_{j+n} = 0. \end{aligned}$$

Noting that the sums involve only the b_{j-n} thru b_{j+n-1} (where $j > n$) and $\pi_n(j + n)$ never vanishes we may solve for b_{j+n} to obtain

$$(3.3) \quad b_{j+n} = - (S_1 + S_2) / a_0(n) \pi_n(j + n),$$

where

$$S_1 = \sum_{i=1}^{2n} a_i(n) \pi_n(j - i + n) b_{j-i+n},$$

and

$$S_2 = \sum_{k=0}^{n-1} \sum_{i=0}^{n+k} a_i(k) \pi_k(j - i + k) b_{j-i+k},$$

for $j > n$.

We now investigate the nature of S_1 and S_2 as polynomials in j . It can be shown that $\pi_n(j + n - 1)$ is a polynomial of degree n in j ,

$$(3.4) \quad \pi_n(j + n - i) = j^n + \left[\frac{n(n + 1)}{2} - in \right] j^{n-1} + \dots,$$

for $i = 1, \dots, 2n$. Using (3.4) in (3.3) we obtain

$$(3.5) \quad \begin{aligned} S_1 = & j^n \sum_{i=1}^{2n} a_i(n) b_{j-i+n} \\ & + j^{n-1} \sum_{i=1}^{2n} a_i(n) \left[\frac{n(n + 1)}{2} - in \right] \\ & + \text{lower powers of } j. \end{aligned}$$

Now consider S_2 . Since $\pi_k(j - i + k)$ is a polynomial of degree k in j , an examination of (3.3) shows that S_2 is a polynomial of degree $n - 1$ in j , and that the only terms which contribute to the coefficient of j^{n-1} are those corresponding to $k = n - 1$. Hence

$$(3.6) \quad \begin{aligned} S_2 = & j^{n-1} \sum_{i=0}^{2n-1} a_i(n - 1) b_{j-i+n-1} \\ & + \text{lower powers of } j. \end{aligned}$$

Combining (3.5) and (3.6) we obtain

$$(3.7) \quad \begin{aligned} S_1 + S_2 = & j^n \sum_{i=1}^{2n} a_i(n) b_{j-i+n} \\ & + j^{n-1} \sum_{i=1}^{2n} \left[a_i(n) \left(\frac{n(n + 1)}{2} - in \right) + a_{i-1}(n - 1) \right] b_{j-i+n} \\ & + \dots, \quad (j > n). \end{aligned}$$

Since $\pi_n(j + n) = j^n + (n(n + 1))/2j^{n-1} + \dots$, is always positive (3.3) yields

$$(3.8) \quad |b_{j+n}| = \frac{|S_1 + S_2|}{|a_0(n)| [j^n + \hat{B}j^{n-1} + \dots]}.$$

We now estimate $|S_1 + S_2|$. Let $M(j) = \text{Max}(|b_{j-n}|, \dots, |b_{j+n-1}|)$, then it follows from (3.1) and (3.7) that $|S_1 + S_2| \leq |a_0(n)| [M(j)Aj^n + M(j)Bj^{n-1} + \dots]$. Hence

$$(3.9) \quad |b_{j+n}| \leq \frac{Aj^n + Bj^{n-1} + \dots}{j^n + \hat{B}j^{n-1} + \dots} M(j)$$

for $j > n$, where A, B , and \hat{B} are given by (3.1).

Consider the estimate (3.9) for $|b_{j+n}|$,

$$(3.10) \quad |b_{j+n}| \leq Q(j)M(j) \quad j > n,$$

where $Q(j) = (Aj^n + Bj^{n-1} + \dots)/(j^n + \hat{B}j^{n-1} + \dots)$. We note that for fixed ζ , $Q(j) \leq 1 + \zeta j^{-1}$ for j sufficiently large if and only if $Aj^n +$

$Bj^{n-1} + \dots \leq j^n + (\hat{B} + \zeta)j^{n-1} + \dots$. Hence if $A < 1$ or $A = 1$ and $B < \hat{B} + \zeta$ we have

$$(3.11) \quad Q(j) \leq 1 + \zeta j^{-1}$$

for j sufficiently large. Now consider the expression

$$(1 + \zeta(j + 1)^{-1})(j - n + 1)^{-1/p},$$

where $\zeta < 0$ and p a positive integer. It is not difficult to see that this is dominated by $(j + n + 1)^{-1/p}$ for j sufficiently large if and only if

$$j^{p+1} + (p + p\zeta + n + 1)j^p + \dots \leq j^{p+1} + (p - n + 1)j^p + \dots,$$

for j sufficiently large. Hence, we have

$$(3.12) \quad (1 + \zeta(j + 1)^{-1})(j - n + 1)^{-1/p} \leq (j + n + 1)^{-1/p}$$

for j sufficiently large if $p \geq -2n\zeta^{-1}$.

We now show that there exists a positive constant K and positive integer p for which $|b_j| \leq Kj^{-1/p}$, j sufficiently large. By hypothesis either $A < 1$ or $A = 1$ and $B < \hat{B}$. If $A < 1$ let $\zeta = -1$ and $p = 2n$, if $A = 1$, select ζ such that $B - \hat{B} < \zeta < 0$ and $p > -2n\zeta^{-1}$. For j sufficiently large, say $j > j_1$, (3.11) and (3.12) hold. Set

$$K = \max_{j \leq j_1+n} |b_j|j^{1/p}$$

so that $|b_j| \leq Kj^{-1/p}$ for $j \leq j_1 + n$. Using (3.10) and (3.11) it follows that

$$|b_{j_1+n+1}| \leq (1 + \zeta(j_1 + 1)^{-1})M(j_1 + 1),$$

where

$$\begin{aligned} M(j_1 + 1) &= \text{Max} (K(j_1 - n + 1)^{-1/p}, \dots, K(j_1 + n)^{-1/p}) \\ &= K(j_1 - n + 1)^{-1/p}. \end{aligned}$$

Hence $|b_{j_1+n+1}| \leq (1 + \zeta(j_1 + 1)^{-1})K(j_1 - n + 1)^{-1/p}$, and using (3.12) this yields

$$(3.13) \quad |b_{j_1+n+1}| \leq K(j_1 + n + 1)^{-1/p}.$$

We now proceed inductively to establish

$$(3.14) \quad |b_{j_1+n+k}| \leq K(j_1 + n + k)^{-1/p} \quad k = 2, 3, \dots$$

Let $K_1 = \max_{j \leq j_1+n+1} |b_j|j^{1/p}$, now $K_1 = \max \{K, |b_{j_1+n+1}|(j_1 + n + 1)^{1/p}\} \leq K$, making use of (3.13). Using (3.11) yields

$$|b_{j_1+n+2}| \leq (1 + \zeta(j_1 + 2)^{-1})M(j_1 + 2)$$

where

$$M(j_1 + 2) = \text{Max} (K(j_i - n + 2)^{-1/p}, \dots, K(j_1 + n + 1)^{-1/p}) \\ = K(j_1 - n + 2)^{-1/p} .$$

Using (3.12) it follows that

$$|b_{j_1+n+2}| \leq K(j_1 + n + 2)^{-1/p} .$$

Continuing on in this manner we establish (3.14) and the theorem is proved.

We note that the conditions (3.1) of Theorem 3.1 involve only the coefficients of the polynomials p_n and p_{n-1} , hence if L satisfies the conditions of (3.1) so do the operators $L \pm i$. Hence we have established the following.

THEOREM 3.2. *Let L be a formally symmetric operator which satisfies (3.1), then the associated operator T_0 has defect indices $n_+ = n_- = n$.*

COROLLARY 3.3. *The operator $L = (c_1z^4 + \bar{c}_1)d^2/dz^2 + (6c_1z^3 + c_3z^2 + a_2z + \bar{c}_3)d/dz + (6c_1z^2 + 2c_3z + a_3)$, where a_3 and a_2 are real and $|c_1| > |c_3| + |a_2|/2$, has self-adjoint extensions.*

Proof. Applying the algorithm given in Theorem 2.3 of [2] the general second order formally symmetric operator has coefficients

$$p_2(z) = c_1z^4 + c_2z^3 + a_1z^2 + \bar{c}_2z + \bar{c}_1 \\ p_1(z) = 6c_1z^3 + (c_3 + 3c_2)z^2 + a_2z + \bar{c}_3 \\ p_0(z) = 6c_1z^2 + 2c_3z + a_3 ,$$

where $a_1, a_2,$ and a_3 are real.

Now $A = (|c_1| + 2|c_2| + |a_1|)/|c_1| \geq 1$ and $A = 1$ if and only if $c_2 = a_1 = 0$. Now $\hat{B} = 3$ and $B = (|c_1| + |a_2| + 2|c_3|)/|c_1| < 3$ if and only if $|c_1| > |c_3| + |a_2|/2$. Hence the result follows from the previous theorem.

REFERENCES

1. N. Dunford and J. Schwartz, *Linear Operators*, Part II, Interscience Publishers, 1963.
2. A. L. Villone, *Self-adjoint differential operators*, Pacific J. Math., **35** (1970), 517-531.

Received September 15, 1972 and in revised form January 5, 1973. This work was supported in part by NSF Grant GP-3594.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$48.00 a year (6 Vols., 12 issues). Special rate: \$24.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by
Pacific Journal of Mathematics
All Rights Reserved

Wm. R. Allaway, <i>On finding the distribution function for an orthogonal polynomial set</i>	305
Eric Amar, <i>Sur un théorème de Mooney relatif aux fonctions analytiques bornées</i>	311
Robert Morgan Brooks, <i>Analytic structure in the spectrum of a natural system</i>	315
Bahattin Cengiz, <i>On extremely regular function spaces</i>	335
Kwang-nan Chow and Moses Glasner, <i>Atoms on the Royden boundary</i>	339
Paul Frazier Duvall, Jr. and Jim Maxwell, <i>Tame Z^2-actions on E^n</i>	349
Allen Roy Freedman, <i>On the additivity theorem for n-dimensional asymptotic density</i>	357
John Griffin and Kelly Denis McKennon, <i>Multipliers and the group L_p-algebras</i>	365
Charles Lemuel Hagopian, <i>Characterizations of λ connected plane continua</i>	371
Jon Craig Helton, <i>Bounds for products of interval functions</i>	377
Ikuko Kayashima, <i>On relations between Nörlund and Riesz means</i>	391
Everett Lee Lady, <i>Slender rings and modules</i>	397
Shozo Matsuura, <i>On the Lu Qi-Keng conjecture and the Bergman representative domains</i>	407
Stephen H. McCleary, <i>The lattice-ordered group of automorphisms of an α-set</i>	417
Stephen H. McCleary, <i>$o - 2$-transitive ordered permutation groups</i>	425
Stephen H. McCleary, <i>o-primitive ordered permutation groups. II</i>	431
Richard Rochberg, <i>Almost isometries of Banach spaces and moduli of planar domains</i>	445
R. F. Rossa, <i>Radical properties involving one-sided ideals</i>	467
Robert A. Rubin, <i>On exact localization</i>	473
S. Sribala, <i>On Σ-inverse semigroups</i>	483
H. M. (Hari Mohan) Srivastava, <i>On the Konhauser sets of biorthogonal polynomials suggested by the Laguerre polynomials</i>	489
Stuart A. Steinberg, <i>Rings of quotients of rings without nilpotent elements</i>	493
Daniel Mullane Sunday, <i>The self-equivalences of an H-space</i>	507
W. J. Thron and Richard Hawks Warren, <i>On the lattice of proximities of Čech compatible with a given closure space</i>	519
Frank Uhlig, <i>The number of vectors jointly annihilated by two real quadratic forms determines the inertia of matrices in the associated pencil</i>	537
Frank Uhlig, <i>On the maximal number of linearly independent real vectors annihilated simultaneously by two real quadratic forms</i>	543
Frank Uhlig, <i>Definite and semidefinite matrices in a real symmetric matrix pencil</i>	561
Arnold Lewis Villone, <i>Self-adjoint extensions of symmetric differential operators</i>	569
Cary Webb, <i>Tensor and direct products</i>	579
James Victor Whittaker, <i>On normal subgroups of differentiable homeomorphisms</i>	595
Jerome L. Paul, <i>Addendum to: "Sequences of homeomorphisms which converge to homeomorphisms"</i>	615
David E. Fields, <i>Correction to: "Dimension theory in power series rings"</i>	616
Peter Michael Curran, <i>Correction to: "Cohomology of finitely presented groups"</i>	617
Billy E. Rhoades, <i>Correction to: "Commutants of some Hausdorff matrices"</i>	617
Charles W. Trigg, <i>Corrections to: "Versum sequences in the binary system"</i>	619