Pacific Journal of Mathematics

TENSOR AND DIRECT PRODUCTS

CARY WEBB

Vol. 49, No. 2

June 1973

TENSOR AND DIRECT PRODUCTS

CARY WEBB

Let R be an associative ring with 1, E a unitary right module, and $(F_i)_{i \in I}$ a family of unitary left modules. Let $f: E \bigotimes_R \prod F_i \to \prod (E \bigotimes_R F_i)$ be the canonical map. THEOREM. f is bijective (surjective) for all families (F_i) iff E is finitely presented (finitely generated). Theorem. If R is a Dedekind domain or is commutative artinian and every F_i is flat, then f is injective. COROLLARY. If R is a Dedekind domain or is commutative artinian, every F_i is flat and $E \bigotimes_R F_i$ is reduced, then $E \bigotimes_R \prod F_i$ is reduced. THEOREM. If R is a Dedekind domain or is commutative artinian, E_j is flat, f is injective for every E_j (e.g. E_j projective) and E is pure in $\prod E_j$, then f is injective. THEOREM. If R is a Dedekind domain and Eis flat then f is injective for E iff f is injective for Hom (F,E) for all modules F. THEOREM. If R is a Dedekind domain and f is injective for E for all families (F_i) then E is reduced. THEOREM. If R is commutative and f is always injective then R must be artinian. The converse holds for serial rings.

Introduction. If E is a right module and $(F_i)_{i \in I}$ is a family of left modules over an associative ring R with 1 then it is always true that the groups $E \bigotimes_R (\bigoplus_{i \in I} F_i)$ and $\bigoplus_{i \in I} (E \bigotimes_R F_i)$ are isomorphic. However, it is not hard to see that the groups $E \bigotimes_R \prod_{i \in I} F_i$ and $\prod_{i \in I} (E \bigotimes_R F_i)$ are not necessarily isomorphic (e.g. if R is the integers let E be the rationals, I the natural numbers and, for a fixed prime p, F_n the cyclic group of order p^n . This example is found in ([8], p. 257, Ex. 10).

It is our purpose to study the relationship between the groups $E \bigotimes_{\mathbb{R}} \prod_{i \in I} F_i$ and $\prod_{i \in I} (E \bigotimes_{\mathbb{R}} F_i)$. We will do so in terms of the natural homomorphism

$$f: E \bigotimes_{R} \prod_{i \in I} F_{i} \longrightarrow \prod_{i \in I} (E \bigotimes_{R} F_{i})$$

which sends a generator $x \bigotimes (y_i)_{i \in I}$ of $E \bigotimes_R \prod_{i \in I} F_i$ onto $(x \bigotimes y_i)_{i \in I}$ in $\prod_{i \in I} (E \bigotimes_R F_i)$.

In §§1 and 2 we investigate the properties of f over an arbitrary ring. It is relatively easy to show that f is surjective (bijective) for all families $(F_i)_{i \in I}$ if and only if E is finitely generated (finitely presented). In §§3 and 4 we study the more difficult problem of when f is injective. In §3 the ring is a Dedekind domain. In §4 the ring is commutative artinian.

We will always denote by R an associative ring with 1, by E a right R-module, and by $(F_i)_{i \in I}$ a family of left R-modules. Modules

are all unitary.

We will often use the notions of flatness and purity. Briefly, a left *R*-module *F* is said to be flat if the functor () $\bigotimes_R F$ is exact. A submodule *E'* of a right *R*-module *E* is said to be pure if () $\bigotimes_R F$ is exact on $0 \to E' \xrightarrow{\text{incl}} E \xrightarrow{\text{proj}} E/E' \to 0$ for every right module *F*.

Over a domain it is well known that torsion free modules are flat iff the domain is Prüfer. It is also known that over a domain, E' is pure in E means $rE' = E' \cap rE$ for all $r \in R$ iff the domain is Prüfer. For this last fact see [13].

Many of our arguments involve a diagram chase. At these points we usually will be able simply to cite the following.

Snake Lemma [2, Ch. 1, §1, No. 4, Prop. 2]. Suppose



is a commutative diagram of abelian groups with exact rows. Then this diagram can be embedded in the following commutative diagram with top and bottom row not necessarily exact.



The following are true of this diagram

- (i) $v_1 \circ u_1 = 0$. If u' is injective then the top row is exact.
- (ii) $v_2 \circ u_2 = 0$. If v is surjective the bottom row is exact.

(iii) If u' is injective and v is surjective there exists a unique homomorphism d such that the sequence (u_1, v_1, d, u_2, v_2) is exact.

For convenience we will omit writing the indexing set whenever there is no possibility of confusion. For example, we will write $\prod F_i$, $\bigoplus F_i$ and (F_i) , omitting the index set *I*. fg and fp will denote, respectively, finitely generated and finitely presented. Instead of writing $E \bigotimes_R F$ we will simply write $E \bigotimes F$. Often an obvious homomorphism will not be defined explicitly or even named. *Z* will be the ring of integers, *Q* the field of rational numbers, and, for a positive integer

n, Z(n) will denote the cyclic group of order n.

This study is a doctoral dissertation prepared under the direction of Professor Edgar Enochs and submitted to the University of Kentucky.

1. Surjectivity and bijectivity.

PROPOSITION 1.1. If E is fg and projective then f is bijective.

Proof. If E' is fg and free then for some positive integer n, $E' \approx R^n$. We have the canonical isomorphisms

$$E' \otimes \prod F_i \approx R^n \otimes \prod F_i \approx (\prod F_i)^n$$

and

$$\prod (E' \otimes F_i) \approx \prod (R^n \otimes F_i) \approx \prod F_i^n.$$

So f is just the canonical isomorphism $(\prod F_i)^n \approx \prod F_i^n$.

If E is fg and projective then E is a direct summand of such an E'. We show later (Prop. 2.1) that if f is injective for E' then f is injective for any summand of E'.

PROPOSITION 1.2. f is surjective (for all families (F_i)) iff E is fg.

Proof. Suppose E is fg, n is a positive integer, and $\mathbb{R}^n \to E \to 0$ is exact. Consider the commutative diagram with exact rows

$$\begin{array}{cccc} R^n \otimes \prod F_i & \longrightarrow E \otimes \prod F_i & \longrightarrow 0 \\ & & \downarrow^{f'} & & \downarrow^f \\ \prod (R^n \otimes F_i) & \longrightarrow \prod (E \otimes F_i) & \longrightarrow 0 \end{array}$$

f' is bijective (Prop. 1.1). The snake lemma tells us f is surjective. On the other hand, suppose E is not fg. Let I = E (E considered as a set), $F_i = {}_{R}R$. Write $E = \{x_i\}$. Then $(x_i \otimes 1)_{i \in I} \in \prod (E \otimes F_i)$ is not in imf. So f is not surjective.

REMARK 1. In order to show E fg all we need to show is that f is surjective for all families (F_i) with $F_i = {}_{\scriptscriptstyle R}R$.

REMARK 2. The part of Proposition 1.2 that supposes E fg is found in [3, Ch. II, Ex. 2].

If g and h are injective homomorphisms of right, respectively, left R-modules, having domains E and F, respectively, we denote im $(g \otimes h)$ as $[E \otimes F]$.

PROPOSITION 1.3. f is bijective (for all families (F_i)) iff E is fp.

Proof. Suppose E is fp and $0 \to K \to L \to E \to 0$ is a finite presentation of E. Consider the commutative diagram with exact rows

 f_1 is surjective because K is fg (Prop. 1.2). f_2 is bijective because L is fg and free (Prop. 1.1). By the snake lemma, therefore, f is bijective.

On the other hand, suppose f is bijective for all families (F_i) . Since f is surjective, E is fg (Prop. 1.2). So we have an exact sequence $0 \to K \to L \to E \to 0$ where L is fg and free. By Remark 1 following Prop. 1.2 to show K fg it suffices to show f is surjective for K for all families (F_i) , $F_i = {}_R R$.

For such a family consider the commutative diagram with exact rows

$$\begin{array}{ccc} K \otimes \prod F_i \longrightarrow L \otimes \prod F_i \longrightarrow E \otimes \prod F_i \longrightarrow 0 \\ & & \downarrow^{f_1} & \downarrow^{f_2} & \downarrow^{f} \\ 0 \longrightarrow \prod (K \otimes F_i) \longrightarrow \prod (L \otimes F_i) \longrightarrow \prod (E \otimes F_i) \longrightarrow 0 \end{array}.$$

 f_2 is bijective because L is fg and free (Prop. 1.1). f is bijective by assumption. Therefore, by the snake lemma, f_1 is surjective.

REMARK 1. It is found in [3, Ch. II, Ex. 2] that if R is right noetherian and E fg then f is bijective. More generally, [2, Ch. 1, Ex. 8] states that if E is fp then f is bijective.

REMARK 2. The first half of the proof of Prop. 1.3 shows that if f is injective for E then f is injective for any quotient of E by a fg submodule.

2. Injectivity.

PROPOSITION 2.1. Suppose J is a set and E_j is a right R-module for every $j \in J$. Let $E \approx \bigoplus E_j$. Then f is injective for E iff f is injective for every E_j . If E' is pure in E and f is injective for E then f is injective for E'.

Proof. It is easy to see that if J is finite and f is injective for every E_j then f is injective for E. Suppose J is infinite,

$$z=\sum\limits_{{\lambda}=1}^n x_{{\lambda}} \otimes y_{{\lambda}} \in E \otimes \prod F_i$$
 ,

and f(z) = 0. Let J' be the support of $\{x_{\lambda}\}_{\lambda=1}^{n}$. Consider the commutative diagram

where g and h are the obvious injective homomorphisms. If $z' = \sum_{\lambda=1}^{n} x_{\lambda} \otimes y_{\lambda} \in (\bigoplus_{J'} E_{j}) \otimes \prod F_{i}$ then g(z') = z. But since J' is finite, f' is injective. Therefore z' = 0, hence z = 0. So f is injective.

For the remainder of the proposition suppose f is injective for Eand E' is pure in E. Consider the commutative diagram with exact rows

$$\begin{array}{ccc} 0 \longrightarrow E' \otimes \prod F_i \longrightarrow E \otimes \prod F_i \\ & & \downarrow^{f'} & \downarrow^f \\ 0 \longrightarrow \prod (E' \otimes F_i) \longrightarrow \prod (E \otimes F_i) \end{array}$$

Since f is injective the snake lemma says f' is injective.

REMARK. It is not hard to see that if E' is a pure submodule of E and f is injective for E' and E/E' then f is injective for E. Merely apply the snake lemma to the following commutative diagram with exact rows

$$\begin{array}{cccc} 0 \longrightarrow E' \otimes \prod F_i \longrightarrow E \otimes \prod F_i \longrightarrow E/E' \otimes \prod F_i \longrightarrow 0 \\ & & \downarrow & & \downarrow \\ 0 \longrightarrow \prod (E' \otimes F_i) \longrightarrow \prod (E \otimes F_i) \longrightarrow \prod (E/E' \otimes F_i) \longrightarrow 0 \end{array}.$$

PROPOSITION 2.2. If E is the quotient of a projective by a fg submodule then f is injective.

Proof. If E is free and $E \approx \bigoplus R$ then we have the canonical isomorphisms

$$E \otimes \prod F_i \approx (\bigoplus R) \otimes \prod F_i \approx \bigoplus \prod F_i$$

and

$$\prod (E \otimes F_i) \approx \prod (\bigoplus R \otimes F_i) \approx \prod \bigoplus F_i.$$

So f is the injection $\bigoplus \prod F_i \to \prod \bigoplus F_i$.

If E is projective then E is a summand of a free module. Hence

f is injective for E (Prop. 2.1). The proposition now follows from Remark 2 following Prop. 1.3.

We want to use the following result of Matlis.

THEOREM 2.3. (Matlis, [12].) If R is right noetherian and E is injective then E is the direct sum of indecomposable, injective submodules.

It will be useful to consider f when every F_i of the family (F_i) is flat. Such a family will be called a flat family.

PROPOSITION 2.4. f is injective for all flat families (F_i) iff R is right noetherian and f is injective for all indecomposable injective E for all flat families (F_i) .

Proof. Suppose f is injective for all flat families (F_i) and E is a right ideal of R. Recall that to show E is fg it suffices to show f is surjective for E for all families (F_i) with $F_i = {}_{R}R$ (Remark 1, following Prop. 1.2). So suppose (F_i) is such a family. Consider the commutative diagram with exact rows

 f_1 is bijective. f_2 is injective by supposition. By the snake lemma f is surjective.

For the converse, suppose (F_i) is a flat family and $E' = \bigoplus E_j$ is the injective envelope of E where E_j is indecomposable injective. By Prop. 2.1, f is injective for E'. Consider the commutative diagram with exact rows

$$\begin{array}{cccc} 0 \longrightarrow E \otimes \prod F_i \longrightarrow E' \otimes \prod F_i \\ & & \downarrow^f & & \downarrow^{f'} \\ 0 \longrightarrow \prod (E \otimes F_i) \longrightarrow \prod (E' \otimes F_i) \ . \end{array}$$

(The top row is exact because over a right noetherian ring the direct product of flat left modules is flat, [3, Ch. VI, Ex. 4].) Since f' is injective the snake lemma says f is also injective.

3. Dedekind domains. Let K be the field of fractions of a domain R. If R is a Dedekind domain the indecomposable injective R-modules are K and, for primes P, the P-primary components of K/R [9, Thm. 7]. Adopting the notation that is standard when R

is the ring of integers we will denote the P-primary component of K/R as $R(P^{\infty})$.

In order to establish that Dedekind domains possess the property of Prop. 2.4 we prove the following lemma.

LEMMA 3.1. If S is a multiplicative subset of an arbitrary commutative ring $R, E = S^{-1}R$ and (F_i) is a family of torsion free R-modules then f is injective.

Proof. If $0 \in S$ then $S^{-1}R = 0$. So assume $0 \notin S$. We have the canonical isomorphisms [2, Ch. II, §2, no. 7, Prop. 18], $S^{-1}R \otimes \prod F_i \approx S^{-1}(\prod F_i)$ and $\prod (S^{-1}R \otimes F_i) \approx \prod (S^{-1}F_i)$. So we can think of f as the canonical map $S^{-1}(\prod F_i) \longrightarrow \prod (S^{-1}F_i)$. Suppose $f((y_i)/s) = (y_i/s) = 0$. Then for every $i \in I$ there exists $t_i \in S$ such that $t_i y_i = 0$. Since $t_i \neq 0$, $y_i = 0$. Hence f is injective.

PROPOSITION 3.2. If R is a Dedekind domain then f is injective for all flat families (F_i) .

Proof. By Prop. 2.4 and Prop. 2.1 it suffices to show f is injective for K and K/R. But since we have the following commutative diagram with exact rows

$$0 \longrightarrow R \otimes \prod F_{i} \longrightarrow K \otimes \prod F_{i} \longrightarrow K/R \otimes \prod F_{i} \longrightarrow 0$$
$$\downarrow f_{1} \qquad \qquad \qquad \downarrow f_{2} \qquad \qquad \qquad \downarrow f_{3}$$
$$0 \longrightarrow \prod (R \otimes F_{i}) \longrightarrow \prod (K \otimes F_{i}) \longrightarrow \prod (K/R \otimes F_{i}) \longrightarrow 0$$

where f_1 is bijective we know from the snake lemma that f_2 is injective iff f_3 is injective. That f_2 is injective is Lemma 3.1. The proof is now complete.

Since a Prüfer domain is Dedekind iff it is noetherian we have an immediate corollary.

COROLLARY 3.3. If R is a Prüfer domain then f is injective for all flat families (F_i) iff R is Dedekind.

There is an interesting generalization of Prop. 3.2. We need the following lemma whose proof is immediate from the definitions.

LEMMA 3.4. If R is a Prüfer domain and (F_i, g_i^i) is an inverse system of flat R-modules then $\lim F_i$ is a pure submodule of $\prod F_i$.

THEOREM 3.5. If R is a Dedekind domain and (F_i, g_i^i) is an

inverse system of flat R-modules then the canonical homomorphism

$$F: E \otimes \varprojlim F_i \longrightarrow \varprojlim (E \otimes F_i)$$

is injective.

Proof. Clearly $(E \otimes F_i, id \otimes g_i^j)$ is an inverse system and F is a homomorphism. To show F injective consider the commutative diagram

where u and v are the obvious maps. u is injective (Lemma 3.4). f is injective (Prop. 3.2). By the Snake lemma F must also be injective.

REMARK. Suppose R is any ring for which f is injective for all flat families (F_i) . Then it is easy to see that F is injective for the inverse system of flat left modules (F_i) iff $\lim_{i \to \infty} F_i$ is pure in $\prod F_i$. Just chase the following commutative diagram

$$\begin{array}{ccc} E \otimes \varprojlim F_i \longrightarrow E \otimes \prod F_i \\ & \swarrow F & \downarrow f \\ \varprojlim (E \otimes F_i) \longrightarrow \prod (E \otimes F_i) \end{array} .$$

In Prop. 2.2 we established that f is injective whenever E is projective. We will see later (Thm. 3.13) that this cannot be extended to the case when E is flat. But when R is a Dedekind domain we do have the following generalization.

THEOREM 3.6. If R is a Dedekind domain, $(E_j)_{j\in J}$ is a family of flat R-modules such that f is injective for every E_j (e.g. E_j is projective), and E is isomorphic to a pure submodule of $\prod E_j$, then f is injective for all families (F_i) .

Proof. Consider the following diagram where all maps are the obvious ones

$$\begin{split} \prod_{J} E_{j} & \otimes \prod_{I} F_{i} \stackrel{f}{\longrightarrow} \prod_{I} (\prod_{J} E_{j} \otimes F_{i}) \stackrel{f_{1}}{\longrightarrow} \prod_{I} \prod_{J} (E_{j} \otimes F_{i}) \\ & \downarrow^{f_{2}} \\ \prod_{J} (E_{j} \otimes \prod_{I} F_{i}) \\ & \downarrow^{f_{3}} \\ \prod_{J} \prod_{I} (E_{j} \otimes F_{i}) \\ & \downarrow^{f_{4}} \\ \prod_{I} \prod_{J} (E_{j} \otimes F_{i}) . \end{split}$$

 f_1 and f_2 are injective (Prop. 3.2) and f_3 is injective by hypothesis. Therefore, $f_4 \circ f_3 \circ f_2 = f_1 \circ f$ is injective. Hence f is injective. Now apply Prop. 2.1.

REMARK 1. We have not used any specific property of Dedekind domains in Thm. 3.6. The theorem holds for any ring for which f is injective for flat families (F_i) .

REMARK 2. If in Thm. 3.6, J is infinite then $\prod_J Z$ is not free (e.g. [8, Thm. 19.2]). Indeed it is not hard to see that in any presentation of $\prod_J Z$ there are infinitely many defining relations [8, p. 95, Ex. 6]. Thus Thm. 3.6 genuinely enlarges the class of modules E as described in Prop. 2.2 for which f is injective.

COROLLARY 3.7. If R is a Dedekind domain, $(E_j)_{j\in J}$ is an inverse system of flat R-modules such that f is injective for every E_j (e.g. E_j is projective), and $E = \lim E_j$ then f is injective.

Proof. By Lemma 3.4, $\lim_{i \to i} E_j$ is pure in $\prod E_j$. Now apply Thm. 3.6.

There is a characterization of modules E for which f is injective in terms of groups of homomorphisms. First a crucial result of D. Lazard.

THEOREM 3.8. (Lazard, [11].) A right R-module E is flat iff E is the direct limit of fg free modules.

DEFINITION 3.9. If R is commutative and E and F are R-modules we will call the module $\operatorname{Hom}_{R}(E, F)$ the F-dual of E.

The *R*-dual of *E* is commonly called, simply, "the dual of *E*". Henceforth we will omit the *R* when writing $\operatorname{Hom}_{R}(E, F)$.

THEOREM 3.10. If R is a Dedekind domain and E is a flat Rmodule then f is injective for E iff f is injective for all E-duals.

Proof. If f is injective for all E-duals then f is injective for the E-dual of R, Hom $(R, E) \approx E$.

Suppose, on the other hand, that f is injective for E. To show f is injective for all E-duals it suffices to consider the E-duals of flat modules since if T is the torsion submodule of an arbitrary module F then Hom $(F, E) \approx \text{Hom } (F/T, E)$. But if F is flat then $F \approx \lim F_j$ where F_j is fg and free, say of rank n_j . Therefore, Hom $(F, \overline{E}) \approx$ Hom $(\lim F_j, E) \approx \lim \text{Hom } (F_j, E)$ where Hom $(F_j, E) \approx E^{n_j}$. Since f is injective for Hom (F_j, E) (Prop. 2.1) and Hom (F_j, E) is flat, f is injective for $\lim \text{Hom } (F_j, E)$ (Cor. 3.7). Hence f is injective for Hom (F, E).

REMARK 1. Suppose R is a Dedekind domain, (E_j) is family of flat modules, E' is a pure submodule of the flat module E, and f is injective for E and every E_j . Then (a) f is injective for $\bigoplus E_j$ (Prop. 2.1), (b) f is injective for $\prod E_j$ (Thm. 3.6), and (c) f is injective for E' (Prop. 2.1). By Thm. 3.10 we also know (a') f is injective for all $\bigoplus E_j$ -duals, (b') f is injective for all $\prod E_j$ -duals, and (c') f is injective for all E'-duals. ((a) and (a') are actually special cases of (c) and (c') respectively.) It is perhaps worthwhile to note that (a'), (b'), and (c') can be proved directly by noting the following where $F \approx \lim_{i \to \infty} F_i$ is an arbitrary flat module, F_i is free and of finite rank n_i .

(i) Hom $(F, \bigoplus E_j)$ is pure in Hom $(F, \prod E_j)$.

(ii) f is injective for Hom $(F, \prod E_j) \approx \prod \text{Hom} (F, E_j) \approx \prod \varprojlim E_j^{n_2}$ by Cor. 3.7 and Thm. 3.6.

(iii) Hom $(F, E') \approx \lim_{t \to 0} (E')^{n_{\lambda}}$ is pure in Hom $(F, E) \approx \lim_{t \to 0} E^{n_{\lambda}}$ and f is injective for $\lim_{t \to 0} E^{n_{\lambda}}$ by Cor. 3.7.

REMARK 2. It is known that the Z-dual of a countable direct product of copies of Z is free [6]. Thus E = Hom(F, Z) when F is such a product does not enlarge beyond our previous knowledge our class of flat groups E for which f is injective.

In general when E is flat and f is injective for E it seems largely unknown what the *E*-duals are. (Hom $(\prod^{\infty} Z, Z) \approx \bigoplus^{\infty} Z$ is the only example we know.) Thus it is unclear how and if Thm. 3.10 does in fact enlarge the class of flat groups E for which f is injective. The identity of such *E*-duals seems a matter of general and independent interest. It is an easy consequence of [3, Ch. I, Ex. 8] that any right module over a right noetherian ring possesses a maximal injective submodule. The complement of this maximal injective submodule has no injective submodules. Therefore, over a Dedekind domain, an arbitrary module E can be written as a direct sum $E_1 \oplus E_2 \oplus E_3$ where E_1 is a direct sum of $R(P^{\infty})$'s, E_2 is a direct sum of quotient fields K, and E_3 has no injective submodules. Since injectivity \equiv divisibility over a Dedekind domain [9, Thm. 6], it is consistent with terminology in abelian groups to say E_3 is reduced. We claim that if f is injective for E then E is reduced, i.e., $E_1 \oplus E_2 = 0$. We need the following lemmas.

LEMMA 3.11. If R is a Dedekind domain, P is a prime ideal and $a \in R - P$ then R/P^n is a-divisible for every positive integer n.

Proof. Define
$$(P^n: a) = \{b \in R: ba \in P^n\}$$
. To show
 $g: R/P^n \longrightarrow R/P^n$

g. 10/1 · 10/1

given by $b + P^n \longrightarrow ab + P^n$ is bijective note that $(P^n: a) = P^n$. Thus g is injective. But R/P^n has a (finite) composition series. Hence g must be surjective.

LEMMA 3.12. If R is a Dedekind domain with prime ideal P, $F = \prod_{n=1}^{n} (R/P^n)$, and h is the canonical homomorphism $R \otimes F \to K \otimes$ F then h is not surjective.

Proof. Note that $K \otimes F$ is divisible. If im h were divisible then for an arbitrary $0 \neq p \in P$ there would exist $1 \otimes (b_n)_{n=1}^{\infty} \in R \otimes F$ such that $1 \otimes (1 - pb_n)_{n=1}^{\infty} \in \ker h$. But ker h is torsion [3, Ch. VII, Prop. 4.6]. (One can see this directly.) Thus there is $0 \neq t \in R$ such that $t(1 - pb_n) = 0$ for all n. Setting $x_n = 1 - pb'_n$ where $b'_n \in R$ and $b'_n + P^n = b_n$, we have $tx_n \in P^n$. Let m be the largest integer such that $t \in P^m$. If $t \notin P$, let m = 0. We claim $x_{m+1} \in P$.

If not there is $y \in R$ such that $x_{m+1}y - 1 = s \in P$. So $yx_{m+1}t = (1 + s)t = t + st \in P^{m+1}$. Thus $t \in P^{m+1}$. Contradiction.

Since $x_{m+1} = 1 - pb'_{m+1}$, $1 = x_{m+1} + pb'_{m+1} \in P$. Contradiction. So im h could not have been divisible.

THEOREM 3.13. If R is a Dedekind domain and f is injective for E then E must be reduced.

Proof. By Prop. 2.1 it suffices to show for a given prime ideal P there exists a family (F_i) such that f is not injective for K and f is not injective for $R(P^{\infty})$. We do this by choosing $F_n = R/P^n$, n =

1, 2, ..., and showing $K \otimes F_n = R(P^{\infty}) \otimes F_n = 0$ but $K \otimes \prod_{n=1}^{\infty} F_n \neq 0 \neq R(P^{\infty}) \otimes \prod_{n=1}^{\infty} F_n$.

Since $R(P^{\infty})$ and K are divisible, $R(P^{\infty}) \otimes F_n = K \otimes F_n = 0$.

Set $F = \prod_{n=1}^{\infty} F_n$ and denote by T the torsion submodule of F. We have the exact sequence

$$K \otimes F \longrightarrow K \otimes F/T \longrightarrow 0$$

where K and F/T are nonzero flat modules. Since $K \otimes F/T \neq 0$, $K \otimes F \neq 0$ by exactness.

Suppose Q is a prime different from P, m is a positive integer and $a \in Q^m - P$. Then F_n is a-divisible for every n (Lemma 3.11). So F is a-divisible. Hence $R(Q^{\infty}) \otimes F = 0$. Therefore, $R(P^{\infty}) \otimes F \approx K/R \otimes F$.

To show $K/R \otimes F \neq 0$ consider the exact sequence

$$R \otimes F \xrightarrow{h} K \otimes F \longrightarrow K/R \otimes F \longrightarrow 0$$

where h is the obvious map. Lemma 3.12 says h is not surjective. By exactness $K/R \otimes F \neq 0$.

REMARK. Over a Dedekind domain a reduced module that is not flat is either (1) a direct sum of a flat module and a finite direct sum of modules of type R/P^n where P is prime, or (2) has an infinite proper chain of direct summands each summand itself a finite direct sum of modules of type R/P^n [9, Thm. 9].

Since f is injective for any module of type R/P^n (Prop. 1.3), injectivity for modules of type (1) reduces to injectivity of reduced flat modules. Although we have some important examples of reduced flat modules for which f is injective, f need not be injective for such a module. This is shown by Prop. 3.15.

It is also true that f need not be injective for modules of type (2). This is shown by Prop. 3.14.

PROPOSITION 3.14. Suppose p is a prime integer and

$$E = \prod_{n=1}^{\infty} Z(p^n)$$
.

Then f is not injective for E. (Obviously E is of type (2).)

Proof. Choose a prime q different from p. Let $F = \prod_{n=1}^{\infty} Z(q^n)$ and denote by S and T the torsion subgroups of E and F respectively. Consider the exact sequence

$$E \otimes F \longrightarrow E/S \otimes F/T \longrightarrow 0$$
.

Since E/S and E/T are nonzero and flat, $E/S \otimes F/T \neq 0$. By exact-

ness, $E \otimes F \neq 0$.

To complete the proof note that $E \otimes Z(q^n) = 0$ for every integer n.

REMARK. It is easy to check that $\prod Z(p)$ where p ranges over the primes is another group of type (2) for which f is not injective.

PROPOSITION 3.15. Suppose E is a reduced torsion free abelian group that is divisible for a prime integer p. Then f is not injective for E.

Proof. Since E is p-divisible, $E \otimes Z(p^n) = 0$ for all positive n. Therefore $\prod_{n=1}^{\infty} (E \otimes Z(p^n)) = 0$.

If $F = \prod_{n=1}^{\infty} Z(p^n)$ and T is the torsion subgroup of F consider the exact sequence

 $E \otimes F \longrightarrow E \otimes F/T \longrightarrow 0$.

Since E and F/T are nonzero and torsion free $E \otimes F/T \neq 0$. By exactness, $E \otimes F \neq 0$.

Since
$$E \otimes \prod_{n=1}^{\infty} Z(p^n) \neq 0$$
 and $\prod_{n=1}^{\infty} (E \otimes Z(p^n)) = 0$,
 $f: E \otimes \prod_{n=1}^{\infty} Z(p^n) \longrightarrow \prod_{n=1}^{\infty} (E \otimes Z(p^n))$

is not injective.

REMARK 1. There are many examples of reduced torsion free groups that are divisible for a prime p. For a prime q different from p, the q-adic integers are such a group. The integers localized at q, Z_q , is another common example.

Concerning Z_q , if E is any torsion free group of rank 1 whose type is (k_1, k_2, \dots) , then E is reduced and divisible for p iff the k_n corresponding to p is ∞ but at least one other k_n is finite. (For the notion of type and a discussion of torsion free groups of rank 1 see [7].)

REMARK 2. We do not know if for every prime integer p, E is not *p*-divisible then f is injective. However, we have since shown this is true for rank 1 torsion free groups.

We have following interesting corollary to Thm. 3.13.

COROLLARY 3.16. If R is a Dedekind domain and (F_i) is a flat family such that $E \otimes F_i$ is reduced then $E \otimes \prod F_i$ is reduced.

Proof. $\prod (E \otimes F_i)$ is reduced and f is injective (Prop. 3.2). Therefore $E \otimes \prod F_i$ is reduced.

REMARK. A special case is when E is reduced. Then $E \otimes \prod R$ is reduced where the product is arbitrary

4. Commutative artinian rings. Matlis [12, Thm. 3.11] has shown that over a commutative artinian ring all indecomposable injective modules are fg. And we have seen that f is injective for all flat families (F_i) iff R is right noetherian and f is injective for all flat families (F_i) whenever E is indecomposable injective (Prop. 2.4). Therefore, by Prop. 1.3 we know the following

THEOREM 4.1. Over a commutative artinian ring f is injective for all flat families (F_i) .

REMARK 1. It is possible to prove this theorem by purely elementary methods, without resorting to Matlis' result.

REMARK 2. Thm. 4.1 endows commutative artinian rings with the property that was necessary to prove Thm. 3.6. (See Remark 1, following that theorem.) We have, therefore, the following

THEOREM 4.2. If R is commutative artinian, (E_j) is a family of flat modules, f is injective for E_j , and E is a pure submodule of $\prod E_j$, then f is injective for all families (F_i) .

REMARK. Thm. 4.2 can be derived in a different way by noting that over a commutative artinian ring every flat module is projective [1] and that the direct product of projective modules is projective [4].

We do not know that if R is commutative artinian then f is always injective. But if f does have this property R must be commutative artinian. To prove this we need two lemmas. A multiplicative subset of R will always be assumed to possess 1 and not 0. If $r \in R$ then (r) will denote the ideal generated by r.

LEMMA 4.3. Suppose S is a multiplicative subset of the commutative ring R. Then

$$S \cap (\bigcap_{s \in S} (s))
eq \oslash \ iff \ S^{-1}R \otimes \prod_{s \in S} R/(s) = 0 \; .$$

Therefore, if f is injective we have $S \cap (\bigcap_{s \in S} (s)) \neq \emptyset$ for all multiplicative subsets S of R.

Proof. For convenience we will write R/(s) as \overline{s} and omit the index set of products and intersections when this set is obviously S. Note that $S^{-1}R \otimes \prod \overline{s} \approx S^{-1}(\prod \overline{s})$, naturally.

If $t \in S \cap (\cap (s))$ and $x/u \in S^{-1}(\prod \overline{s})$ then x/u = tx/tu = 0/tu = 0. So $S \cap (\cap (s)) \neq \emptyset$ implies $S^{-1}R \otimes \prod \overline{s} = 0$.

If, on the other hand, $S^{-1}R \otimes \prod \overline{s} = 0$ then $(1 + (s))s \in S/1 = 0$ in $S^{-1}(\prod \overline{s})$. So there exists $t \in S$ such that $t \in \cap (s)$. Hence

$$S \cap (\cap (s)) \neq \emptyset$$
.

For the rest, note that for $s \in S$, $S^{-1}R \otimes \overline{s} = 0$. So $\prod (S^{-1}R \otimes \overline{s}) = 0$. Hence if f is injective, it must be true that $S^{-1}R \otimes \prod \overline{s} = 0$. Therefore, by the first part of the lemma, $S \cap (\cap (s)) \neq \emptyset$.

LEMMA 4.4. If R is a commutative ring such that for every prime ideal P, $(R - P) \cap (\bigcap_{s \notin P} (s)) \neq \emptyset$, then every prime ideal is maximal. In particular, if R is a domain then R is a field.

Proof. If $t \in (R - P) \cap (\bigcap_{s \notin P} (s))$ then $t + P \in \bigcap_{s \notin P} ((s) + P)/P$. So $t + P \in ((t^2) + P)/P$. Hence there exists $r \in R$ such that $t - rt^2 = t(1 - rt) \in P$. So $1 - rt \in P$. This says t + P is a unit in R/P.

If $s \in R - P$ then $t + P \in ((s) + P)/P$. Since t + P is a unit in R/P, (s) + P = R. Choose $b \in R$ such that $sb - 1 \in P$. Then (s + P)(b + P) = 1 + P, i.e., s + P is a unit in R/P. Hence P is maximal.

THEOREM 4.5. If R is a commutative ring and f is always injective then R is artinian.

Proof. By Lemma 4.3 $(R - P) \cap (\bigcap_{s \notin P} (s)) \neq \emptyset$ for all prime ideals *P*. Therefore, every prime ideal is maximal (Lemma 4.4). But *R* must be noetherian (Prop. 2.4). Since *R* is noetherian with every prime ideal maximal, it is a standard result that *R* must be artinian.

Since an artinian domain is necessarily a field we have the following immediate corollary.

COROLLARY 4.6. If R is a domain and f is always injective then R is a field.

As already remarked we do not know if the converse to Thm. 4.5 is true. There are, however, important classes of artinian rings over which f is injective. For example, suppose R is a proper quotient of a Dedekind domain. Then any R-module can be considered as a module, say E, over a Dedekind domain, this module having nontrivial annihilator. A classical result of Prüfer [10, Thm. 6] tells us that E is a direct sum of cyclic submodules. Since f is injective on each summand (Prop. 1.3), f is injective on all of E (Prop. 2.1).

A proper quotient of a Dedekind domain is a direct sum of rings each of whose lattices of ideals is finite and totally ordered by inclusion.

Any ring, not necessarily commutative, whose left and right free modules of rank 1 have unique composition series has been called a serial ring [5].

THEOREM 4.7. (Eisenbud and Griffith, [5].) If R is a serial ring then any right R-module is a direct sum of submodules with unique composition series.

In particular every module over a serial ring is a direct sum of finitely generated submodules. We have as an immediate consequence of this and Propositions 1.3 and 2.1, the following.

PROPOSITION 4.8. If R is a serial ring then f is always injective.

References

1. H. Bass, Finitistic homological dimension and a homological generalization of semiprimary rings, Trans. Amer. Math. Soc., **95** (1960), 466-488.

2. N. Bourbaki, Algèbre Commutative, Ch. 1 and 2, Hermann, Paris, 1961.

3. H. Cartan and S. Eilenberg, Homological Algebra, Princeton, 1956.

4. S. Chase, Direct products of modules, Trans. Amer. Math. Soc., 97 (1960), 457-473.

5. D. Eisenbud and P. Griffith, Serial rings, J. of Algebra, 17 (1971), 389-400.

6. E. Enochs, A note on reflexive modules, Pacific J. Math., 14 (1964), 879-881.

7. L. Fuchs, Abelian Groups, Pergamon Press, New York, 1960.

8. ____, Infinite Abelian Groups, Vol. I, Academic Press, New York, 1970.

9. I. Kaplansky, Modules over Dedekind rings and valuation rings, Trans. Amer. Math. Soc., 72 (1952), 327-340.

10. ____, Infinite Abelian Groups, U. of Michigan Press, Ann Arbor, 1969.

D. Lazard, Sur les modules plats, C. R. Acad. Sc. Paris, t. 258 (29 Juin 1964), 6313-16.
 E. Matlis, Injective modules over noetherian rings, Pacific J. Math., 8 (1958), 511-528.

13. R. Warfield, Jr., Purity and algebraic compactness for modules, Pacific J. Math., 28 (1969), 699-719.

Received August 21, 1972. This study is a doctoral dissertation prepared under the direction of Professor Edgar Enochs and submitted to the University of Kentucky. For the inspiration, guidance, and understanding by Professor Enochs throughout the development of this paper I am deeply grateful.

CHICAGO STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor) University of California

Los Angeles, California 90024

R. A. BEAUMONT

University of Washington Seattle, Washington 98105 J. DUGUNDJI*

Department of Mathematics University of Southern California Los Angeles, California 90007

D. GILBARG AND J. MILGRAM Stanford University Stanford, California 94305

K. YOSHIDA

ASSOCIATE EDITORS

E.F. BECKENBACH

B.H. NEUMANN

F. Wolf

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA UNIVERSITY OF SOUTHERN CALIFORNIA CALIFORNIA INSTITUTE OF TECHNOLOGY STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF NEVADA UNIVERSITY OF WASHINGTON NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY * * UNIVERSITY OF OREGON AMERICAN MATHEMATICAL SOCIETY **OSAKA UNIVERSITY** NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The Pacific Journal of Mathematics is issued monthly as of January 1966. Regular subscription rate: \$48.00 a year (6 Vols., 12 issues). Special rate: \$24.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by Pacific Journal of Mathematics All Rights Reserved

Pacific Journal of Mathematics Vol. 49, No. 2 June, 1973

Wm. R. Allaway, On finding the distribution function for an orthogonal polynomial	305
Frie Amar Sur un théorème de Moonen relatif aux fonctions analytiques bornées	311
Robert Morgan Brooks Analytic structure in the spectrum of a natural system	315
Robert Morgan Brooks, Analytic structure in the spectrum of a natural system	335
Kwang nan Chow and Mosas Clasper. Atoms on the Powlen houndary	330
Paul Frazier Duvall. Ir. and Jim Maxwell. <i>Tame</i> 7 ² actions on F ⁿ	339
Allen Pox Freedman. On the additivity theorem for n dimensional asymptotic	549
Anen Köy Meedman, On the additivity theorem for n-dimensional asymptotic density	357
John Griffin and Kelly Denis McKennon Multipliers and the aroun L -algebras	365
Charles Lemuel Hagonian Characterizations of λ connected plane continua	371
Ion Craig Helton, Rounds for products of interval functions	377
Jon Craig Tickon, Dounds for products of interval functions	301
Everett Lee Ledy Slender rings and modules	307
Shaza Matauura On the Lu Oi Kana conjecture and the Baroman representative	591
domains	407
Stephen H. McCleary. The lattice ordered group of automorphisms of an a-set	417
Stephen H. McCleary, $a = 2$ -transitive ordered normutation arouns	425
Stephen H. McCleary, a primitive ordered permutation groups	431
Dishord Dochborg, Almost isometries of Banach spaces and moduli of planar	431
domains	445
R F Rossa Radical properties involving one-sided ideals	467
Robert A Rubin On exact localization	473
S Sribala On Σ -inverse semigroups	483
H M (Hari Mohan) Srivastava On the Konhauser sets of hiorthogonal polynomials	105
suggested by the Laguerre polynomials	489
Stuart A. Steinberg, <i>Rings of quotients of rings without nilpotent elements</i>	493
Daniel Mullane Sunday The self-equivalences of an H-space	507
W I Thron and Richard Hawks Warren <i>On the lattice of proximities of Čech</i>	201
compatible with a given closure space	519
Frank Uhlig. The number of vectors jointly annihilated by two real quadratic forms	
determines the inertia of matrices in the associated pencil	537
Frank Uhlig, On the maximal number of linearly independent real vectors annihilated	
simultaneously by two real quadratic forms	543
Frank Uhlig, Definite and semidefinite matrices in a real symmetric matrix pencil	561
Arnold Lewis Villone, Self-adjoint extensions of symmetric differential operators	569
Cary Webb, Tensor and direct products	579
James Victor Whittaker, On normal subgroups of differentiable	
homeomorphisms	595
Jerome L. Paul, Addendum to: "Sequences of homeomorphisms which converge to	
homeomorphisms"	615
David E. Fields, <i>Correction to: "Dimension theory in power series rings"</i>	616
Peter Michael Curran, Correction to: "Cohomology of finitely presented groups"	617
Billy E. Rhoades, Correction to: "Commutants of some Hausdorff matrices"	617
Charles W. Trigg, Corrections to: "Versum sequences in the binary system"	619