

Pacific Journal of Mathematics

THE CENTER OF A SIMPLE ALGEBRA

TIMOTHY V. FOSSUM

THE CENTER OF A SIMPLE ALGEBRA

T. V. FOSSUM

The following theorem is proved: *Let A be a finite-dimensional simple K -algebra, K a field. If E is an extension of K and if M is an absolutely irreducible left $A \otimes_K E$ -module with character $\chi: A \otimes_K E \rightarrow E$, then $\chi(A)$ is a subfield of E which is K -isomorphic to the center of A .*

The purpose of this note is to give a short demonstration of the above theorem, proved first by Brauer [1] and later by Fein [3, 4] in case K is a perfect field (or, more generally, when A is a separable K -algebra). We make no assumptions on separability.

All algebras are assumed to be finite-dimensional, and all modules are unital left modules. Let B be a K -algebra, K a field, and let M be a B -module. We say M is *absolutely irreducible* if $M \otimes_K E$ is an irreducible $B \otimes_K E$ -module for all extensions E of K ; an extension E of K is said to be a *splitting field* for B if every irreducible $B \otimes_K E$ -module is absolutely irreducible (cf. [2, p. 202]). We will always identify B with its natural image in $B \otimes_K E$.

LEMMA. *Let A be a central simple L -algebra, L a field, and let F be a Galois extension of L . If N is an irreducible $A \otimes_L F$ -module with character $\chi: A \otimes_L F \rightarrow F$ such that $\chi \neq 0$, then $\chi(A) = L$.*

Proof. For each $\sigma \in G(F/L)$, the Galois group of F over L , define an L -automorphism (still denoted by σ) of $A \otimes_L F$ by $\sigma(\sum_i a_i \otimes f_i) = \sum_i a_i \otimes \sigma(f_i)$. Each such L -automorphism of $A \otimes_L F$ gives rise to an irreducible $A \otimes_L F$ -module σN : The additive group of $\sigma N = \{\sigma n: n \in N\}$ is the same as that of N , but the module structure on σN is defined by $(\sigma x)(\sigma n) = \sigma(xn)$ for all $x \in A \otimes_L F$ and $n \in N$. One checks that the character of σN is $\sigma\chi\sigma^{-1}$. Since $A \otimes_L F$ is simple [2, (68.1)], $\sigma N \cong N$, and so $\sigma\chi\sigma^{-1} = \chi$. This says that for each $a \in A$, $\sigma\chi(a) = \chi(a)$ for all $\sigma \in G(F/L)$; hence $\chi(A) \subseteq L$. Since $\chi(A)$ is a nonzero L -subspace of L , it follows that $\chi(A) = L$, as desired.

LEMMA. *Let A be a central simple L -algebra, L a field, and let E be an extension of L . If M is an absolutely irreducible $A \otimes_L E$ -module with character $\zeta: A \otimes_L E \rightarrow E$, then $\zeta(A) = L$.*

Proof. It is well known that there is a Galois extension F of L which is a splitting field for A . Let N be an irreducible $A \otimes_L F$ -module with character $\chi: A \otimes_L F \rightarrow F$. Then N is absolutely irredu-

cible, and $\chi \neq 0$. By the above lemma, $\chi(A) = L$.

Let W be a compositum of E and F . Now $(A \otimes_L E) \otimes_E W$ and $(A \otimes_L F) \otimes_F W$ are both isomorphic to $A \otimes_L W$, and $M \otimes_E W$ and $N \otimes_F W$ are irreducible $A \otimes_L W$ -modules with characters ζ and χ , respectively, on A . Since $A \otimes_L W$ is simple, $M \otimes_E W \cong N \otimes_F W$, so $\zeta = \chi$ on A . It follows that $\zeta(A) = \chi(A) = L$, as desired.

Observe that the restriction ζ_A of ζ to A is the reduced trace of A into its center L .

THEOREM. *Let A be a simple K -algebra with center L . Let E be an extension of K , and let M be an absolutely irreducible $A \otimes_K E$ -module with character $\chi: A \otimes_K E \rightarrow E$. Then $\chi(A)$ is a K -subfield of E , and $\chi(A) \cong L$ as K -algebras.*

Proof. Since L is contained in the center of $A \otimes_K E$, L is K -isomorphic to a subfield of $\text{End}_{A \otimes_K E}(M) \cong E$, and we regard this as an identification [2, (29.13)]. It follows that M can be made into an $A \otimes_L E$ -module, and that the diagram

$$\begin{array}{ccc}
 A & \longrightarrow & A \otimes_L E \\
 \parallel & & \uparrow \\
 A & \longrightarrow & A \otimes_K E
 \end{array}
 \begin{array}{l}
 \searrow \alpha \\
 \nearrow \beta
 \end{array}
 \begin{array}{c}
 \\
 \text{End}_E(M) \\
 \longrightarrow T \\
 E
 \end{array}$$

commutes, where T is the trace map, and where α and β are the E -algebra homomorphisms afforded by the module structures on M . Since M is an absolutely irreducible $A \otimes_K E$ -module, it follows that β is an epimorphism, and so α is also an epimorphism. Thus M is an absolutely irreducible $A \otimes_L E$ -module, with character $T\alpha: A \otimes_L E \rightarrow E$. By the previous lemma, $T\alpha(A) = L$. Now $\alpha(A) = \beta(A)$, so $\chi(A) = T\beta(A) = T\alpha(A) = L$, as desired.

With a little extra effort, it is possible to generalize this result to orders. In particular, let R be a Krull domain with quotient field K , and let A be a simple K -algebra. An R -order \mathcal{A} in A is a unital R -subalgebra of A which spans A over K , and each element of \mathcal{A} is integral over R . Let E be an extension of K , and let M be an absolutely irreducible $A \otimes_K E$ -module with character $\chi: A \otimes_K E \rightarrow E$. If \mathcal{A} is an R -order in A which is separable over its center, then one can prove that $\chi(\mathcal{A})$ is R -isomorphic to the center of \mathcal{A} .

REFERENCES

1. R. Brauer, *Über systeme hypercomplexer Zahlen*, Math. Zeit., **30** (1929), 79-107.
2. C. W. Curtis and I. Reiner, *Representation Theory of Finite Groups and Associative Algebras*, Interscience, New York, 1962.

3. B. Fein, *Representations of direct products of finite groups*, Pacific J. Math., **20** (1967), 45-58.
4. ———, *The Schur index for projective representations of finite groups*, Pacific J. Math., **28** (1969), 87-100.

Received September 28, 1972.

THE UNIVERSITY OF UTAH

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Pacific Journal of Mathematics

Vol. 50, No. 1

September, 1974

Gail Atneosen, <i>Sierpinski curves in finite 2-complexes</i>	1
Bruce Alan Barnes, <i>Representations of B^*-algebras on Banach spaces</i>	7
George Benke, <i>On the hypergroup structure of central $\Lambda(p)$ sets</i>	19
Carlos R. Borges, <i>Absolute extensor spaces: a correction and an answer</i>	29
Tim G. Brook, <i>Local limits and tripleability</i>	31
Philip Throop Church and James Timourian, <i>Real analytic open maps</i>	37
Timothy V. Fossum, <i>The center of a simple algebra</i>	43
Richard Freiman, <i>Homeomorphisms of long circles without periodic points</i>	47
B. E. Fullbright, <i>Intersectional properties of certain families of compact convex sets</i>	57
Harvey Charles Greenwald, <i>Lipschitz spaces on the surface of the unit sphere in Euclidean n-space</i>	63
Herbert Paul Halpern, <i>Open projections and Borel structures for C^*-algebras</i>	81
Frederic Timothy Howard, <i>The numer of multinomial coefficients divisible by a fixed power of a prime</i>	99
Lawrence Stanislaus Husch, Jr. and Ping-Fun Lam, <i>Homeomorphisms of manifolds with zero-dimensional sets of nonwandering points</i>	109
Joseph Edmund Kist, <i>Two characterizations of commutative Baer rings</i>	125
Lynn McLinden, <i>An extension of Fenchel's duality theorem to saddle functions and dual minimax problems</i>	135
Leo Sario and Cecilia Wang, <i>Counterexamples in the biharmonic classification of Riemannian 2-manifolds</i>	159
Saharon Shelah, <i>The Hanf number of omitting complete types</i>	163
Richard Staum, <i>The algebra of bounded continuous functions into a nonarchimedean field</i>	169
James DeWitt Stein, <i>Some aspects of automatic continuity</i>	187
Tommy Kay Teague, <i>On the Engel margin</i>	205
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable, V</i>	215
Kung-Wei Yang, <i>Isomorphisms of group extensions</i>	299