

Pacific Journal of Mathematics

**THE NUMER OF MULTINOMIAL COEFFICIENTS DIVISIBLE
BY A FIXED POWER OF A PRIME**

FREDERIC TIMOTHY HOWARD

THE NUMBER OF MULTINOMIAL COEFFICIENTS DIVISIBLE BY A FIXED POWER OF A PRIME

F. T. Howard

In this paper some results of L. Carlitz and the writer concerning the number of binomial coefficients divisible by p^j but not by p^{j+1} are generalized to multinomial coefficients. In particular $\theta_j(k; n)$ is defined to be the number of multinomial coefficients $n!/n_1! \cdots n_k!$ divisible by exactly p^j , and formulas are found for $\theta_j(k; n)$ for certain values of j and n . Also the generating function technique used by Carlitz for binomial coefficients is generalized to multinomial coefficients.

1. Introduction. Let p be a fixed prime and let n and j be nonnegative integers. L. Carlitz [2], [3] has defined $\theta_j(n)$ as the number of binomial coefficients

$$\binom{n}{r} \quad (r = 0, 1, \dots, n)$$

divisible by exactly p^j and he has found formulas for $\theta_j(n)$ for certain values of j and n . In particular, if we write

$$(1.1) \quad n = a_0 + a_1p + \cdots + a_s p^s \quad (0 \leq a_i < p)$$

then

$$\theta_0(n) = (a_0 + 1)(a_1 + 1) \cdots (a_s + 1)$$

$$\theta_1(n) = \sum_{i=0}^{s-1} (a_0 + 1) \cdots (a_{i-1} + 1)(p - a_i - 1)a_{i+1}(a_{i+2} + 1) \cdots (a_s + 1).$$

The writer [5], [6] has also considered the problem of evaluating $\theta_j(n)$.

The purpose of this paper is to consider the analogous problem for multinomial coefficients and to generalize some of the formulas developed by Carlitz and the writer. Thus we define $\theta_j(k; n)$ as the number of multinomial coefficients

$$(n_1, \dots, n_k) = \frac{n!}{n_1! \cdots n_k!} (n_1 + \cdots + n_k = n)$$

divisible by exactly p^j . In this definition the order of the terms n_1, \dots, n_k is important. We are distinguishing, for example, between $(1, 2, 3)$ and $(2, 1, 3)$. Clearly $\theta_j(2; n) = \theta_j(n)$.

In this paper we find formulas for $\theta_0(k; n)$, $\theta_1(k; n)$, and $\theta_2(k; n)$. We also show how the generating function method used by Carlitz

can be generalized to multinomial coefficients, and we evaluate $\theta_j(k; n)$ for special values of j and n .

Throughout this paper we assume p is a fixed prime number and k is a fixed positive integer, $k > 1$.

2. Preliminaries. Let $E(n_1, \dots, n_k)$ denote the largest value of w such that p^w divides (n_1, \dots, n_k) . To determine $E(n_1, \dots, n_k)$ we shall make use of an analogue [4] of Kummer's famous theorem for binomial coefficients:

LEMMA 2.1. *Let n have expansion (1.1), let $n = n_1 + \dots + n_k$ and let*

$$(2.1) \quad n_i = a_{i,0} + a_{i,1}p + \dots + 2_{i,s}p^s \quad (0 \leq a_{i,r} < p)$$

for $i = 1, \dots, k$. If

$$\begin{aligned} a_{1,0} + \dots + a_{k,0} &= \varepsilon_0 p + a_0 \\ \varepsilon_0 + a_{1,1} + \dots + a_{k,1} &= \varepsilon_1 p + a_1 \\ &\dots\dots\dots \\ \varepsilon_{s-1} + a_{1,s} + \dots + a_{k,s} &= a_s \end{aligned}$$

where each $\varepsilon_i = 0, 1, \dots$, or $k - 1$, then

$$E(n_1, \dots, n_k) = \varepsilon_0 + \varepsilon_1 + \dots + \varepsilon_{s-1}.$$

If n has expansion (1.1) and if $\nu(n)$ is the largest value of w such that p^w divides $n!$, then it is familiar [1, p. 55] that

$$\nu(n) = \frac{n - S(n)}{p - 1}$$

where $S(n) = a_0 + a_1 + \dots + a_s$. Thus we have

LEMMA 2.2. *If $n = n_1 + \dots + n_k$ then*

$$E(n_1, \dots, n_k) = \frac{S(n_1) + \dots + S(n_k) - S(n)}{p - 1}.$$

Furthermore, if $E_t(n_1, \dots, n_k)$ is the largest value of w such that p^w divides

$$(n + t) \cdots (n + 1) \quad (n_1, \dots, n_k),$$

then

$$E_t(n_1, \dots, n_k) = \frac{S(n_1) + \dots + S(n_k) - S(n + t) + t}{p - 1}.$$

Compositions, or ordered partitions, are important in evaluating $\theta_j(k; n)$. We define a composition of a nonnegative integer u into r parts to be an ordered sequence of r nonnegative integers whose sum is u . This is more general than the usual definition of composition in that we allow 0 to be one or more of the parts. See [7, pp. 124-125] for example.

Throughout this paper we shall let $C(u)$ denote the number of compositions of u into exactly k parts, with no part larger than $p - 1$. We define $C(u) = 0$ if $u < 0$.

LEMMA 2.3. $C(u)$ is the coefficient of x^u in the expansion of

$$(1 + x + x^2 + \dots + x^{p-1})^k = \left[\sum_{i=0}^{\infty} \binom{k+i-1}{i} x^i \right] (1 - x^p)^k.$$

It is clear from Lemma 2.3 that if $0 \leq a < p$ and if $0 \leq b$, then

$$(2.2) \quad C(a + bp) = \sum_{i=0}^b (-1)^i \binom{k}{i} \binom{k-1+a+(b-i)p}{k-1}.$$

In particular, for $0 \leq a < p$,

$$C(a) = \binom{k-1+a}{k-1},$$

$$C(a + p) = \binom{k-1+a+p}{k-1} - k \binom{k-1+a}{k-1},$$

$$C(a + 2p) = \binom{k-1+a+2p}{k-1} - k \binom{k-1+a+p}{k-1} + \binom{k}{2} \binom{k-1+a}{k-1}.$$

3. Evaluation of $\theta_0(k; n)$, $\theta_1(k; n)$, $\theta_2(k; n)$.

THEOREM 3.1. If n has expansion (1.1) then

$$\theta_0(k; n) = C(a_0)C(a_1) \dots C(a_s).$$

Proof. We use Lemma 2.1. If $E(n_1, \dots, n_k) = 0$ then we must have

$$\sum_{i=1}^k a_{i,r} = a_r \quad (r = 0, \dots, s).$$

For a given r , the total number of ways we can have this equality is equal to $C(a_r)$.

Note that by Lemma 2.3 we have

$$C(a_r) = \binom{a_r + k - 1}{k - 1} \quad (r = 0, \dots, s).$$

THEOREM 3.2. *If n has expansion (1.1) then*

$$\theta_1(k; n) = \sum_{\hat{i}=0}^{s-1} C(a_0) \cdots C(a_{\hat{i}-1})C(a_{\hat{i}} + p)C(a_{\hat{i}+1} - 1)C(a_{\hat{i}+2}) \cdots C(a_s).$$

Proof. Using Lemma 2.1, we see that if $E(n_1, \dots, n_k) = 1$ then we must have exactly one $\varepsilon_i = 1$, $0 \leq i < s$. So for some i we have

$$\begin{aligned} a_{1,i} + \cdots + a_{k,i} &= a_i + p, \\ a_{1,i+1} + \cdots + a_{k,i+1} &= a_{i+1} - 1, \\ a_{1,r} + \cdots + a_{k,r} &= a_r \quad (r \neq i, i+1). \end{aligned}$$

Clearly the total number of ways we can have these equalities is

$$C(a_0) \cdots C(a_{i-1})C(a_i + p)C(a_{i+1} - 1)C(a_{i+2}) \cdots C(a_s).$$

To simplify the formula for $\theta_2(k; n)$ we introduce the following notation. Let

$$\begin{aligned} A_i &= \left[\prod_{t=0}^s C(a_t) \right] / \left[C(a_i)C(a_{i+1})C(a_{i+2}) \right], \\ B_i &= \left[\prod_{t=0}^s C(a_t) \right] / \left[C(a_i)C(a_{i+1}) \right], \\ H_{i,r} &= \left[\prod_{t=0}^s C(a_t) \right] / \left[C(a_i)C(a_{i+1})C(a_r)C(a_{r+1}) \right]. \end{aligned}$$

THEOREM 3.3. *If n has expansion (1.1) then*

$$\begin{aligned} \theta_2(k; n) &= \sum_{\hat{i}=0}^{s-2} C(p + a_i)C(p + a_{i+1} - 1)C(a_{i+2} - 1)A_i \\ &\quad + \sum_{\hat{i}=0}^{s-1} C(2p + a_i)C(a_{i+1} - 2)B_i \\ &\quad + \sum_{r=\hat{i}+2}^{s-1} \sum_{\hat{i}=0}^{s-3} C(p + a_i)C(a_{i+1} - 1)C(p + a_r)C(a_{r+1} - 1)H_{i,r}. \end{aligned}$$

Proof. The proof is similar to the proof of Theorem 3.2. We determine the number of ways we can have exactly two of the ε 's equal to 1 or exactly one ε equal to 2, and all other ε 's equal to 0.

For example, let $p = 5$, $k = 3$, and $n = 278 = 3 + 5^2 + 2 \cdot 5^3$. We have

$$\begin{aligned} \theta_0(3; 278) &= C(3)C(0)C(1)C(2) = 180; \\ \theta_1(3; 278) &= C(3)C(5)C(0)C(2) + C(3)C(0)C(6)C(1) = 1650, \\ \theta_2(3; 278) &= C(8)C(4)C(0)C(2) + C(3)C(5)C(5)C(1) \\ &\quad + C(3)C(0)C(11)C(0) = 11,100. \end{aligned}$$

In each example we have used (2.2) to evaluate $C(u)$.

4. Generating functions for $\theta_i(k; n)$. Let $\psi_{r,t,j}(k; n)$ denote the

number of products $(n + t) \cdots (n + 1)(n_1, \dots, n_k)$, $n_1 + \cdots + n_k = n$, divisible by exactly p^j . Clearly

$$(4.1) \quad \psi_{t,j}(k; n) = \theta_{j-r}(k; n)$$

if p^r is the highest power of p dividing $(n + t) \cdots (n + 1)$.

Also

$$\psi_{t,j}(k; n) = 0$$

if p^{j+1} divides $(n + t) \cdots (n + 1)$. We introduce the following generating functions:

$$F_0(x, y) = \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} \theta_j(k; n) x^n y^j,$$

$$F_t(x, y) = \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} \psi_{t,j}(k; n) x^n y^j \quad (t > 0).$$

Using an argument analogous to that of Carlitz [3], we obtain

$$(4.2) \quad F_0(x, y) = \sum_{t=0}^m y^t f_t(x) F_t(x^p, y)$$

where m is the integer such that

$$(4.3) \quad mp \leq k(p - 1) < (m + 1)p$$

and

$$f_t(x) = \sum_{a=tp}^{tp+p-1} C(a) x^a \quad (0 \leq t < m),$$

$$f_m(x) = \sum_{a=mp}^{kp-k} C(a) x^a.$$

Comparing coefficients of $x^n y^j$ on both sides of (4.2), we have, for $0 \leq a < p$,

$$(4.4) \quad \theta_j(k; a + bp) = C(a) \theta_j(k; b) + \sum_{t=1}^m C(a + tp) \psi_{t,j-t}(k; b - t).$$

In (4.4) it is understood that $\psi_{t,j}(k; u) = 0$ if $u < 0$ and $\psi_{t,-1}(k; u) = 0$.

Also, for $t < p$,

$$F_t(x, y) = \sum_{r=1}^h y^r g_r(x) F_r(x^p, y)$$

where h is the integer such that

$$(4.5) \quad hp - t \leq k(p - 1) < (h + 1)p - t,$$

and

$$\begin{aligned}
 g_0(x) &= \sum_{a=0}^{p-t-1} C(a)x^a, \\
 g_r(x) &= \sum_{a=rp-t}^{(r+1)p-t-1} C(a)x^a \quad (r = 1, \dots, h-1), \\
 g_h(x) &= \sum_{a=hp-t}^{kp-k} C(a)x^a.
 \end{aligned}$$

Thus for $0 \leq a < p-t$, $hp+a \leq kp-k$, we have

$$(4.6) \quad \psi_{t,j}(k; a+bp) = C(a)\theta_j(k; b) + \sum_{r=1}^h C(a+rp)\psi_{r,j-r}(k; b-r).$$

For $0 \leq a < p-t$, $hp+a > kp-k$, we have

$$(4.7) \quad \psi_{t,j}(k; a+bp) = C(a)\theta_j(k; b) + \sum_{r=1}^{h-1} C(a+rp)\psi_{r,j-r}(k; b-r).$$

For $p-t \leq a < p$, we have

$$(4.8) \quad \psi_{t,j}(k; a+bp) = \sum_{r=1}^h C(a+(r-1)p)\psi_{r,j-r}(k; a-r+1).$$

Here again it is understood that $\psi_{r,j}(k; u) = 0$ if $u < 0$. We remark that in all of these formulas specific values for $C(u)$ can be found from formula (2.2).

Using (4.4) we can compute $\theta_j(k; n)$ for special values of n . By (4.4) and (4.1) we have, for $0 \leq a < p$, $0 \leq b < p$,

$$\begin{aligned}
 \theta_j(k; a+bp) &= C(a+jp)\theta_0(k; b-j) \\
 &= C(a+jp) C(b-j) \quad \text{if } j \leq m, \\
 &= 0 \quad \text{if } j > m
 \end{aligned}$$

where m is defined by (4.3).

Also, if $0 \leq a < p$,

$$\begin{aligned}
 \theta_j(k; a+p^2) &= C(a)C(1) \quad \text{if } j = 0, \\
 &= C(a+(j-1)p)C(p-j+1) \quad \text{if } 1 \leq j \leq m+1, \\
 &= 0 \quad \text{if } j > m+1.
 \end{aligned}$$

If $0 \leq a < p$, $p > 2$,

$$\begin{aligned}
 &\theta_j(k; a+2p^2) \\
 &= C(a+(j-2)p)\theta_1(k; 2p-j+2) \\
 &\quad + C(a+(j-1)p)\theta_0(k; 2p-j+1) \quad (1 < j \leq p+1, j \leq m+1), \\
 &= C(a+(j-2)p)\theta_1(k; 2p-j+2) \quad (j = m+2 \leq p+1), \\
 &= C(a+(j-2)p)\theta_1(k; p) \quad (j = p+2 \leq m+2), \\
 &= C(a+(j-2)p)\theta_0(k; p-r+2) \quad (j = p+r \leq m+2, 2 < r \leq p+2), \\
 &= 0 \quad \text{if } j > m+2.
 \end{aligned}$$

If $0 \leq a < p$, $0 \leq b < p$,

$$\begin{aligned}
 \theta_j(k; a + bp + p^2) &= C(a + (j - 1)p)\theta_1(k; p + b - j + 1) \\
 &\quad + C(a + jp)\theta_0(k; p + b - j) \quad (b \geq j; m + 1 > j) \\
 &= C(a + (j - 1)p)\theta_0(k; p + b - j + 1) \quad (b < j \leq p + b, m + 1 > j), \\
 &= C(a + mp)\theta_1(k; p + b - m) \quad (j = m + 1, b \geq m), \\
 &= C(a + mp)\theta_0(k; p + b - m) \quad (j = m + 1, b < m), \\
 &= 0 \quad \text{if } j > m + 1.
 \end{aligned}$$

Some of the results in [2] can also be generalized. We use the symbols $E(n_1, \dots, n_k)$ and $E_t(n_1, \dots, n_k)$ as they are used in Lemma 2.2.

Let

$$\begin{aligned}
 F_j(n; x_1, \dots, x_k) &= \sum_{\substack{a_1 + \dots + a_k = n \\ E(a_1, \dots, a_k) = j}} x_1^{a_1} \dots x_k^{a_k}, \\
 G_{t,j}(n; x_1, \dots, x_k) &= \sum_{\substack{a_1 + \dots + a_k = n \\ E_t(a_1, \dots, a_k) = j}} x_1^{a_1} \dots x_k^{a_k} \quad (t > 0), \\
 G_{0,j}(n; x_1, \dots, x_k) &= F_j(n; x_1, \dots, x_k).
 \end{aligned}$$

Note that

$$\begin{aligned}
 F_j(n; x, \dots, x) &= x^n \theta_j(k; n), \\
 G_{t,j}(n; x, \dots, x) &= x^{n\psi_{t,j}(k; n)}.
 \end{aligned}$$

By generalizing Carlitz's work in [2] in the natural way, we obtain

$$\begin{aligned}
 (4.9) \quad &F_j(a + bp; x_1, \dots, x_k) \\
 &= \sum_{s=0}^m c_{sp+a}(x_1, \dots, x_k) G_{s,j-s}(b - s; x_1^p, \dots, x_k^p)
 \end{aligned}$$

where $0 \leq a < p$, m is defined by (4.3), and

$$c_r(x_1, \dots, x_k) = \sum_{s_1 + \dots + s_k = r} x_1^{s_1} \dots x_k^{s_k}.$$

Also, if h is defined by (4.5),

$$\begin{aligned}
 (4.10) \quad &G_{t,j}(a + bp; x_1, \dots, x_k) \\
 &= \sum_{s=0}^h c_{sp+a}(x_1, \dots, x_k) G_{s,j-s}(b - s; x_1^p, \dots, x_k^p) \\
 &\quad (hp + a \leq kp - k, 0 \leq a < p - t), \\
 &= \sum_{s=0}^{h-1} c_{sp+a}(x_1, \dots, x_k) G_{s,j-s}(b - s; x_1^p, \dots, x_k^p) \\
 &\quad (hp + a > kp - k, 0 \leq a < p - t), \\
 &= \sum_{s=1}^h c_{(s-1)p+a}(x_1, \dots, x_k) G_{s,j-s}(a - s + 1; x_1^p, \dots, x_k^p) \\
 &\quad (p - t \leq a < p - 1).
 \end{aligned}$$

5. Some special evaluations. If $j > \nu(n)$, where $\nu(n)$ is the exponent of the highest power of p that divides $n!$, then it is clear that $\theta_j(k; n) = 0$. For example, if $0 \leq a < p$, $0 \leq b < p$ then

$$\theta_j(k; a + bp) = 0 \quad (j > b).$$

Let n have expansion (1.1). By Lemma 2.1 it is clear that $\theta_j(k; n) = 0$ for $j > M$, where

$$\begin{aligned} M &= s(k - 1) \quad \text{if } k \leq a_s + 1 \\ &= (s - 1)(k - 1) + a_s \quad \text{if } k > a_s + 1. \end{aligned}$$

Also,

$$\begin{aligned} \theta_M(k; n) &= C(a_0 + (k - 1)p)C(a_s - k + 1) \prod_{i=1}^{s-1} C(a_i - k + 1 + (k - 1)p) \\ &\hspace{20em} (k \leq a_s + 1), \\ &= C(a_0 + (k - 1)p)C(a_{s-1} - k + 1 + a_s p) \prod_{i=1}^{s-2} C(a_i - k + 1 + (k - 1)p) \\ &\hspace{15em} (k > a_s + 1, s > 1), \\ &= C(a_0 + a_1 p) \hspace{15em} (k > a_s + 1, s = 1). \end{aligned}$$

For example, if $k = 2$ and $a_s \neq 0$ then $M = s$. This is the case for ordinary binomial coefficients. We have in this case

$$\theta_s(2; n) = (p - a_0 - 1)(p - a_1) \cdots (p - a_{s-1})a_s.$$

For $p = 2$ we can generalize the method used in [6]. Let

$$(5.1) \quad n = 2^{e_1} + \cdots + 2^{e_r}, \quad 0 \leq e_1 < \cdots < e_r,$$

$$(5.2) \quad n_i = 2^{e_{i,1}} + \cdots + 2^{e_{i,S(i)}}, \quad 0 \leq e_{i,1} < \cdots < e_{i,S(i)}.$$

Consider all the different compositions $n = n_1 + \cdots + n_k$ such that (5.1) and (5.2) hold, such that

$$S(n_1) + \cdots + S(n_k) = r + j,$$

and such that there are a total of $r + j - t$ $e_{i,w}$'s having the property that $e_{i,w} \neq e_{x,y}$ for all x, y (except for the one case $i = x, w = y$). Let $b_{j,t}$ be the sum over all these compositions of the number of different ways of distributing the remaining t $e_{i,w}$'s into k distinct cells with no two identical objects in the same cell. Then for $p = 2, j > 0$,

$$(5.3) \quad \theta_j(k; n) = b_{j,2}k^{m+j-2} + b_{j,3}k^{m+j-3} + \cdots + b_{j,m+j}.$$

Using the convention that $e_1 - e_0 = t$ means $e_1 = t - 1$ and that $e_1 - e_0 > t$ means $e_1 > t - 1$, let

$$\begin{aligned}
e_i - e_{i-1} &> 1 \quad \text{for } q_1 \text{ terms } e_i, \\
&> 2 \quad \text{for } q_2 \text{ terms } e_i, \\
&= 1, e_{i-1} - e_{i-2} = 1 \quad \text{for } q_3 \text{ terms } e_i \quad (i \neq 2), \\
&= 1, e_{i-1} - e_{i-2} > 1 \quad \text{for } q_4 \text{ terms } e_i, \\
&= 2 \quad \text{for } q_5 \text{ terms } e_i \quad (i \neq 1), \\
&= 1 \quad \text{for } q_6 \text{ terms } e_i \quad (i \neq 1).
\end{aligned}$$

Then, by (5.3), for $p = 2$,

$$\theta_0(k; n) = k^r,$$

$$\theta_1(k; n) = q_1 \binom{k}{2} k^{r-1} + q_6 \binom{k}{3} k^{r-2},$$

$$\begin{aligned}
\theta_2(k; n) &= q_2 \binom{k}{2} k^r + q_5 \binom{k}{3} k^{r-1} \\
&\quad + \left[\binom{q_1}{2} + q_4 \right] \binom{k}{2}^2 k^{r-2} \\
&\quad + \left[q_4(q_1 - 1) + q_3 \right] \binom{k}{3} \binom{k}{2} k^{r-3} \\
&\quad + \left[\binom{q_4}{2} + q_5(q_3 - 1) + q_4 q_3 \right] \binom{k}{3}^2 k^{r-4}.
\end{aligned}$$

For example, let $n = 2^4 + 2^5 + 2^{20} + 2^{26} + 2^{28}$. Then $q_1 = 4$, $q_2 = 3$, $q_3 = 0$, $q_4 = 1$, $q_5 = 1$ and $q_6 = 1$. Thus

$$\theta_0(k; n) = k^5$$

$$\theta_1(k; n) = 4 \binom{k}{2} k^4 + \binom{k}{3} k^3,$$

$$\theta_2(k; n) = 3 \binom{k}{2} k^5 + \binom{k}{3} k^4 + 7 \binom{k}{2}^2 k^3 + 3 \binom{k}{3} \binom{k}{2} k^3.$$

REFERENCES

1. P. Bachmann, *Niedere Zahlentheorie*, vol. 1, Leipzig, 1902.
2. L. Carlitz, *Distribution of binomial coefficients*, Riv. Mat. Univ. Parma, (2) **11** (1970), 45-64.
3. ———, *The number of binomial coefficients divisible by a fixed power of a prime*, Rend. Circ. Mat. Palermo, (2) **16** (1967), 299-320.
4. Robert D. Fray, *Congruence properties of ordinary and q-binomial coefficients*, Duke Math. J., **34** (1967), 467-480.
5. F. T. Howard, *Formulas for the number of binomial coefficients divisible by a fixed power of a prime*, Proc. Amer. Math. Soc., **37** (1973), 358-362.

6. F. T. Howard, *The number of binomial coefficients divisible by a fixed power of 2*, Proc. Amer. Math. Soc., **29** (1971), 236-242.
7. J. Riordan, *An Introduction to Combinatorial Analysis*, New York, 1958.

Received September 11, 1972.

WAKE FOREST UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Gail Atneosen, <i>Sierpinski curves in finite 2-complexes</i>	1
Bruce Alan Barnes, <i>Representations of B^*-algebras on Banach spaces</i>	7
George Benke, <i>On the hypergroup structure of central $\Lambda(p)$ sets</i>	19
Carlos R. Borges, <i>Absolute extensor spaces: a correction and an answer</i>	29
Tim G. Brook, <i>Local limits and tripleability</i>	31
Philip Throop Church and James Timourian, <i>Real analytic open maps</i>	37
Timothy V. Fossum, <i>The center of a simple algebra</i>	43
Richard Freiman, <i>Homeomorphisms of long circles without periodic points</i>	47
B. E. Fullbright, <i>Intersectional properties of certain families of compact convex sets</i>	57
Harvey Charles Greenwald, <i>Lipschitz spaces on the surface of the unit sphere in Euclidean n-space</i>	63
Herbert Paul Halpern, <i>Open projections and Borel structures for C^*-algebras</i>	81
Frederic Timothy Howard, <i>The numer of multinomial coefficients divisible by a fixed power of a prime</i>	99
Lawrence Stanislaus Husch, Jr. and Ping-Fun Lam, <i>Homeomorphisms of manifolds with zero-dimensional sets of nonwandering points</i>	109
Joseph Edmund Kist, <i>Two characterizations of commutative Baer rings</i>	125
Lynn McLinden, <i>An extension of Fenchel's duality theorem to saddle functions and dual minimax problems</i>	135
Leo Sario and Cecilia Wang, <i>Counterexamples in the biharmonic classification of Riemannian 2-manifolds</i>	159
Saharon Shelah, <i>The Hanf number of omitting complete types</i>	163
Richard Staum, <i>The algebra of bounded continuous functions into a nonarchimedean field</i>	169
James DeWitt Stein, <i>Some aspects of automatic continuity</i>	187
Tommy Kay Teague, <i>On the Engel margin</i>	205
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable, V</i>	215
Kung-Wei Yang, <i>Isomorphisms of group extensions</i>	299