

Pacific Journal of Mathematics

ISOMORPHISMS OF GROUP EXTENSIONS

KUNG-WEI YANG

ISOMORPHISMS OF GROUP EXTENSIONS

KUNG-WEI YANG

To my parents

Let $0 \rightarrow G \rightarrow E \rightarrow \Pi \rightarrow 1$ and $0 \rightarrow G \rightarrow E' \rightarrow \Pi \rightarrow 1$ be two crossed product extensions given by the crossed product groups $E = [G, \varphi, f, \Pi]$ and $E' = [G, \varphi', f', \Pi]$ respectively. A homomorphism $\Gamma: E \rightarrow E'$ is stabilizing if the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & G & \longrightarrow & E & \longrightarrow & \Pi \longrightarrow 1 \\ & & \parallel & & \downarrow \Gamma & & \parallel \\ 0 & \longrightarrow & G & \longrightarrow & E' & \longrightarrow & \Pi \longrightarrow 1 \end{array}$$

commutes. In this paper, a necessary and sufficient condition for the existence of a stabilizing homomorphism (hence isomorphism) between any two crossed product extensions is obtained.

The result is applied to obtain a necessary and sufficient condition for the existence of an automorphism $\phi: E \rightarrow E$ making the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & G & \longrightarrow & E & \longrightarrow & \Pi \longrightarrow 1 \\ & & \downarrow \tau & & \downarrow \phi & & \downarrow \sigma \\ 0 & \longrightarrow & G & \longrightarrow & E & \longrightarrow & \Pi \longrightarrow 1 \end{array}$$

commutative, given $(\sigma, \tau) \in \text{Aut } \Pi \times \text{Aut } G$.

NOTATION. In general, we use the notation in [3]. Throughout the paper, G and Π denote two fixed groups. G will be written in additive notation and Π in multiplicative notation. $\text{Aut } G$, $\text{Out } G$, and ZG are the automorphism group, the outer automorphism group, and the center of G , respectively. For any element $a \in G$, $\mu(a)$ denotes the inner automorphism $\mu(a)(g) = a + g - a$ given by conjugation with a . When X is a group the natural image of an element $x \in X$ in a quotient group of X is denoted \bar{x} . When φ is a map, $\bar{\varphi}$ denotes the map $\bar{\varphi}(x) = \overline{\varphi(x)}$.

Given groups G, Π , and functions $\varphi: \Pi \rightarrow \text{Aut } G, f: \Pi \times \Pi \rightarrow G$ satisfying

$$(1) \quad \varphi(x)f(y, z) + f(x, yz) = f(x, y) + f(xy, z),$$

$$(2) \quad \varphi(x)\varphi(y) = \mu[f(x, y)]\varphi(xy),$$

and the normalization conditions $\varphi(1) = 1, f(x, 1) = 0 = f(1, y)$, the set $G \times \Pi$ under the sum defined by

$$(3) \quad (g, x) + (h, y) = (g + \varphi(x)h + f(x, y), xy)$$

is a group. The group so constructed is called a *crossed product group*, and is denoted $[G, \varphi, f, \Pi]$, or simply E . With the homomorphism $G \rightarrow E$ defined by $g \mapsto (g, 1)$, and $E \rightarrow \Pi$ defined by $(g, x) \mapsto x$, we have an extension of G by Π

$$0 \longrightarrow G \longrightarrow E \longrightarrow \Pi \longrightarrow 1 .$$

The extension is called a *crossed product extension*.

Results. Let $E = [G, \varphi, f, \Pi]$ and $E' = [G, \varphi', f', \Pi]$ be two crossed product groups. Define a stabilizing homomorphism $\Gamma: E \rightarrow E'$ as in the abstract. Notice that, by the "5 lemma" for groups, a stabilizing homomorphism is an isomorphism. Clearly, a homomorphism $\Gamma: E \rightarrow E'$ is stabilizing if and only if Γ is of the form

$$(4) \quad \Gamma(g, x) = (g + \gamma(x), x) ,$$

and

$$(5) \quad \varphi(x)g + f(x, y) + \gamma(xy) = \gamma(x) + \varphi'(x)[g + \gamma(y)] + f'(x, y) .$$

Because of the normalization conditions, $\gamma(1) = 0$. Setting $y = 1$ in (5), we obtain

$$(6) \quad \varphi(x)g + \gamma(x) = \gamma(x) + \varphi'(x)g .$$

Setting $g = 0$ in (5), we obtain

$$(7) \quad f(x, y) + \gamma(xy) = \gamma(x) + \varphi'(x)\gamma(y) + f'(x, y) .$$

Conversely, (5) can immediately be derived from (6) and (7). Summarizing, we have

LEMMA. *If $E = [G, \varphi, f, \Pi]$ and $E' = [G, \varphi', f', \Pi]$ are two crossed product groups, then $\Gamma: E \rightarrow E'$ is a stabilizing isomorphism if and only if Γ is of the form (4), where the map $\gamma: \Pi \rightarrow G$ satisfies (6) and (7).*

In particular, when $\varphi = \varphi'$, and $f = f'$, we see that by (6), $\gamma(x) \in ZG$, and by (7), $\gamma \in Z^1(\Pi, ZG)$. $Z^1(\Pi, ZG)$ is the group of normalized 1-cocycles, and the Π -module structure on ZG is given by φ .

COROLLARY. *If $E = [G, \varphi, f, \Pi]$ is a crossed product group, then $\Gamma: E \rightarrow E$ is a stabilizing automorphism if and only if Γ is of the form (4), where $\gamma \in Z^1(\Pi, ZG)$.*

We remark that both the lemma and the corollary are well-known. See for instance, [5, p. 127], [2, 17.1 Satz, p. 119], [4].

It is obvious from (6) that, as homomorphisms from Π to $\text{Out } G$,

$$(8) \quad \bar{\varphi} = \bar{\varphi}' .$$

Trivially, (7) implies that

$$(9) \quad k(x, y) = -f(x, y) + \gamma(x) + \varphi'(x)\gamma(y) + f'(x, y) - \gamma(xy)$$

is equal to 0.

Conversely, given crossed product groups $E = [G, \varphi, f, \Pi]$ and $E' = [G, \varphi', f', \Pi]$, we can ascertain the existence of a stabilizing isomorphism from E to E' by the following procedure. First, we decide whether condition (8) is fulfilled. If not, the question is settled. If (8) is satisfied, then there is a function $\gamma: \Pi \rightarrow G$ such that (6) holds, and ZG acquires a well-defined Π -module structure with operators $xc = \varphi(x)c (= \varphi'(x)c)$, for $c \in ZG$. We set $\gamma(1) = 0$. It is now meaningful to speak of the group $Z^2(\Pi, ZG)$ (resp., $B^2(\Pi, ZG)$) of the 2-dimensional normalized cocycles (resp., coboundaries) of Π with values in ZG . Define $k(x, y)$ by (9). We claim that $k(x, y) \in Z^2(\Pi, ZG)$. Trivially, $k(x, 1) = 0 = k(1, y)$. To see $k(x, y) \in ZG$, we merely observe that conjugating $\varphi'(xy)g$ with $\gamma(xy)$ and conjugating $\varphi'(xy)g$ with $-f(x, y) + \gamma(x) + \varphi'(x)\gamma(y) + f'(x, y)$ give the same result $\varphi'(xy)g$. $k(x, y)$, being the difference of these two elements, is therefore in ZG . To verify the identity

$$xk(y, z) - k(xy, z) + k(x, yz) - k(x, y) = 0 ,$$

we observe that

$$\begin{aligned} k(x, yz) - k(xy, z) &= -f(x, yz) + \gamma(x) + \varphi'(x)\gamma(yz) + f'(x, yz) - f'(xy, z) \\ &\quad - \varphi'(xy)\gamma(z) - \gamma(xy) + f(xy, z) \\ &= -f(x, yz) + \gamma(x) + \varphi'(x)[\gamma(yz) - f'(y, z) - \varphi'(y)\gamma(z)] \\ &\quad + f'(x, y) - \gamma(xy) + f(xy, z) \\ &= -f(x, yz) + \gamma(x) - \varphi'(x)k(y, z) - \varphi'(x)f(y, z) - \gamma(x) \\ &\quad + f(x, y) + k(x, y) + f(xy, z) \\ &= k(x, y) - xk(y, z). \end{aligned}$$

We made use of the identities (1), (2), (9), (6), and (1), in that order.

Finally, if $k(x, y) \in B^2(\Pi, ZG)$, then $k(x, y) = x\beta(y) - \beta(xy) + \beta(x)$. The function $\gamma': \Pi \rightarrow G$ defined by $\gamma'(x) = \gamma(x) - \beta(x)$ satisfies (6) and (7). Therefore, the map Γ defined by (4), using γ' instead of γ , is a stabilizing isomorphism. If $k(x, y) \notin B^2(\Pi, ZG)$, then there could not exist a stabilizing isomorphism $\Gamma: E \rightarrow E'$. For, if there were to exist such an isomorphism, the discussion leading up to the above lemma would show that $\Gamma(g, x) = (g + \gamma'(x), x)$, with γ' satisfying (6) and (7).

Since γ and γ' both satisfy (6), $\beta(x) = \gamma(x) - \gamma'(x) \in ZG$. By (7) we have $k(x, y) = x\beta(y) - \beta(xy) + \beta(x)$ showing $k(x, y) \in B^2(\Pi, ZG)$. This discussion also shows that $\overline{k(x, y)}$ in $H^2(\Pi, ZG)$ is independent of the choice of γ . These results may now be summarized as follows.

THEOREM 1. *Let $E = [G, \varphi, f, \Pi]$ and $E' = [G', \varphi', f', \Pi]$ be two crossed product groups. Then there exists a stabilizing isomorphism $\Gamma: E \rightarrow E'$, if and only if*

- (A) $\overline{\varphi} = \overline{\varphi}'$, and
- (B) $\overline{k} = 0$ in $H^2(\Pi, ZG)$,

where $k(x, y)$ is defined as above.

We note that Theorem 1 is well-known (and is easily seen to be true) in the case where $\varphi = \varphi'$ [3, Theorem 8.8, p. 128, Lemma 8.2].

An application. Let $0 \rightarrow G \rightarrow E \rightarrow \Pi \rightarrow 1$ be a group extension. Call an automorphism of E taking G onto G an automorphism *over* G . Clearly, any automorphism of E over G induces automorphisms τ on G and σ on Π . It is easy to see that, in general, not every pair $(\sigma, \tau) \in \text{Aut } \Pi \times \text{Aut } G$ can be so induced by an automorphism of E over G . In [4], Charles Wells defined an exact sequence which gives a necessary and sufficient condition for a pair $(\sigma, \tau) \in \text{Aut } \Pi \times \text{Aut } G$ to be inducible by an automorphism of E over G . We now apply Theorem 1 to prove a similar result. We hope our method will also help clarify the nature of the map $C \rightarrow H^2_\alpha(\Pi, ZG)$ as defined in [4].

Let $0 \rightarrow G \rightarrow E \rightarrow \Pi \rightarrow 1$ be a group extension. We may (and do) assume that E is of the form $E = [G, \varphi, f, \Pi]$ with homomorphisms $G \rightarrow E, E \rightarrow \Pi$ of the form as defined in the definition of a crossed product extension at the beginning of this paper. We say that a pair $(\sigma, \tau) \in \text{Aut } \Pi \times \text{Aut } G$ is *inducible* if there exists an automorphism $\phi: E \rightarrow E$ such that the diagram

$$(10) \quad \begin{array}{ccccccc} 0 & \longrightarrow & G & \longrightarrow & E & \longrightarrow & \Pi \longrightarrow 1 \\ & & & & \downarrow \tau & & \downarrow \sigma \\ & & & & G & \longrightarrow & E \longrightarrow \Pi \longrightarrow 1 \end{array}$$

is commutative.

For $(\sigma, \tau) \in \text{Aut } \Pi \times \text{Aut } G$, let $\varphi_\sigma(x) = \varphi(\sigma x), f_\sigma(x, y) = f(\sigma x, \sigma y); \varphi_\tau(x) = \tau\varphi(x)\tau^{-1}, f_\tau(x, y) = \tau f(x, y)$. If $\overline{\varphi}_\sigma = \overline{\varphi}_\tau$, there exists a map $\gamma: \Pi \rightarrow G$ such that $\varphi_\tau(x) + \gamma(x) = \gamma(x) + \varphi_\sigma(x)$. Choose γ so that $\gamma(1) = 0$. In this case, define

$$(11) \quad k_{\sigma, \tau}(x, y) = -f_\tau(x, y) + \gamma(x) + \varphi_\sigma(x)\gamma(y) + f_\sigma(x, y) - \gamma(xy).$$

THEOREM 2. *The pair $(\sigma, \tau) \in \text{Aut } \Pi \times \text{Aut } G$ is inducible if and*

only if

- (A) $\bar{\varphi}_\sigma = \bar{\varphi}_\tau$, and
- (B) $\bar{k}_{\sigma,\tau} = 0$ in $H^2(\Pi, ZG)$,

where $k_{\sigma,\tau}(x, y)$ is defined as in (11), and the Π -module structure on ZG is induced by the homomorphism $\bar{\varphi}_\sigma = \bar{\varphi}_\tau$.

Proof. Let $0 \rightarrow G \rightarrow E \rightarrow \Pi \rightarrow 1$ be a group extension. Set $E = [G, \varphi, f, \Pi]$. Let $\varphi_\sigma, f_\sigma, \varphi_\tau, f_\tau, k_{\sigma,\tau}$ be as defined in the paragraph preceding Theorem 2. Let $E_\tau = [G, \varphi_\tau, f_\tau, \Pi]$, $E_\sigma = [G, \varphi_\sigma, f_\sigma, \Pi]$. Define $T: E \rightarrow E_\tau$ by $T(g, x) = (\tau g, x)$, and $\Sigma: E_\sigma \rightarrow E$ by $\Sigma(g, x) = (g, \sigma x)$. It is a straightforward matter to check that T and Σ are both group homomorphisms and that the following two diagrams

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & G & \longrightarrow & E & \longrightarrow & \Pi & \longrightarrow & 1 \\
 & & \downarrow \tau & & \downarrow T & & \parallel & & \\
 0 & \longrightarrow & G & \longrightarrow & E_\tau & \longrightarrow & \Pi & \longrightarrow & 1 \\
 & & & & & & & & \\
 0 & \longrightarrow & G & \longrightarrow & E_\sigma & \longrightarrow & \Pi & \longrightarrow & 1 \\
 & & \parallel & & \downarrow \Sigma & & \downarrow \sigma & & \\
 0 & \longrightarrow & G & \longrightarrow & E & \longrightarrow & \Pi & \longrightarrow & 1
 \end{array}$$

commute.

If there is a stabilizing isomorphism $\Gamma: E_\tau \rightarrow E_\sigma$, then (σ, τ) is clearly inducible.

Conversely, if (σ, τ) is inducible, then there exists an automorphism $\Phi: E \rightarrow E$ such that the diagram (10) is commutative. Such an automorphism is necessarily of the form $\Phi(g, x) = (\tau g + \gamma(x), \sigma x)$, where γ satisfies (6) and (7) with φ replaced by φ_τ and φ' replaced by φ_σ . Therefore, $\Gamma = \Sigma^{-1}\Phi T^{-1}$ is a stabilizing isomorphism from E_τ to E_σ . By Theorem 1, (σ, τ) is inducible if and only if (A) and (B) are satisfied.

Condition (A) of Theorem 2 can be stated more explicitly as follows: For any $x \in \Pi$, $\bar{\tau}\bar{\varphi}(x)\bar{\tau}^{-1} = \bar{\varphi}(\sigma x)$.

As direct consequences of Theorem 2, we have

COROLLARY 1. *If $\text{Out } G = 1$ and $H^2(\Pi, ZG) = 0$, then every pair $(\sigma, \tau) \in \text{Aut } \Pi \times \text{Aut } G$ is inducible.*

COROLLARY 2. *If $\text{Out } G = 1$ and $ZG = 0$, then for any group E such that $G \triangleleft E$ and for each $\tau \in \text{Aut } G$, there exists $\Phi \in \text{Aut } E$ such that restriction of Φ to G is equal to τ .*

The second corollary also follows directly from [1, Theorem 1].

REFERENCES

1. R. Baer, *Absolute retracts in group theory*, Bull. Amer. Math. Soc., **52** (1946), 501-6.
2. B. Huppert, *Endliche Gruppen I, Die Grundlehren der math. Wissenschaften, Band 134*, Springer-Verlag, Berlin and New York, 1967.
3. S. MacLane, *Homology, Die Grundlehren der math. Wissenschaften, Band 114*, Academic Press, New York, Springer Verlag, Berlin, Grottingen, Heidelberg, 1963.
4. C. Wells, *Automorphisms of group extensions*, Trans. Amer. Math. Soc., **155** (1970), 189-194.
5. H. J. Zassenhaus, *The Theory of Groups*, 2nd ed., Chelsea Publishing Company, 1958.

Received September 28, 1972.

WESTERN MICHIGAN UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Pacific Journal of Mathematics

Vol. 50, No. 1

September, 1974

Gail Atneosen, <i>Sierpinski curves in finite 2-complexes</i>	1
Bruce Alan Barnes, <i>Representations of B^*-algebras on Banach spaces</i>	7
George Benke, <i>On the hypergroup structure of central $\Lambda(p)$ sets</i>	19
Carlos R. Borges, <i>Absolute extensor spaces: a correction and an answer</i>	29
Tim G. Brook, <i>Local limits and tripleability</i>	31
Philip Throop Church and James Timourian, <i>Real analytic open maps</i>	37
Timothy V. Fossum, <i>The center of a simple algebra</i>	43
Richard Freiman, <i>Homeomorphisms of long circles without periodic points</i>	47
B. E. Fullbright, <i>Intersectional properties of certain families of compact convex sets</i>	57
Harvey Charles Greenwald, <i>Lipschitz spaces on the surface of the unit sphere in Euclidean n-space</i>	63
Herbert Paul Halpern, <i>Open projections and Borel structures for C^*-algebras</i>	81
Frederic Timothy Howard, <i>The numer of multinomial coefficients divisible by a fixed power of a prime</i>	99
Lawrence Stanislaus Husch, Jr. and Ping-Fun Lam, <i>Homeomorphisms of manifolds with zero-dimensional sets of nonwandering points</i>	109
Joseph Edmund Kist, <i>Two characterizations of commutative Baer rings</i>	125
Lynn McLinden, <i>An extension of Fenchel's duality theorem to saddle functions and dual minimax problems</i>	135
Leo Sario and Cecilia Wang, <i>Counterexamples in the biharmonic classification of Riemannian 2-manifolds</i>	159
Saharon Shelah, <i>The Hanf number of omitting complete types</i>	163
Richard Staum, <i>The algebra of bounded continuous functions into a nonarchimedean field</i>	169
James DeWitt Stein, <i>Some aspects of automatic continuity</i>	187
Tommy Kay Teague, <i>On the Engel margin</i>	205
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable, V</i>	215
Kung-Wei Yang, <i>Isomorphisms of group extensions</i>	299