

Pacific Journal of Mathematics

ON A “SQUARE” FUNCTIONAL EQUATION

ARJUN K. GUPTA

ON A "SQUARE" FUNCTIONAL EQUATION

A. K. GUPTA

In a recent paper Stanton and Cowan have generalized the Pascal's triangle to a tableau. They have developed several expressions for these numbers, using combinatorial techniques. In the present paper we derive some of their results very simply, by using the calculus of finite differences. We further obtain the relations of these numbers to hypergeometric function and derive many relations among these numbers which are useful in constructing the tableau.

1. Introduction. The triangular array of binomial coefficients, well-known as Pascal's triangle, has been much studied. Basically, it depends on the recursion relation

$$(1) \quad f(n + 1, r) = f(n, r) + f(n, r - 1).$$

In a recent paper Stanton and Cowan [5], have considered a generalization of this situation by defining a tableau by the recurrence relation

$$(2) \quad g(n + 1, r + 1) = g(n, r + 1) + g(n + 1, r) + g(n, r).$$

This formula, together with the boundary conditions, $g(n, 0) = g(0, r) = 1$, uniquely determines $g(n, r)$. The lower half of the first portion of this tableau is presented in Table 1, the upper half can be obtained by symmetry in n and r (see § 2).

TABLE 1

g(n, r)							
1							
1	3						
1	5	13					
1	7	25	63.				
1	9	41	129	321			
1	11	61	231	681	1683		
1	13	85	377	1289	3653	8989	
1	15	113	575	2241	7183	19825	48639

Stanton and Cowan [5], have developed several expressions for these numbers $g(n, r)$, and indicated that they have a further com-

binatorial interpretation, namely, $g(n, r)$ is the volume of an r -sphere in n -space under the Lee metric. In this paper we obtain some of the results of Stanton and Cowan by using the calculus of finite differences and obtain some additional properties of the function $g(n, r)$.

2. Main results. In order to obtain an explicit formula for $g(n, r)$ the calculus of finite differences, [4], is efficacious. Let E denote the shift operator such that $EU(r) = U(r + 1)$, and Δ denote the difference operator defined by $\Delta U(r) = U(r + 1) - U(r) = (E - 1)U(r)$. We use the convention, [1], that the binomial coefficient $\binom{n}{r}$ is defined only for r an integer, and that $\binom{n}{r}$ vanishes for $r < 0$, and for $r > n$.

With this notation we prove the following results.

LEMMA 1.

$$(3) \quad g(n, r) = \sum_{\alpha} \binom{n}{\alpha} \binom{r + \alpha}{n} = \sum_{\alpha} \binom{r + \alpha}{\alpha} \binom{r}{n - \alpha}.$$

Proof. From (2), we get

$$(4) \quad \Delta g(n + 1, r) = (E + 1)g(n, r),$$

which yields,

$$\begin{aligned} g(n, r) &= (E + 1)^n \Delta^{-n} g(0, r) = \sum_{\alpha} \binom{n}{\alpha} E^{\alpha} \Delta^{-n} g(0, r) \\ &= \sum_{\alpha} \binom{n}{\alpha} E^{\alpha} \binom{r}{n}, \quad \text{since } \Delta^j \binom{r}{j} = 1 \text{ (see (3))}, \\ &= \sum_{\alpha} \binom{n}{\alpha} \binom{r + \alpha}{n} = \sum_{\alpha} \binom{r + \alpha}{\alpha} \binom{r}{n - \alpha}. \end{aligned}$$

LEMMA 2.

$$(5) \quad g(n, r) = \sum_{\alpha} 2^{\alpha} \binom{n}{\alpha} \binom{r}{\alpha}.$$

Proof. From (4), we have

$$g(n + 1, r) = \left(\frac{E + 1}{E - 1} \right) g(n, r),$$

which can be written as,

$$g(n, r) = (1 + 2\Delta^{-1})^n g(0, r) \\ = \sum_{\alpha} \binom{n}{\alpha} 2^{\alpha} \Delta^{-\alpha} g(0, r) = \sum_{\alpha} 2^{\alpha} \binom{n}{\alpha} \binom{r}{\alpha}.$$

It is obvious that $g(n, r)$ is symmetric in n and r . The effective upper limit of the summations in (5) is at $K = \min. (n, r)$.

In view of the fact that the numbers $g(n, r)$ possess some applications, in addition to their intrinsic interest, it may be of interest to discuss some additional properties, and their relation to some well-known functions.

LEMMA 3.

$$(6) \quad g(n, r + s) = \sum_k g(k, r) \{g(n - k, s) - g(n - k - 1, s)\}.$$

Proof. It is known, [5], that $g(n, r)$ is the coefficient of x^n in the expansion of $(1 + x)^n / (1 - x)^{n+1}$. Or, since $g(n, r) = g(r, n)$, we might use the coefficient of x^n in $(1 + x)^r / (1 - x)^{r+1}$. Let $f_r(x) = (1 + x)^r / (1 - x)^{r+1}$. Then it is easy to verify that

$$(7) \quad (1 - x)f_r(x)f_s(x) = f_{r+s}(x).$$

Now equate the coefficients of x^n in the expansion of (7) and we get the desired result.

If we consider the identity (7) as

$$(8) \quad f_r(x)f_s(x) = (1 - x)^{-1}f_{r+s}(x)$$

and equate the coefficients of x^n , we get

$$\sum_{k=0}^n g(k, r + s) = \sum_{k=0}^n g(k, r)g(n - k, s).$$

In the following lemma we prove that $g(n, r)$ is a special case of the hypergeometric function ${}_2F_1$ defined by

$${}_2F_1(a, b; c; x) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k} \frac{x^k}{k!},$$

where,

$$(a)_k = a(a + 1) \cdots (a + k - 1).$$

LEMMA 4.

$$(9) \quad g(n, r) = {}_2F_1(-n, -r; 1; 2).$$

Proof.

$$\begin{aligned}
 g(n, r) &= \sum_{\alpha} 2^{\alpha} \binom{n}{\alpha} \binom{r}{\alpha} = \sum_{\alpha} \frac{2^{\alpha} n! r!}{\alpha! \alpha! (n - \alpha)! (r - \alpha)!} \\
 &= \sum_{\alpha} \frac{2^{\alpha}}{\alpha!} \frac{(-n)_{\alpha} (-r)_{\alpha}}{(1)_{\alpha}} = {}_2F_1(-n, -r; 1; 2).
 \end{aligned}$$

Note that the diagonal sums become, $g(n, n) = {}_2F_1(-n, -n; 1; 2)$. In the case of binomial coefficients such sums provide the Fibonacci numbers.

LEMMA 5.

$$(10) \quad g(n, r) = \frac{1}{n - r} \{n g(n - 1, r) - r g(n, r - 1)\}, \quad n \neq r.$$

Proof. We have,

$$\begin{aligned}
 n g(n - 1, r) - r g(n, r - 1) &= n \sum_{\alpha} 2^{\alpha} \binom{n - 1}{\alpha} \binom{r}{\alpha} - r \sum_{\alpha} 2^{\alpha} \binom{n}{\alpha} \binom{r - 1}{\alpha} \\
 &= \sum_{\alpha} 2^{\alpha} \binom{n}{\alpha} \binom{r}{\alpha} (n - r) = (n - r) g(n, r).
 \end{aligned}$$

For actual computations of $g(n, r)$ the result (10) is as easy as the basic definition (2).

The result of Lemma 5, can also be obtained from the corresponding identity for hypergeometric functions, $(b - a) {}_2F_1(a, b; c; x) + a {}_2F_1(a + 1, b; c; x) - b {}_2F_1(a, b + 1; c; x) = 0$ where now $a = -n$, $b = -r$, $c = -1$, and $x = 2$.

LEMMA 6.

$$(11) \quad g(n, r) = \frac{2r + 1}{n} g(n - 1, r) + \frac{n - 1}{n} g(n - 2, r)$$

$$(12) \quad = \frac{2n + 1}{r} g(n, r - 1) + \frac{r - 1}{r} g(n, r - 2).$$

Proof. The result (11) follows from the following identity for hypergeometric functions,

$$\begin{aligned}
 (c - a) {}_2F_1(a - 1, b; c; x) &+ (2a - c - ax + bx) {}_2F_1(a, b; c; x) \\
 &+ a(x - 1) {}_2F_1(a + 1, b; c; x) = 0
 \end{aligned}$$

where now $a = -n$, $b = -r$, $c = 1$, and $x = 2$. Similarly the result (12) can be obtained from the symmetry or from the corresponding identity for hypergeometric functions, viz,

$$\begin{aligned}
 (c - b) {}_2F_1(a, b - 1; c; x) &+ (2b - c - bx + ax) {}_2F_1(a, b; c; x) \\
 &+ b(x - 1) {}_2F_1(a, b + 1; c; x) = 0.
 \end{aligned}$$

Lemma 6 is useful to compute the numbers $g(n, r)$ in any one column or row of the tableau and for this purpose has the advantage over the formulae (2) or (10) of Lemma 5.

Some more relations between the numbers $g(n, r)$, which can be easily obtained from the corresponding identities for hypergeometric functions, are the following.

$$(n + r + 1)g(n, r) + ng(n - 1, r) - (r + 1)g(n, r + 1) = 0 ,$$

or equivalently,

$$(n + r + 1)g(n, r) + rg(n, r - 1) - (n + 1)g(n + 1, r) = 0 ,$$

and,

$$(n + 1)g(n + 1, r) - (r - n)g(n, r) - (r + 1)g(n, r + 1) = 0 .$$

3. Additional remarks. It may be noted that the combinatorial interpretation of $g(n, r)$; the value $\sum_{\alpha} 2^{\alpha} \binom{n}{\alpha} \binom{r}{\alpha}$, is the volume (i.e., the number of lattice points in Euclidean n -space within a regular cross-polytope) of a sphere of radius r in n -dimensions (or a sphere of radius n in r -dimensions) using the Lee metric as deduced by Golomb and Welch [3]. Golomb [2], has also derived the generating function for $g(n, r)$,

$$\frac{1}{1 - x - y - xy} = \sum_{n=0}^{\infty} \sum_{r=0}^{\infty} g(n, r) x^n y^r$$

which may be used to evaluate $g(n, r)$ either explicitly or asymptotically and is simpler than our results. Applications of these numbers in sphere packing, coding metrics, and chess puzzles are also described by Golomb [2]. Thus the numbers $g(n, r)$ have many applications in addition to their intrinsic interest. The relationship with the hypergeometric functions further illustrates their usefulness.

Thanks are due to the referee for pointing out the reference [2].

REFERENCES

1. William Feller, *An Introduction to Probability Theory and Its Applications*, Second, ed., Vol. I, John Wiley and Sons, Inc. New York, 1957.
2. S. W. Golomb, *Sphere Packing, Coding Metrics, and Chess Puzzles*, Proceedings of the Second Chapel Hill Conference on Combinatorial Mathematics and its Applications, University of North Carolina, 1970, 176-189.
3. S. W. Golomb and L. R. Welch, *Algebraic Coding and the Lee Metric*, Error Correcting Codes, (edited by H. B. Mann), John Wiley and Sons, Inc.; New York, 1968, 175-194.
4. L. M. Milne-Thomson, *The Calculus of Finite Differences*, Macmillan and Co., Ltd., London, 1960.

5. R. G. Stanton and D. D. Cowan, *Note on a 'square' functional equation*, Siam Review, **12** (1970), 277-279.

Received October 5, 1972.

THE UNIVERSITY OF MICHIGAN

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$60.00 a year (6 Vols., 12 issues). Special rate: \$30.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Mustafa Agah Akcoglu, John Philip Huneke and Hermann Rost, <i>A counter example to the Blum Hanson theorem in general spaces</i>	305
Huzihiro Araki, <i>Some properties of modular conjugation operator of von Neumann algebras and a non-commutative Radon-Nikodym theorem with a chain rule</i>	309
E. F. Beckenbach, Fook H. Eng and Richard Edward Tafel, <i>Global properties of rational and logarithmico-rational minimal surfaces</i>	355
David W. Boyd, <i>A new class of infinite sphere packings</i>	383
K. G. Choo, <i>Whitehead Groups of twisted free associative algebras</i>	399
Charles Kam-Tai Chui and Milton N. Parnes, <i>Limit sets of power series outside the circles of convergence</i>	403
Allan Clark and John Harwood Ewing, <i>The realization of polynomial algebras as cohomology rings</i>	425
Dennis Garbanati, <i>Classes of circulants over the p-adic and rational integers</i>	435
Arjun K. Gupta, <i>On a "square" functional equation</i>	449
David James Hallenbeck and Thomas Harold MacGregor, <i>Subordination and extreme-point theory</i>	455
Douglas Harris, <i>The local compactness of vX</i>	469
William Emery Haver, <i>Monotone mappings of a two-disk onto itself which fix the disk's boundary can be canonically approximated by homeomorphisms</i>	477
Norman Peter Herzberg, <i>On a problem of Hurwitz</i>	485
Chin-Shui Hsu, <i>A class of Abelian groups closed under direct limits and subgroups formation</i>	495
Bjarni Jónsson and Thomas Paul Whaley, <i>Congruence relations and multiplicity types of algebras</i>	505
Lowell Duane Loveland, <i>Vertically countable spheres and their wild sets</i>	521
Nimrod Megiddo, <i>Kernels of compound games with simple components</i>	531
Russell L. Merris, <i>An identity for matrix functions</i>	557
E. O. Milton, <i>Fourier transforms of odd and even tempered distributions</i>	563
Dix Hayes Pettey, <i>One-one-mappings onto locally connected generalized continua</i>	573
Mark Bernard Ramras, <i>Orders with finite global dimension</i>	583
Doron Ravdin, <i>Various types of local homogeneity</i>	589
George Michael Reed, <i>On metrizable complete Moore spaces</i>	595
Charles Small, <i>Normal bases for quadratic extensions</i>	601
Philip C. Tonne, <i>Polynomials and Hausdorff matrices</i>	613
Robert Earl Weber, <i>The range of a derivation and ideals</i>	617