Pacific Journal of Mathematics

THE LOCAL COMPACTNESS OF vX

DOUGLAS HARRIS

Vol. 50, No. 2

October 1974

THE LOCAL COMPACTNESS OF vX

DOUGLAS HARRIS

Necessary and sufficient conditions are given for the local compactness of the Hewitt realcompactification vX of a completely regular Hausdorff space X; the conditions are expressed in terms of the space X alone. In addition, the local compactness of other extensions is considered.

Introduction. There has been much recent interest in determining conditions on a completely regular Hausdorff space X that are equivalent to the local compactness of its Hewitt realcompactification vX. This interest stems primarily from the fact that the seemingly artificial hypothesis "vX is locally compact" enters quite naturally into the examination of the relation $vX \times vY = v(X \times Y)$. Apparently the only known condition equivalent to the local compactness of vX is one discussed by Comfort in [1] and [2]. As remarked by Comfort, the condition is not on X alone, but involves vX essentially in its statement.

In the present paper a condition on X is given which is equivalent to the local compactness of vX (Theorem 2.7) and a number of known results are obtained as corollaries of this characterization theorem. Another characterization (Theorem 2.3) is given of the local compactness of vX in terms of real maximal ideals.

It was shown by Comfort in [1] and [2] that the local pseudocompactness of X plays an important role in connection with the local compactness of νX . The precise role is established below, where it is shown that the local pseudocompactness of X is equivalent to the local compactness of the extension ηX of X constructed by Johnson and Mandelker in [9]. In addition a characterization is given of those spaces for which the extension ψX constructed by Johnson and Mandelker is locally compact.

Our attention will be restricted entirely to completely regular Hausdorff spaces. The terminology and notation of [4] will be used without further comment.

Given $f \in C(X)$ the symbols N(f) and S(f) represent respectively $\{x \in X: f(x) \neq 0\}$ and $cl_x \{x \in X: f(x) \neq 0\}$; these sets are called the *cozero set* and the *support* of f. If A and B are subsets of X, write $A \ll B$ if A is completely separated from X - B. We shall frequently apply [4, 1.15] to construct additional separating zero sets when $A \ll B$.

The symbol M^p will denote the maximal ideal in C(X) which corresponds to the point p of βX , and \mathscr{M}^p will denote the corresponding *z*-ultrafilter (written A^p in [4]). Similarly O^p represents the ideal defined in [4, 7.12] and \mathscr{O}^p the corresponding *z*-filter. The referee deserves acknowledgment for a number of useful suggestions. Thanks are also due to Marquette University for a Summer Faculty Fellowship which supported a portion of this work.

2. The local compactness of νX . The family $C_{\psi}(X)$ of functions with pseudocompact support, discussed at length in [9] and [11], plays the major role in our condition for local compactness of νX . This is to be expected since by 2.1(d) below the isomorphism $f \to f^{\nu}$ is an isomorphism of $C_{\psi}(X)$ with $C_{\kappa}(\nu X)$. We write $\mathscr{C}_{\psi}(X)$ for the corresponding collection of zero sets.

The following results are either found in [4] or may be established using results from [4].

2.1. (a) $Z(f^{\circ}) = \operatorname{cl}_{\circ_X} Z(f)$. (b) $N(f^{\circ}) = \operatorname{int}_{\circ_X} (\circ X - Z(f))$. (c) $S(f^{\circ}) = \operatorname{cl}_{\circ_X} S(f) = \operatorname{cl}_{\circ_X} N(f)$. (d) $S(f^{\circ})$ is compact if and only if S(f) is pseudocompact. The following result from [11] is frequently useful.

2.2. Every support in a pseudocompact space is pseudocompact. Since the isomorphism $f \to f^{\circ}$ induces a bijection between real maximal ideals in C(X) and fixed ideals in C(vX), the following result is an immediate consequence of [4, 4D3].

THEOREM 2.3. The space $\cup X$ is locally compact if and only if $C_{\psi}(X)$ is not contained in any real maximal ideal.

We turn now toward a condition expressible in terms of X alone. For each $\varepsilon > 0$ and each $f \in C(X)$, define $U_{\varepsilon}(f) = \{x \in X : |f(x)| \ge \varepsilon\}$; this is a zero set in X. The following results are essential for our characterization theorem.

2.4. (a) If $\varepsilon > \delta > 0$ then $N(f) \gg U_{\delta}(f) \gg U_{\varepsilon}(f)$.

(b) If $p \in \beta X$ and $f \in C^*(X)$ then $f^{\beta}(p) = 0$ if and only if $U_{\varepsilon}(f) \notin \mathcal{M}^p$ for each $\varepsilon > 0$. (Also \mathcal{M}^p may be replaced by \mathcal{O}^p in this condition). (c) For each $f \in C(X)$, $U_{\varepsilon}(f^{\circ}) = \operatorname{cl}_{\gamma X} U_{\varepsilon}(f)$.

Proof. The proofs of (a) and (c) are straightforward. (b) For any $\varepsilon > 0$, if $U_{\varepsilon}(f) \in \mathscr{M}^{p}$, then $p \in \operatorname{cl}_{\beta_{X}} U_{\varepsilon}(f)$ and $f(p) \ge \varepsilon$, a contradiction. Hence $U_{\varepsilon}(f) \notin \mathscr{M}^{p}$.

Conversely let $U_{\varepsilon}(f) \notin \mathcal{O}^p$ for each $\varepsilon > 0$. For every $\varepsilon > 0$ we have $\{x \in X : |f(x)| \leq \varepsilon\} \in \mathcal{M}^p$ by [4, 7.12(b)]; hence $f^{\beta}(p) \leq \varepsilon$. Thus $f^{\beta}(p) = 0$.

Let U(X) represent the set of units in C(X); by [4, 1.12], these are the functions with empty zero set. Clearly the image of U(X) under the isomorphism $f \to f^{\circ}$ is U(vX).

A scale on X is a function $\varepsilon: f \to \varepsilon(f)$ from U(X) to the positive real numbers. If ε is a scale, put $\mathscr{C}(\varepsilon) = \{U_{\varepsilon(f)}(f): f \in U(X)\}.$

THEOREM 2.5. A z-ultrafilter is real if and only if it contains $\mathscr{C}(\varepsilon)$ for some scale ε .

Proof. Let ε be a scale on X and let \mathscr{M}^p be a hyper-real zultrafilter. By [4, 8.8] there is a bounded unit f in C(X) with $f^{\beta}(p) = 0$; hence $U_{\varepsilon(f)}(f) \notin \mathscr{M}^p$. Thus \mathscr{M}^p does not contain $\mathscr{C}(\varepsilon)$.

Let $p \in \upsilon X$. For any $f \in U(X)$ put $\varepsilon(f) = |f^{\upsilon}(p)|$. Since f^{υ} is a unit of $C(\upsilon X)$, $\varepsilon(f) > 0$, and thus ε is a scale on X. Since $p \in U_{\varepsilon(f)}(f^{\upsilon}) = \operatorname{cl}_{\upsilon X} U_{\varepsilon(f)}(f) \subset \operatorname{cl}_{\beta X} U_{\varepsilon(f)}(f)$, it follows that $U_{\varepsilon(f)}(f) \in \mathscr{M}^p$. Thus $\mathscr{C}(\varepsilon) \subset \mathscr{M}^p$.

COROLLARY 2.6. A filter \mathscr{F} on X is contained in a real z-ultrafilter if and only if every member of \mathscr{F} meets every member of $\mathscr{E}(\varepsilon)$ for some scale ε .

THEOREM 2.7. The space $\cup X$ is locally compact if and only if X satisfies the following condition: (RL). For every scale ε there are $f_1, \dots, f_k \in U(X)$ and $g \in C_{\psi}(X)$ such that $Z(g) \cap (\bigcap_{i=1}^k U_{\varepsilon(f_i)}(f_i)) = \phi$.

Proof. Let X satisfy (*RL*). For any $p \in \upsilon X$, by 2.5 there is a scale ε on X such that $\mathscr{C}(\varepsilon) \subset \mathscr{M}^p$. By (*RL*), $\mathscr{C}_{\psi}(X) \not\subset \mathscr{M}^p$; thus υX is locally compact by 2.3.

Suppose νX is locally compact and ε is a scale on X. By 2.3, $\mathscr{C}_{\psi}(X) \not\subset \mathscr{M}^{p}$ for any $p \in \nu X$. Thus, by 2.5, $\mathscr{C}_{\psi}(X) \cup \mathscr{C}(\varepsilon)$ lacks the finite intersection property; that is, condition RL is satisfied.

REMARK 2.8. It is clear that we need consider in condition RLonly those scales for which the family $\mathscr{C}(\varepsilon)$ has the finite intersection property, since the condition is trivially fulfilled when some finite subfamily of $\mathscr{C}(\varepsilon)$ has empty intersection. The condition is also fulfilled trivially when $\mathscr{C}_{\psi}(X)$ contains a unit of C(X), and this occurs precisely when X is pseudocompact.

Certainly if $\mathscr{C}_{\psi}(X)$ lacks the countable intersection property then it is not contained in a real z-ultrafilter. It will now be shown that $\mathscr{C}_{\psi}(X)$ lacks the property precisely when υX is locally compact and σ -compact. Our condition is shown to be related to one given by Hager in [7].

THEOREM 2.9. The following are equivalent for a space X. (a) $\cup X$ is locally compact and σ -compact.

DOUGLAS HARRIS

(b) $\mathscr{C}_{\psi}(X)$ lacks the countable intersection property.

(c) (Hager) $X = \bigcup_{n=1}^{\infty} A_n$, where each A_n is pseudocompact and $A_n \ll A_{n+1}$ for each n.

Proof. (a) implies (c). If νX is locally compact and σ -compact then [3, XI, 7.2] $\nu X = \bigcup_{n=1}^{\infty} U_n$, where each U_n is open and has compact closure and $\operatorname{cl}_{\nu X} U_n \subset U_{n+1}$ for each n. By [4, 3.11(a)], $U_n \ll U_{n+1}$. Setting $A_n = \operatorname{cl}_X (U_n \cap X)$ it follows from [2, 4.1] that each A_n is pseudo-compact.

(c) implies (b). Let $X = \bigcup_{n=1}^{\infty} A_n$, with A_n pseudocompact and $A_n \ll A_{n+1}$ for each n. Choose for each n a function f_n such that $A_n \subset N(f_n) \subset S(f_n) \subset A_{n+1}$. Then, by [2.2] each $f_n \in C_{\psi}(X)$, and clearly $\bigcap_{n=1}^{\infty} Z(f_n) = \phi$.

(b) implies (a). Let $\bigcap_{n=1}^{\infty} Z(f_n) = \phi$, where $f_n \in C_{\psi}(X)$ for each *n*. Then, by 2.1(a) and [4, 8.7], $\bigcap_{n=1}^{\infty} Z(f_n^{\circ}) = \phi$, and thus $\bigcup_{n=1}^{\infty} S(f_n^{\circ}) = \upsilon X$. By 2.1(d) each $S(f_n^{\circ})$ is compact, thus υX is σ -compact. By 2.3, υX is locally compact.

Comfort [2, 4.6] gives another condition (C) which is equivalent to the local compactness of vX. A direct proof of the equivalence of (C) with the condition of Theorem 2.3 will now be given.

2.10. $\mathscr{C}_{\psi}(X)$ is not contained in any real z-ultrafilter if and only if: (C) For each $p \in vX$ there exist pseudocompact subsets A and B of X such that $p \in cl_{\nu X} A$ and $A \ll B$.

Proof. If X satisfies condition (C) and \mathscr{M}^p is a real maximal ideal then there are pseudocompact sets A and B and functions $f, g \in C(X)$ such that $p \in \operatorname{cl}_{_{v_X}} A$ and $A \subset Z(f) \subset N(g) \subset B$. It follows from 2.2 that $g \in C_{\psi}(X)$. Since $p \in \operatorname{cl}_{_{v_X}} A$ then $f \in M^p$, and thus $g \notin M^p$. Thus $C_{\psi}(X) \not\subset M^p$.

Conversely, for any $p \in vX$ there is $f \in C_{\psi}(X)$ and $g \in \mathscr{M}^{p}$ such that $Z(f) \cap Z(g) = \phi$; thus $Z(g) \ll N(f)$ and there exist $h, k \in C(X)$ such that $Z(g) \subset N(k) \subset Z(h) \subset N(f)$. Put A = S(k) and B = S(f). Since $A \subset S(f)$, it follows from 2.2 that A is pseudocompact. Also $p \in cl_{\beta X}Z(g)$, since $g \in \mathscr{M}^{p}$, so $p \in cl_{v X} A$. Finally, $A \subset Z(h)$ and $X - B \subset Z(f)$, with $Z(h) \cap Z(f) = \phi$, so $A \ll B$. Thus condition (C) is satisfied.

3. The local pseudocompactness of X. The space X is locally pseudocompact if every point has a pseudocompact neighborhood. Locally pseudocompact spaces are discussed in [1] and [2]. The results in this section clarify the relationship between the local pseudocompactness of X and the local compactness of vX.

472

3.1. The space X is locally pseudocompact if and only if $C_{\psi}(X)$ is not contained in any fixed maximal ideal.

Proof. Let X be locally pseudocompact. Then any point x in X has a pseudocompact neighborhood A. Therefore, there is $f \in C(X)$ with $x \in N(f) \subset A$. Thus $x \notin Z(f)$ and by 2.2, the set S(f) is pseudocompact, so $f \in C_{\psi}(X)$. Therefore, $C_{\psi}(X)$ is not contained in any fixed maximal ideal. Conversely suppose $C_{\psi}(X)$ is contained in no fixed maximal ideal. Then for each $x \in X$ there is $f \in C_{\psi}(X)$ with $x \in N(f)$, and thus S(f) is a pseudocompact neighborhood of x.

For any space Y denote by L(Y) the set of all points of Y that have a compact neighborhood in Y; i.e., L(Y) = Y - R(Y), where R(Y)is as defined in [8, p. 87]. Clearly L(Y) is locally compact. For any space X define $\kappa X = \{p \in \beta X : \mathscr{C}_{\psi}(X) \not\subset \mathscr{M}^p\}$; equivalently, $\kappa X = \beta X - \theta(\mathscr{C}_{\psi}(X))$, where $\theta(\mathscr{C}_{\psi}(X))$ is as defined in [4, 70].

THEOREM 3.2. For each space $X, \kappa X = L(\upsilon X) = \operatorname{int}_{\beta_X} \upsilon X$, and thus κX is locally compact.

Proof. The relation $L(\upsilon X) = \operatorname{int}_{\beta_X} \upsilon X$ follows from [4, 3.15(b)]. By [9, 3.1], $\beta X - \kappa X = \theta(\mathscr{C}_{\psi}(X)) = \operatorname{cl}_{\beta_X}(\beta X - \upsilon X)$, so $\kappa X = \operatorname{int}_{\beta_X} \upsilon X$.

COROLLARY 3.3. The space X is locally pseudocompact if and only if $X \subset \kappa X$. In this case κX is the largest locally compact space between X and υX .

The following result is due to Comfort ([1] and [2]).

COROLLARY 3.4. The space X is locally pseudocompact if and only if there is a locally compact space Y between X and $\cup X$.

4. Functions with small support. Another ideal in C(X) plays an important role in connection with local compactness. Before discussing this ideal, the class of *small* sets will be examined, where a set $A \subset X$ is *small* if any zero set contained in A is compact.

4.1. The set A is small if and only if every zero set that intersects X - A in a compact set is compact.

Proof. Certainly in the latter condition holds then A is small. Now suppose A is small and Z is a zero set such that $Z \cap (X - A)$ is compact. If α is a cover of Z by cozero sets then finitely many of the cozero sets cover $Z \cap (X - A)$. Their union is a cozero set N(g) and $Z(g) \cap Z$ is compact, since it is a zero set. Thus, finitely many additional members of α can be chosen to complete the choice of a finite subcover of Z.

4.2. The finite union of small cozero sets is small.

Proof. Let N(f) and N(g) be small, and let $Z(h) \subset N(f) \cup N(g)$. Then $Z(h) \cap (X - N(g)) = Z(h) \cap Z(g) \subset N(f)$. Since N(f) and N(g) are small it follows from 4.1 that Z(h) is compact.

A function $f \in C(X)$ has small support if and only if N(f) is small. Equivalently, according to [4, 4E2], the function f belongs to every free maximal ideal in C(X). It is clear from this latter characterization that the collection $C_s(X)$ of functions with small support is an ideal; this can also be shown directly from 4.2.

REMARK 4.3. The term small support may be misleading; the condition applies to N(f) and not S(f). The ideal $C_s(X)$ contains the ideal $C_{\kappa}(X)$ [4, 4D5 and 4E2]. Spaces for which $C_{\kappa}(X) = C_s(X)$ are called μ -compact and are fully discussed in [9] and [11]; in [9] the ideal $C_s(X)$ is called I(X).

The following result should be compared with [4, 4D1 and 4D3], as well as with Theorem 2.3.

THEOREM 4.4. The space X is locally compact if and only if $C_s(X)$ is not contained in any fixed maximal ideal.

Proof. If X is locally compact then $C_{\kappa}(X)$ is not contained in any fixed maximal ideal; since $C_{\kappa}(X) \subset C_{s}(X)$ then $C_{s}(X)$ is not contained in any fixed maximal ideal.

Now if $C_s(X)$ is not contained in any fixed maximal ideal then for each $x \in X$ there is $f \in C_s(X)$ such that $x \in N(f)$. Thus, there is a zero set neighborhood of x such that $Z \subset N(f)$, and it follows that Z is compact. Thus X is locally compact.

REMARK 4.5. One sense in which Theorem 4.4 is more appropriate than the characterization [4, 4D3] of local compactness is when the generalization to T_1 spaces and T_1 compactifications is considered. In [5] the compact small sets of a space X are defined as those sets such that any closed set contained in A is compact. It is shown there that the spaces for which each point has a compact-small neighborhood are appropriate generalizations of locally compact completely regular spaces. It is shown in [6] that results analogous to Theorem 4.4 hold for locally compact-small spaces.

5. The local compactness of ηX and ψX . Two additional

subspaces of νX are of special interest in connection with local compactness. Mandelker defines (in [11]) a space X to be ψ -compact if $C_{\kappa}(X) = C_{\psi}(X)$, and Mandelker and Johnson define (in [9]) a space X to be η -compact if $C_s(X) = C_{\psi}(X)$; they construct extensions ηX and ψX as the intersections respectively of the η - compact and the ψ -compact subspaces of βX .

The following results are shown in [9].

5.1. (a) $\eta X = X \cup \operatorname{int}_{\beta_X} \upsilon X$.

(b) $\psi X - X = \bigcup_{f \in C \psi(X)} [S(f^{\upsilon}) - S(f)].$

The next results are immediate from 5.1(a) and Theorem 3.2.

5.2. (a) $\kappa X = \operatorname{int}_{\beta_X} \eta X = \operatorname{int}_{\beta_X} \psi X$.

(b) $\eta X = X \cup \kappa X$

The next theorem characterizes the local compactness of ηX . The proof is immediate from 5.2 and Corollary 3.3.

THEOREM 5.3. The space ηX is locally compact if and only if X is locally pseudocompact.

THEOREM 5.4. The space ψX is locally compact if and only if X is locally pseudocompact and $\mathscr{C}_{\psi}(X)$ is round.

Proof. Let ψX be locally compact. Then X is locally pseudocompact by Corollary 3.4. Also ψX is open in βX , so $\beta X - \psi X$ is closed. By [9, 5.3], $C_{\psi}(X) = M^{\beta X - \psi X}$ and thus $\beta X - \psi X$ is round; hence by [10, 4.2] $\mathscr{C}_{\psi}(X)$ is round.

Let X be locally pseudocompact and let $\mathscr{C}_{\psi}(X)$ be round. By 3.3 and 5.2(a), $X \subset \kappa X \subset \psi X$. Let $p \in \upsilon X - X$. Using 5.1(b) choose $f \in C_{\psi}(X)$ so that $p \in S(f^{\upsilon})$; since $\mathscr{C}_{\psi}(X)$ is round there is $g \in C_{\psi}(X)$ with $Zg \ll Zf$. By [4, 7.14], $\operatorname{cl}_{\beta_X} Z(f)$ is a neighborhood of $\operatorname{cl}_{\beta_X} Z(g)$, and thus there is a compact set F with $\beta X - \operatorname{cl}_{\beta_X} Z(f) \subset F \subset \beta X - \operatorname{cl}_{\beta_X} Z(g)$. Since $N(f) \subset \beta X - \operatorname{cl}_{\beta_X} Z(f)$ and (by [9, 3.1]) $\beta X - \upsilon X \subset \operatorname{cl}_{\beta_X} Z(g)$, it follows that $p \in S(f^{\upsilon}) \subset F \subset \beta X - \operatorname{cl}_{\beta_X} Z(g) \subset \upsilon X$, and hence

$$p \in \operatorname{int}_{\beta_X} \upsilon X = \kappa X$$
.

Thus $\psi X = \kappa X$ and so ψX is locally compact.

It is instructive in the use of scales to deduce directly from Condition (RL) that X is locally pseudocompact and $\mathscr{C}_{\psi}(X)$ is round.

5.5. If X satisfies Condition (*RL*) then X is locally pseudocompact and $\mathscr{C}_{\psi}(X)$ is round.

Proof. The first paragraph of the proof of Theorem 2.7 shows that X will be locally pseudocompact. Now suppose $f \in C_{\psi}(X)$. Then

DOUGLAS HARRIS

S(f) is pseudocompact, and it follows that every $h \in U(X)$ is bounded aways from zero on N(f). Choose a scale ε so that $|h| \ge \varepsilon(h)$ on N(f), for each $h \in U(X)$. Since (RL) is satisfied, there are $h_1, \dots, h_k \in U(X)$ and $g \in C_{\psi}(X)$ such that $Zg \cap (\cap U_{\varepsilon(h_i)}(h_i)) = \emptyset$. Clearly

$$Z(g) \subset igcup_{i=1}^k \left\{ x \in X
ight: |h_i(x)| < arepsilon(h_i)
ight\} \subset Z(f)$$
 .

It follows that $g \ll f$.

References

1. W. W. Comfort, Locally compact realcompactifications, General Topology and its Relations to Modern Analysis and Algebra II, Proceedings of the Second Prague Topological Symposium, (1966), 95-100.

2. ____, On the Hewitt realcompactification of a product space, Trans. Amer. Math. Soc., **131** (1968), 107-118.

3. J. Dugundji, Topology, Allyn and Bacon, Boston, 1966.

4. L. Gillman and M. Jerison, *Rings of Continuous Functions*, Van Nostrand, Princeton, 1960.

5. D. Harris, Closed images of the Wallman compactification, Proc. Amer. Math. Soc., 42 (1974), 312-319.

6. _____, Semirings and T_1 -compactifications I, Trans. Amer. Math. Soc., to appear. 7. A. W. Hager, On the tensor product of function rings, Doctoral Dissertation, Pennsylvania State University, University Park, Pa., 1965.

8. M. Henriksen and J. Isbell, Some properties of compactifications, Duke Math. J., 25 (1958), 83-106.

9. D. G. Johnson and M. Mandelker, *Functions with pseudocompact support*, General Topology and Appl., to appear.

10. M. Mandelker, Round z-filters and round subsets of βX , Israel J. Math., 7 (1969), 1-8.

11. ____, Supports of continuous functions, Trans. Amer. Math. Soc., 156 (1971), 73-83.

Received October 27, 1972 and in revised form July 25, 1973.

MARQUETTE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor) University of California Los Angeles, California 90024 J. Dugundji*

Department of Mathematics University of Southern California Los Angeles, California 90007

D. GILBARG AND J. MILGRAM Stanford University Stanford, California 94305

ASSOCIATE EDITORS

E.F. BECKENBACH

R. A. BEAUMONT

University of Washington

Seattle, Washington 98105

B.H. NEUMANN

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIAUNICALIFORNIA INSTITUTE OF TECHNOLOGYSTAUNIVERSITY OF CALIFORNIAUNIMONTANA STATE UNIVERSITYUNIUNIVERSITY OF NEVADAWASNEW MEXICO STATE UNIVERSITYUNIOREGON STATE UNIVERSITYUNIUNIVERSITY OF OREGONAMIOSAKA UNIVERSITYNAV

UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON * * * * AMERICAN MATHEMATICAL SOCIETY

K. YOSHIDA

NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$60.00 a year (6 Vols., 12 issues). Special rate: \$30.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by Pacific Journal of Mathematics Manufactured and first issued in Japan

Pacific Journal of Mathematics Vol. 50, No. 2 October, 1974

Mustafa Agah Akcoglu, John Philip Huneke and Hermann Rost, <i>A counter</i> <i>example to the Blum Hanson theorem in general spaces</i>	305
Huzihiro Araki, Some properties of modular conjugation operator of von Neumann algebras and a non-commutative Radon-Nikodym theorem with a chain rule	309
E. F. Beckenbach, Fook H. Eng and Richard Edward Tafel, <i>Global</i>	355
David W Boyd A new class of infinite sphere packings	383
K. G. Choo. Whitehead Groups of twisted free associative algebras	399
Charles Kam-Tai Chui and Milton N. Parnes, <i>Limit sets of power series</i> outside the circles of convergence	403
Allan Clark and John Harwood Ewing, The realization of polynomial	
algebras as cohomology rings	425
Dennis Garbanati, Classes of circulants over the p-adic and rational	
integers	435
Arjun K. Gupta, On a "square" functional equation	449
David James Hallenbeck and Thomas Harold MacGregor, Subordination	
and extreme-point theory	455
Douglas Harris, The local compactness of vX	469
William Emery Haver, <i>Monotone mappings of a two-disk onto itself which</i> <i>fix the disk's boundary can be canonically approximated by</i>	
homeomorphisms	477
Norman Peter Herzberg, On a problem of Hurwitz	485
Chin-Shui Hsu, A class of Abelian groups closed under direct limits and subgroups formation	495
Bjarni Jónsson and Thomas Paul Whaley, <i>Congruence relations and</i> multiplicity types of algebras	505
I owell Duane I oveland Vertically countable spheres and their wild	505
sets	521
Nimrod Megiddo, Kernels of compound games with simple components	531
Russell L. Merris, An identity for matrix functions	557
E. O. Milton, Fourier transforms of odd and even tempered distributions	563
Dix Hayes Pettey, One-one-mappings onto locally connected generalized	
continua	573
Mark Bernard Ramras, Orders with finite global dimension	583
Doron Ravdin, Various types of local homogeneity	589
George Michael Reed, On metrizability of complete Moore spaces	595
Charles Small, Normal bases for quadratic extensions	601
Philip C. Tonne, Polynomials and Hausdorff matrices	613
Robert Earl Weber, <i>The range of a derivation and ideals</i>	617