

Pacific Journal of Mathematics

**MONOTONE MAPPINGS OF A TWO-DISK ONTO ITSELF
WHICH FIX THE DISK'S BOUNDARY CAN BE CANONICALLY
APPROXIMATED BY HOMEOMORPHISMS**

WILLIAM EMERY HAVER

MONOTONE MAPPINGS OF A TWO-DISK ONTO ITSELF
WHICH FIX THE DISK'S BOUNDARY CAN BE
CANONICALLY APPROXIMATED BY
HOMEOMORPHISMS

WILLIAM E. HAVER

The theorem stated in the title is proven. As a corollary it is shown that the space of all such monotone mappings is an absolute retract.

1. Introduction. Let D^n denote the standard n -ball of radius one in E^n and $H(D^n)$ the space of all homeomorphisms of D^n onto itself which equal the identity on the boundary of D^n . Let $\overline{H(D^n)}$ denote the space of all mappings of D^n onto itself which can be approximated arbitrarily closely by elements of $H(D^n)$. Under the supremum topology, $H(D^n)$ and $\overline{H(D^n)}$ are separable metric spaces; $\overline{H(D^n)}$ is complete under the supremum metric. It is known that $\overline{H(D^n)}$ is locally contractible [7], $\overline{H(D^n)} \times l_2 \approx \overline{H(D^n)}$ [4], $\overline{H(D^n)}$ is homogeneous [7], and that $\overline{H(D^1)} \approx l_2$ [3]. In this paper we shall be concerned with the case $n = 2$, and to simplify notation we shall write D for D^2 . It is well-known (cf. [8]) that $\overline{H(D)}$ is the space of all monotone mappings of D onto itself which equal the identity when restricted to the boundary of D .

We shall show [Theorem 1] that the elements of $\overline{H(D)}$ can be "canonically approximated" by elements of $H(D)$ and [Theorem 2] that $\overline{H(D)}$ is an absolute retract. The work of this paper depends heavily on W. K. Mason's paper, "The space of all self-homeomorphisms of a two-cell which fix the cell's boundary is an absolute retract", [9]. The crux of Mason's paper is the definition of a basis for $H(D)$ which can be shown to possess some particularly nice properties. We shall review the definition of this basis in the following section and then define a basis for $\overline{H(D)}$. Familiarity will be assumed with the notation and basic definitions of [9].

2. Mason's basis for $H(D)$. Consider D to be a rectangle in R^2 with horizontal and vertical sides. A grating, P , on D consists of a finite number of spanning segments (crosscuts) across D , parallel to its sides, with the same number of horizontal and vertical crosscuts. Let P_1, P_2, \dots be a sequence of gratings on D such that (a) the mesh of P_i approaches 0 as i increases and (b) if l is a crosscut of P_i and $j \geq i$, then l is a crosscut of P_j .

Let \mathcal{H} be the collection of all polyhedral disks H contained in D

such that $\text{Bd}(H)$ is the union of a vertical segment in the left side of $\text{Bd}(D)$, a vertical segment in the right side of $\text{Bd}(D)$, a polygonal spanning arc of D, H^T , that is contained in the closure of the same component of $H(D) - H$ as the top of $\text{Bd}(D)$, and a polygonal spanning arc of D, H^B , that is contained in the closure of the same component of $H(D) - H$ as the bottom of $\text{Bd}(D)$. Let \mathcal{V} be the collection of all polyhedral disks V contained in D such that $\text{Bd}(V)$ is the union of a horizontal segment in the top of $\text{Bd}(D)$, a horizontal segment in the bottom of $\text{Bd}(D)$, a polygonal spanning arc of D, V^L , that is contained in the closure of the same component of $H(D) - V$ as the left side of $\text{Bd}(D)$ and a polygonal spanning arc of D, V^R , that is contained in the closure of the same component of $H(D) - V$ as the right side of $\text{Bd}(D)$.

Let P_j be a grating from the sequence P_1, P_2, \dots . Let $\{l_1, \dots, l_n\}$ be the set of horizontal crosscuts of P_j and $\{m_1, \dots, m_n\}$ the set of vertical crosscuts. Let $\{H_1, \dots, H_n\} \subset \mathcal{H}$ satisfy $H_i \cap H_j = \emptyset$ if $i \neq j$ and $\{V_1, \dots, V_n\} \subset \mathcal{V}$ satisfy $V_i \cap V_j = \emptyset$ if $i \neq j$. Then define

$$\begin{aligned} O(P_j; H_1, \dots, H_n; V_1, \dots, V_n) &= \{f \in H(D) \mid f(l_i) \subset H_i - \{\text{Cl}(D - H_i)\} \text{ and} \\ &\quad f(m_i) \subset V_i - \{\text{Cl}(D - V_i)\} \text{ for } 1 \leq i \leq n\}. \end{aligned}$$

Then the basis for $H(D)$, which Mason denotes HVT , is the collection of all such open sets.

3. A Basis for $\overline{H(D)}$. In this section we define a basis, β , for $\overline{H(D)}$ and demonstrate that it possesses some nice properties. Let $P_j, \{H_1, \dots, H_n\}$ and $\{V_1, \dots, V_n\}$ be as in the definition of HVT . The basis, β , will consist of all sets of the following form:

$$\begin{aligned} B(P_j; H_1, \dots, H_n; V_1, \dots, V_n) &= \{f \in \overline{H(D)} \mid f^{-1}(l_i) \subset H_i - \{\text{Cl}(D - H_i)\} \text{ and} \\ &\quad f^{-1}(m_i) \subset V_i - \{\text{Cl}(D - V_i)\}, \text{ for } 1 \leq i \leq n\}. \end{aligned}$$

We note that $f \in B(P_j; H_1, \dots, H_n; V_1, \dots, V_n) \cap H(D)$ if and only if $f^{-1} \in O(P_j; H_1, \dots, H_n; V_1, \dots, V_n)$. To see that the elements of β are open subsets of $\overline{H(D)}$, let f be an arbitrary element of $B(P_j; H_1, \dots, H_n; V_1, \dots, V_n)$. Then let

$$\varepsilon = \min_{1 \leq i \leq n} \{d(f(H_i^T \cup H_i^B), \bigcup_{j=1}^n l_j), d(f(V_i^L \cup V_i^R), \bigcup_{j=1}^n m_j)\}.$$

Let g be an arbitrary element of $\overline{H(D)}$ satisfying $d(f, g) < \varepsilon$. Suppose that $g \notin B(P_j; H_1, \dots, H_n; V_1, \dots, V_n)$. Then, without loss of generality, we can assume that there exists an integer i such that $g^{-1}(l_i)$

is not contained in $H_i - \text{Cl}(D - H_i)$. Since (i) $g^{-1}(l_i) \cap (\text{Bd } D) = f^{-1}(l_i) \cap (\text{Bd } D) \subset H_i - \text{Cl}(D - H_i)$, (ii) $g^{-1}(l_i)$ is a connected set and (iii) $H_i^T \cup H_i^B$ separates $H_i - \text{Cl}(D - H_i)$ from $D - H_i$, there is an $x \in H_i^T \cup H_i^B$ such that $g(x) \in l_i$. But this implies that $d(f(x), g(x)) \geq d(f(H_i^T \cup H_i^B), l_i) \geq \varepsilon$ and hence $d(f, g) \geq \varepsilon$.

LEMMA 1. β is a basis for $\overline{H(D)}$.

Proof. Let $f \in \overline{H(D)}$ and $\varepsilon > 0$ be given. We wish to find $B \in \beta$ such that $f \in B$ and $d(f, g) < \varepsilon$ for all $x \in B$. Pick a grating, P_j , such that $\text{diam } |st(x, P_j)| < \varepsilon$ for every $x \in D$. Let l_i be the i th crosscut from the top of D . Choose H_1^T to be a polygonal spanning arc of D with one endpoint in each side of D that separates the top of D from $f^{-1}(l_1)$. Choose H_1^B to be a polygonal spanning arc of D that separates $f^{-1}(l_1)$ from $f^{-1}(l_2)$. The polyhedral disk H_1 is thus uniquely defined. Assume inductively that disks H_1, \dots, H_{i-1} have been defined in such a manner that $H_j \cap H_k = \emptyset$ if $1 \leq j \leq i-1, 1 \leq k \leq i-1$, and $j \neq k$ and that for each $j, 1 \leq j \leq i-1, H_j^T$ separates H_{j-1}^B from $f^{-1}(l_j)$ and H_j^B separates $f^{-1}(l_j)$ from $f^{-1}(l_{j+1})$. Choose H_i^T to be a polygonal spanning arc of D that separates H_{i-1}^B from $f^{-1}(l_i)$. Finally choose H_i^B to be a polygonal spanning arc of D that separates $f^{-1}(l_i)$ from $f^{-1}(l_{i+1})$ (or from the bottom of D if $i = n$). We have thus uniquely defined H_i in such a way that the inductive hypothesis is satisfied. Define V_1, \dots, V_n in a similar manner. Now, by construction $f \in B(P_j; H_1, \dots, H_n; V_1, \dots, V_n)$ and if $g \in B(P_j; H_1, \dots, H_n; V_1, \dots, V_n)$ then $d(f, g) < \varepsilon$.

LEMMA 2. Let B_1, \dots, B_j be elements of β . Then $B = \bigcap_{k=1}^j B_k$ is an element of β .

Proof. Assume $B \neq \emptyset$ and that P_k is the grating associated with $B_k, 1 \leq k \leq j$. Hence for any $k, 1 \leq k \leq j$, every crosscut of P_k is a crosscut of P_j . Let l_1 be the first horizontal crosscut of P_j . For each $k, 1 \leq k \leq j$, let $H_{1,k}$ be the element of \mathcal{H} associated with l_1 and B_k (if there is one). Let H_1 be the component of $D - (\bigcup_{k=1}^j H_{1,k}^T \cup \bigcup_{k=1}^j H_{1,k}^B)$ that contains $f^{-1}(l_1)$. Define in an analogous manner H_2, \dots, H_n and V_1, \dots, V_n . It is clear that $B(P_j; H_1, \dots, H_n; V_1, \dots, V_n) = \bigcap_{k=1}^j B_k$. The elements of $\{H_1, \dots, H_n\}$ are pairwise disjoint since each H_i is contained in $H_{i,n}$ and the elements of $\{H_{1,j}, \dots, H_{n,j}\}$ are pairwise disjoint.

Lemma 3 will follow as a corollary to the following theorem of Mason. (The proof of this theorem constitutes the bulk of [9].)

THEOREM (Mason). Let U be an element of HVT and K a finite

dimensional compact subset of U . Then there is an embedding ψ of the cone over K into U such that $\psi(f, 0) = f$, for all $f \in K$.

LEMMA 3. Let $B = B(P_j; H_1, \dots, H_n; V_1, \dots, V_n)$ be an element of β and K a finite dimensional compact subset of $B \cap H(D)$. Then there is an embedding λ of the cone over K into $B \cap H(D)$ such that $\psi(f, 0) = f$, for all $f \in K$.

Proof. Since $H(D)$ is a topological group, the function $G: H(D) \rightarrow H(D)$ defined by $G(f) = f^{-1}$ is a homeomorphism. Therefore, by the note following the definition of β , $G(K)$ is a finite dimensional compact subset of $U(P_j; H_1, \dots, H_n; V_1, \dots, V_n)$. Hence by Mason's theorem there is an embedding ψ of the cone over $G(K)$ into $U(P_j; H_1, \dots, H_n; V_1, \dots, V_n)$ such that $\psi(f, 0) = f$, for all $f \in G(K)$. Define $\lambda: K \times I \rightarrow B \cap H(D)$ by $\lambda(k, t) = G^{-1}(\psi(G(k), t))$.

4. The main results. The following theorem shows that the elements of $\overline{H(D)}$ can be canonically approximated by elements of $H(D)$.

THEOREM 1. Let α be an open cover of $\overline{H(D)}$. Then there exists a locally finite polyhedron, \mathcal{P} , and maps $b: \overline{H(D)} \rightarrow \mathcal{P}$, $\psi: \mathcal{P} \rightarrow H(D)$, and $\theta: \overline{H(D)} \times I \rightarrow \overline{H(D)}$ such that

- (a) for each $f \in \overline{H(D)}$, there is an element, U_f , of α such that $\theta(f, t) \in U_f$, for each $t \in I$,
- (b) $\theta(f, 1) = f$, for each $f \in \overline{H(D)}$,
- (c) $\theta(f, 0) = \psi b(f)$, for each $f \in \overline{H(D)}$,
- (d) $\theta(f, t) \in H(D)$ for each $f \in \overline{H(D)}$ and $t \in [0, 1)$.

Proof. Let α' be an open barycentric refinement α (i.e., if $f \in \overline{H(D)}$, $|st(f, \alpha')|$ is contained in some element of α). For each positive integer, k , let α_k be an open cover of $\overline{H(D)}$ such that

- (i) α_k is a refinement of α' ,
- (ii) if $V \in \alpha_k$, $\text{diam } V < 1/k$,
- (iii) if $V \in \alpha_k$, then $V \in \beta$.

We next define an open cover, η , of $\overline{H(D)} \times [0, 1)$.

$$\text{Let } \eta = \{V \times [0, 1/2) \mid V \in \alpha_i\} \cup \bigcup_{k=2}^{\infty} \left\{ V \times \left(\frac{2^k - 3}{2^k}, \frac{2^k - 1}{2^k} \right) \mid V \in \alpha_k \right\}.$$

Let γ be a countable refinement of η such that

- (a) if $h \in \gamma$, $st(h, \gamma)$ is a finite set,
- (b) if $h \in \gamma$, then there is an element of η , $V \times J$, such that $|st(h, \gamma)| \subset V \times J$.

Let \mathcal{P} be the nerve of γ and $B: \overline{H(D)} \times [0, 1) \rightarrow \mathcal{P}$ be the standard

barycentric map. Order the element of γ , and for each $h_i \in \gamma$, let $V_i \times J_i$ be an element of η such that $|st(h_i, \gamma)| \subset V_i \times J_i$. Note that $V_i \in \beta$.

We shall define a map $\psi: \mathcal{S} \rightarrow H(D)$ by induction on the skeletons of \mathcal{S} . For each vertex (h_i) of \mathcal{S} , let $\psi^0((h_i))$ be an arbitrary element of $H(D)$ intersected with the projection of h_i onto $\overline{H(D)}$.

Now assume that for $m = 1, 2, \dots, n$ we have defined $\psi^m: \mathcal{S}^m \rightarrow H(D)$ such that ψ^m extends ψ^{m-1} and for each simplex $\sigma^m = (h_{\lambda_0}, \dots, h_{\lambda_m})$:

- (a) $\psi^m(\sigma^m)$ is finite dimensional,
- (b) $\psi^m(\sigma^m) \subset H(D) \cap \{V_i \mid h_i \subset \bigcap_{j=0}^m st(h_{\lambda_j}, \gamma)\}$.

Now let $\sigma^{n+1} = \langle h_0, \dots, h_{n+1} \rangle$ be any simplex of \mathcal{S}^{n+1} . Let

$$U = \bigcap \{V_i \mid h_i \subset \bigcap_{j=0}^{n+1} st(h_j, \gamma)\}.$$

Since each V_i is an element of β , by Lemma 2, $U \in \beta$. By the inductive hypothesis the image under ψ^n of the boundary of σ^{n+1} is a finite dimensional compact subset of $U \cap H(D)$, denoted K . By Lemma 3 there is an embedding

$$\lambda: \subset (K) \rightarrow U \cap H(D)$$

such that $\lambda(f, 0) = f$ for all $f \in K$. We consider σ^{n+1} to be the cone over its boundary, and so for $(x, t) \in \sigma^{n+1}$, let $\psi^{n+1}(x, t) = \lambda(\psi^n(x), t)$.

Extending over each $n + 1$ simplex in this manner gives $\psi^{n+1}: \mathcal{S}^{n+1} \rightarrow H(D)$ and completes the induction. Hence $\lim_{n \rightarrow \infty} \psi^n = \psi: \mathcal{S} \rightarrow H(D)$ is continuous by the continuity of each ψ^n and the local finiteness of \mathcal{S} .

Let $b: \overline{H(D)} \rightarrow \mathcal{S}$ be defined by $b(f) = B((f, 0))$.

We next define the homotopy $\theta: \overline{H(D)} \times I \rightarrow \overline{H(D)}$ in the following manner:

$$\theta(f, t) = \begin{cases} \psi(B(f, t)), & t \neq 1 \\ f, & t = 1. \end{cases}$$

Conditions (b), (c), and (d) are obviously satisfied. We show simultaneously that θ is continuous and that for each $f \in \overline{H(D)}$ there is an element U_f of α such that for each $t \in I$, $\theta(f, t) \in U_f$.

Suppose that $(f, t) \in \overline{H(D)} \times [0, 1)$ and that $(2^k - 3)/2^k < t < (2^k - 1)/2^k$. Let h_0 be any element of γ which contains (f, t) . By the definition of ψ , $\psi B(f, t) \in V_0$. But $V_0 \in \alpha_{k-1} \cup \alpha_k \cup \alpha_{k+1}$ and therefore the diameter of V_0 is less than $1/(k - 1)$ which implies that $d(\psi B(f, t), f) < 1/(k - 1)$ and thereby that θ is continuous. Since each α_k is a refinement of α' , there exists an element of α' , $U_{(f,t)}$, such that

$\{f\} \cup \{\psi B(f, t)\} \subset V_0 \subset U_{(f,t)}$. Since α' is a barycentric refinement of α , there is some element, U_f , of α such that $\bigcup_{t \in [0,1]} U_{(f,t)} \subset U_f$ and hence $\theta(f, t) \in U_f$, for each $t \in I$.

The following result is an immediate corollary of Theorem 1 and a theorem of Hanner [5] which states that a metric space X is an ANR if given an arbitrary cover, α , of X there exists a locally finite polyhedron \mathcal{P} , maps $b: X \rightarrow \mathcal{P}$, $\psi: \mathcal{P} \rightarrow X$, and $\theta: X \times I \rightarrow X$ such that $\theta(x, 0) = \psi b(x)$ for all $x \in X$, $\theta(x, 1) = x$ for all $x \in X$ and for each $x \in X$ there is an element U of α such that $\theta(x, t) \in U$ for all $t \in [0, 1]$.

THEOREM 2. $\overline{H(D)}$ is an absolute retract.

Proof. By the preceding comments, $\overline{H(D)}$ is an ANR. But $\overline{H(D)}$ is contractible by the Alexander isotopy [1] applied to $\overline{H(D)}$. The theorem follows since every contractible absolute neighborhood retract is an absolute retract.

5. Applications. (a) The author has shown [6] that $\overline{H(M)}$, the space of all mappings of a compact manifold onto itself which can be approximated arbitrarily closely by homeomorphisms, is weakly locally contractible. Theorem 1 can be used [7] to show that for any compact 2-manifold, M^2 , $\overline{H(M^2)}$ is locally contractible.

(b) A problem of current interest is whether $H(D)$ is homeomorphic to l_2 ; it can easily be shown using a result of Anderson [2] that if $\overline{H(D)}$ is homeomorphic to l_2 , then $H(D)$ is homeomorphic to l_2 . Perhaps the results of this paper and the fact that $\overline{H(D)}$ is complete under the usual metric will be helpful in showing that $\overline{H(D)}$ is homeomorphic to l_2 .

(c) L. C. Siebenmann [10] has asked whether the inclusion map $i: H(M) \rightarrow \overline{H(M)}$ is a homotopy equivalence. Theorem 1 provides an affirmative answer to the question for the special case $i: H(D) \rightarrow \overline{H(D)}$.

Added in proof. Recent work of Toruńczyk ("Absolute retracts as factors of normed linear spaces," Fund. Math., to appear) implies that since $\overline{H(D)}$ is an AR and $H(D) \times l_2 \approx \overline{H(D)}$, $\overline{H(D)}$ is in fact homeomorphic to l_2 .

REFERENCES

1. J. W. Alexander, *On the deformation of an n -cell*, Proc. Nat. Acad. Sci., U.S.A., **9** (1923), 406-407.
2. R. D. Anderson, *Strongly negligible sets in Frechet manifolds*, Bull. Amer. Math. Soc., **75** (1969), 64-67.
3. R. Geoghegan, *On spaces of homeomorphisms, embeddings, and functions I*, Topology, **11** (1972), 159-177.

4. R. Geoghegan and D. W. Henderson, *Stable function spaces*, to appear.
5. O. Hanner, *Some Theorems on Absolute Neighborhood Retracts*, Arkiv Fur Matematik, **1** (1951), 389-408.
6. W. Haver, *Homeomorphisms and UV^∞ maps*, Proc. of SUNY Bingham. Conf. on Monotone Mappings and Open Mappings, (1970), 112-121.
7. ———, *The Closure of the Space of Homeomorphisms on a Manifold*, Trans. Amer. Math. Soc., to appear.
8. J. Hocking, *Approximations to monotone mappings on noncompact two-dimensional manifolds*, Duke Math. J., **21** (1954), 639-651.
9. W. K. Mason, *The space of all self-homeomorphisms of a 2-cell which fix the cell's boundary is an absolute retract*, Trans. Amer. Math. Soc., **161** (1971), 185-206.
10. L. C. Siebenmann, *Approximating cellular maps by homeomorphisms*, Topology, **11** (1972), 271-294.

Received September 6, 1972. Research partially supported by NSF Grant GP. 33872.

UNIVERSITY OF TENNESSEE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$60.00 a year (6 Vols., 12 issues). Special rate: \$30.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

| | |
|--|-----|
| Mustafa Agah Akcoglu, John Philip Huneke and Hermann Rost, <i>A counter example to the Blum Hanson theorem in general spaces</i> | 305 |
| Huzihiro Araki, <i>Some properties of modular conjugation operator of von Neumann algebras and a non-commutative Radon-Nikodym theorem with a chain rule</i> | 309 |
| E. F. Beckenbach, Fook H. Eng and Richard Edward Tafel, <i>Global properties of rational and logarithmico-rational minimal surfaces</i> | 355 |
| David W. Boyd, <i>A new class of infinite sphere packings</i> | 383 |
| K. G. Choo, <i>Whitehead Groups of twisted free associative algebras</i> | 399 |
| Charles Kam-Tai Chui and Milton N. Parnes, <i>Limit sets of power series outside the circles of convergence</i> | 403 |
| Allan Clark and John Harwood Ewing, <i>The realization of polynomial algebras as cohomology rings</i> | 425 |
| Dennis Garbanati, <i>Classes of circulants over the p-adic and rational integers</i> | 435 |
| Arjun K. Gupta, <i>On a "square" functional equation</i> | 449 |
| David James Hallenbeck and Thomas Harold MacGregor, <i>Subordination and extreme-point theory</i> | 455 |
| Douglas Harris, <i>The local compactness of vX</i> | 469 |
| William Emery Haver, <i>Monotone mappings of a two-disk onto itself which fix the disk's boundary can be canonically approximated by homeomorphisms</i> | 477 |
| Norman Peter Herzberg, <i>On a problem of Hurwitz</i> | 485 |
| Chin-Shui Hsu, <i>A class of Abelian groups closed under direct limits and subgroups formation</i> | 495 |
| Bjarni Jónsson and Thomas Paul Whaley, <i>Congruence relations and multiplicity types of algebras</i> | 505 |
| Lowell Duane Loveland, <i>Vertically countable spheres and their wild sets</i> | 521 |
| Nimrod Megiddo, <i>Kernels of compound games with simple components</i> | 531 |
| Russell L. Merris, <i>An identity for matrix functions</i> | 557 |
| E. O. Milton, <i>Fourier transforms of odd and even tempered distributions</i> | 563 |
| Dix Hayes Pettey, <i>One-one-mappings onto locally connected generalized continua</i> | 573 |
| Mark Bernard Ramras, <i>Orders with finite global dimension</i> | 583 |
| Doron Ravdin, <i>Various types of local homogeneity</i> | 589 |
| George Michael Reed, <i>On metrizable complete Moore spaces</i> | 595 |
| Charles Small, <i>Normal bases for quadratic extensions</i> | 601 |
| Philip C. Tonne, <i>Polynomials and Hausdorff matrices</i> | 613 |
| Robert Earl Weber, <i>The range of a derivation and ideals</i> | 617 |