

Pacific Journal of Mathematics

VERTICALLY COUNTABLE SPHERES AND THEIR WILD SETS

LOWELL DUANE LOVELAND

VERTICALLY COUNTABLE SPHERES AND THEIR WILD SETS

L. D. LOVELAND

A 2-sphere S in E^3 is said to have vertical order n if the intersection of each vertical line with S contains no more than n points. It is shown that $S \cup \text{Int } S$ is a 3-cell that is locally tame from $\text{Ext } S$ modulo a 0-dimensional set if S has vertical order 5. A subset X of E^3 is said to have countable (finite) vertical order if the intersection of X with each vertical line consists of countably (finitely) many points. A 2-sphere in E^3 with countable vertical order can have a wild set of dimension no larger than one.

For each 2-sphere S in E^3 there is a homeomorphism $h: E^3 \rightarrow E^3$ such that each vertical line intersecting $h(S)$ does so in a 0-dimensional set [2, Theorem 10.1]; thus the condition that a 2-sphere be "vertically 0-dimensional" imposes no restriction on the wildness of the 2-sphere. A study of vertically finite 2-spheres (spheres with finite vertical order) was begun in [10] where it was proven that a 2-sphere in E^3 having vertical order 3 is tame. Even though there are wild 2-spheres having vertical order 4, it is known that $S \cup \text{Int } S$ is a 3-cell if S has vertical order 5 [11]. We extend this result to show that the set $W(S)$ of points where the 2-sphere S fails to be locally tame must be 0-dimensional if S has vertical order 5. An example is given at the end of the paper to show that 5 is the largest integer for which this result is true. We also show that the wildness of a vertically countable sphere is limited to a 1-dimensional set.

In the remainder of the paper we use $\pi: E^3 \rightarrow E^2$ to denote the vertical projection of E^3 onto the horizontal plane E^2 . For convenience, we always assume that E^2 is located vertically below the sphere or cube under investigation. We use $L(x)$ to denote the vertical line containing the point x .

A vertical line L is said to pierce a subdisk D of a 2-sphere S if there is an interval I in L such that $I \cap S$ is a point $p \in D$ and I intersects both $\text{Int } S$ and $\text{Ext } S$. We say that L links the boundary $\text{Bd } D$ of a disk D if L intersects every disk bounded by $\text{Bd } D$.

2. Spheres having countable vertical order.

THEOREM 2.1. *If S is a 2-sphere in E^3 having countable vertical order, then $W(S)$ contains no open subset of S .*

Proof. Suppose that $W(S)$ contains a disk D in S . We shall

produce a contradiction by exhibiting a vertical line L whose intersection with D contains a Cantor set.

Assertion A. If D' is a subdisk of D , then there is an open subset U of E^3 such that $\pi(U) \subset \pi(D')$.

To prove Assertion A it suffices to show that $\pi(D')$ is not one-dimensional. This follows from [9, Theorem VI.7, p. 91] since the map $\pi \upharpoonright D': D' \rightarrow \pi(D')$ is closed.

Assertion B. If D' is a subdisk of D and U is an open subset of E^3 such that $\pi(U) \subset \pi(D')$, then there exist disjoint disks D_1 and D_2 in D' and an open subset N of U such that each vertical line through $\text{cl}(N)$ intersects both D_1 and D_2 .

In order to select the disks D_i in Assertion B we first show the existence of a vertical line L containing two points r and t in D' and containing two sequences $\{u_i\}$ and $\{l_i\}$ of points such that

- (1) $\{u_i\}$ converges to r from above,
- (2) $\{l_i\}$ converges to r from below,
- (3) there is a component V_1 of $E^3 - S$ containing every u_i , and
- (4) $E^3 - (S \cup V_1) = V_2$ contains every l_i .

Notice that some vertical line L' intersects D' in more than two points [7, Theorem 2.3], so we may choose two points r' and t' in $L' \cap D'$. Let B be an open ball centered at r' such that $B \cap S \subset D'$. If r' does not satisfy the four conditions above relative to L' , it must be because some interval I in $L' \cap B$ has r' as its midpoint and lies, except for r' , in a single component, say V_1 , of $E^3 - S$. Let B_1 and B_2 be disjoint round open balls of equal radius centered at points of L' above and below r' , respectively such that $B_1 \cup B_2 \subset V_1 \cap B$. Now close to r' and vertically between B_1 and B_2 , there must exist a point e of V_2 . Then $L = L(e)$ intersects V_2 between its two intersections with $V_1 \cap (B_1 \cup B_2)$, so L intersects D' at least twice. Let r be the lowest point of the component of $L \cap (S \cup V_1)$ containing $L \cap B_1$, and choose t to be some other point of $L \cap S$. Since S has countable vertical order it is clear that r is a limit point of $L \cap V_1$ from above and of $L \cap V_2$ from below. Thus conditions (1), (2), (3), and (4) are satisfied.

Choose a disk D_1 in D' such that $r \in \text{Int } D_1$ and $t \notin D_1$. We claim that there is an open set U_1 containing r such that every vertical line through U_1 intersects D_1 . Suppose there is no such open set, and for each i let E_i be a horizontal disk centered at l_i and lying in V_2 . There must be a sequence $\{x_i\}$ such that $x_i \in E_i$, for each i , no $L(x_i)$ intersects D_1 , and $\{L(x_i)\}$ converges to $L(r)$. For each i let y_i be the first point of S above x_i on $L(x_i)$ (such a point will exist for suf-

ficiently large integers i since u_i and l_i are different components of $E^3 - S$), and let I_i be the vertical interval $[x_i, y_i]$ in $S \cup V_2$. Since some subsequence of $\{y_i\}$ converges, we assume for notational convenience that $\{y_i\}$ converges to a point y . Of course $y \in L(r) \cap S$. It is clear that y is not above r on $L(r)$ because $\{r, y\} \subset \liminf I_i \subset S \cup V_2$ whereas $\{u_i\} \rightarrow r$ and $u_i \in V_1$. Nor is y below r on $L(r)$ because $\{l_i\} \rightarrow r$, $\{l_i, x_i\} \subset E_i$, and x_i lies vertically below y_i . Thus $\{y_i\}$ converges to r , and we have the contradiction that most of the y_i 's must belong to D' while $L(y_i) \cap D'$ was supposed to be empty. The existence of U_1 is established.

Now choose a disk D_2 such that $D_1 \cap D_2 = \emptyset$, $t \in \text{Int } D_2$, $D_2 \subset D'$, and $\pi(D_2) \subset \pi(U_1)$. From Assertion A there is an open set U_2 such that every vertical line through U_2 intersects D_2 . Such a line will also intersect U_1 and hence D_1 . Choose N to be any open subset of U such that $\pi(\text{cl}(N) \subset \pi(U_1) \cap \pi(U_2))$.

Now that the two assertions have been proven it might be clear how to proceed inductively to produce a vertical line containing uncountably many points of S ; nevertheless, we give a brief outline. From Assertion A there is an open set U such that every vertical line through U intersects D . Now we apply Assertion B to obtain an open set U_1 , whose closure lies in U , and two disjoint disks D_1 and D_2 in D such that every vertical line through $\text{cl}(U_1)$ intersects both D_1 and D_2 . This ends the first step in the construction. Assertion B can now be applied to D_1 to obtain two disjoint disks D_{11} and D_{12} in D_1 and an open set N_1 such that vertical lines through $\text{cl}(N_1)$ intersect both D_{11} and D_{12} . Now B is applied to D_2 and N_1 so that at the completion of step 2 we have an open set U_2 whose closure lies in U_1 and four disjoint disks D_{11}, D_{12}, D_{21} , and D_{22} in D where each vertical line through $\text{cl}(U_2)$ intersects each of the four disks. When the construction is finished it is clear that a vertical line through $\bigcap_1^\infty \text{cl}(U_i)$ will intersect each of the 2^n disks at the n th step. Thus such a line contains an uncountable set of points of S . This contradiction establishes the theorem.

COROLLARY 2.2. *If S is a 2-sphere in E^3 having countable vertical order, then S is locally tame modulo a 1-dimensional subset.*

3. Spheres of vertical order order 5. The following four lemmas are used to establish the main result (Theorem 3.5).

LEMMA 3.1. *If S has vertical 5, then S is locally tame at each point of S that is vertically above or below a point of $\text{Int } S$; that is, $\pi(\text{Int } S) \cap \pi(W(S)) = \emptyset$.*

Proof. Let p be a point of S such that $L(p) \cap \text{Int } S \neq \emptyset$. Thus

$L(p)$ must link the boundaries of each of two disjoint disks D_1 and D_2 in S . Let B be a ball lying in $\text{Int } S$ such that each vertical line through B links both $\text{Bd } D_1$ and $\text{Bd } D_2$. If $p \notin D_1 \cup D_2$, then there is a disk D_3 in S such that $p \in \text{Int } D_3$, $D_3 \cap (D_1 \cup D_2) = \emptyset$, and $\pi(D_3) \subset \pi(B)$. Then each vertical line intersecting D_3 also intersects both D_1 and D_2 . Since D has vertical order 5 it is clear that D_3 has vertical order 3. Thus D is locally tame at p [7, Theorem 2.3] and so is S .

We may now assume that $p \in \text{Int } D_1$. Let D'_1 be a subdisk of D_1 such that $\pi(D'_1) \subset \pi(B)$, and, for each $\xi > 0$, let X^ξ be the union of all vertical intervals of diameter no less than ξ in $S \cup \text{Int } S$ that intersect D'_1 . It is an exercise to see that X^ξ is closed, and it follows from [6, Theorem 5] that X^ξ is a $*$ -taming set. Now consider a point q in D'_1 but not in $X^{1/i}$ for any i . It follows that q lies in no vertical interval in $S \cup \text{Int } S$. Thus $L(q)$ does not pierce D'_1 at q , and $L(q)$ must pierce D'_1 at some other point t by the choice of B . Let D be a disk in D'_1 with t in its interior such that $q \notin D$ and $L(q)$ links $\text{Bd } D$. Then there is a disk D_q in $D'_1 - D$ such that $q \in \text{Int } D_q$ and each vertical line through D_q links $\text{Bd } D$. Thus such a line intersects both D and D_2 . This means that D_q has vertical order 3 and is tame [7, Theorem 2.3]. Now we see that each point of D'_1 either lies in the interior of a tame disk in D'_1 or lies in $\bigcup_1^\infty X^{1/i}$. Since a tame disk is a $*$ -taming set and a countable number of tame disks suffice to cover $D'_1 - \bigcup_1^\infty X^{1/i}$, we see that D'_1 lies in a $*$ -taming set of the form $(\bigcup_1^\infty X^{1/i}) \cup$ (a countable collection of tame disks) in $S \cup \text{Int } S$ [5, Theorem 3.7 and Corollary 3.8]. Thus S is locally tame at p from $E^3 - (S \cup \text{Int } S)$ by the definition of a $*$ -taming set. Since S is locally tame from $\text{Int } S$ [11], it follows that S is locally tame at p .

LEMMA 3.2. *If M is a continuum in $W(S)$ and S is a 2-sphere having vertical order 5, then M is tame.*

Proof. We may assume that M is nondegenerate since singleton sets always lie on tame spheres. From the previous lemma it is clear that $\pi(M) \subset \text{Bd } \pi(\text{Int } S)$. Let $U = \text{Int } S$ and let X be the component of $\text{Bd } \pi(U)$ containing $\pi(M)$. We shall show the existence of a space homeomorphism $H: E^3 \rightarrow E^3$ such that $\pi(H(M))$ is either an arc or a simple closed curve. Then $H(M)$ is clearly tame since it lies in $\pi^{-1}(\pi(H(M)))$.

The continuum X can be shown locally connected as in [7, Part 0.2]. Notice that $\pi(U)$ is open and connected. We let U' be the component of $E^2 - X$ containing $\pi(U)$ and for convenience in what follows we assume that U' is bounded. Notice that $\text{cl}(U') = X \cup U'$ since every point of S is accessible from $\text{Int } S$. Let $B^2 = \{(x, y) \mid x^2 + y^2 \leq 1\} \subset E^2$. There is a continuous function $f: B^2 \rightarrow \text{cl}(U')$ such that

$f|_{\text{Int } B^2}$ is a homeomorphism of $\text{Int } B^2$ onto U' and $f^{-1}(x)$ is a totally disconnected subset of $S^1 = \text{Bd } B^2$ for each $x \in X$ (see [12, p. 186]). Now we follow [7, §§ 2.1, 2.2, 2.3, and 2.4] to find a homeomorphism H of E^3 onto E^3 such that $\pi(H(\pi^{-1}(X) \cap S))$ is a simple closed curve. Thus $\pi(H(M))$ is either an arc or a simple closed curve since $\pi(H(M)) \subset \pi(H(\pi^{-1}(X) \cap S))$.

In the case where U' is not bounded the map f above takes $E^2 - \text{Int } B^2$ onto $\text{cl}(U')$ and causes some notational difficulties when we try to follow [7] as above. However, [7] still serves as an outline and we leave the details to the reader.

LEMMA 3.3. *If M is a nondegenerate continuum in $W(S)$ and S is a 2-sphere having vertical order 5, then each point of M is a limit point of $W(S) - M$.*

Proof. Suppose some point $p \in M$ is not a limit point of $W(S) - M$, and choose a disk D on S such that $p \in \text{Int } D$, $\text{Bd } D$ is tame [3], and $D \cap W(S) \subset M$. Let $X = M \cup (\text{Bd } D)$, and let S' be a 2-sphere containing $M \cup D$ that is locally tame modulo X [1]. From Lemma 3.2 we see that X is a taming set [4, Theorem 8.1.6, p. 320]. Thus S' is tame. This is a contradiction and the result follows.

LEMMA 3.4. *If D is a disk in a 2-sphere S , S has vertical order 5, $p \in \text{Int } D$, and V is an open subset of E^3 such that $p \in V$ and, for each vertical line L piercing D at a point in V , $L \cap \text{Int } S$ has exactly one component whose closure intersects D , then D is locally tame at p .*

Proof. If $L(p)$ intersects $\text{Int } S$, then the conclusion of Lemma 3.4 follows from Lemma 3.1. Thus we now assume $L(p) \cap \text{Int } S = \emptyset$. Choose a 2-sphere H in the shape of a right circular cylinder such that $p \in \text{Int } H$, $H \cap S \subset D$, $\text{Bd } D \subset \text{Ext } H$, $[L \cap (\text{Int } H)] \cap S = \{p\}$, the top and bottom disks T and D of H lie in $\text{Ext } S$, and each vertical line intersecting H also intersects V .

Let X be a component of $(\text{Int } S) \cap H$, and let $K = \text{Bd } X$. We shall show that $X \cup K$ is a disk by showing that K is a simple closed curve. To show that K is connected it suffices to prove that each simple closed curve J in X bounds a disk in X . Such a curve J cannot be essential on the annulus $H - D \cup T$ since J would link $L(p)$ while $L(p) \subset (\text{Ext } S) \cup S$ and $J \subset \text{Int } S$. Thus J must bound a disk E in $H - D \cup T$. From the hypothesis of Lemma 3.4 it is clear that $E \subset X$. Thus K is connected. The fact that K has vertical order 5 insures that K is arcwise accessible from both its complementary domains in H , and this implies that K is a simple closed curve.

Thus the closure of each component of $(\text{Int } S) \cap H$ is a spanning

disk for the 3-cell $C = S \cup \text{Int } S$. There can be at most a countable collection $\{D_1, D_2, \dots\}$ of these spanning disks since their interiors are pairwise disjoint. The fact that D has vertical order 5 insures that $\{D_i\}$ is a null sequence. We use these spanning disks to construct a 2-sphere S' containing p and lying in $D \cup (\bigcup_1^\infty D_i)$ and in $H \cup \text{Int } H$. From the hypothesis on D we see that the interior of S' is vertically connected; thus S' is tame [7, Main Theorem]. This means that D is locally tame at p .

THEOREM 3.5. *If a 2-sphere S in E^3 has vertical order 5, then $S \cup \text{Int } S$ is a 3-cell and S is locally tame from $\text{Ext } S$ modulo a 0-dimensional set.*

Proof. That $C = S \cup \text{Int } S$ is a 3-cell follows from [11]. It remains to show that the set W of wild points of S is 0-dimensional. Suppose to the contrary that there is a nondegenerate continuum M lying in W . Since C is a 3-cell there is an embedding $g: M \times [0, 1] \rightarrow C$ such that $G = g(M \times [0, 1]) \subset \text{Int } S$ and $g(m, 0) = m$ for every $m \in M$. We let $F = g(M \times [0, 1])$, and we note that it follows from Lemma 3.1 that $\pi(M)$ lies in the boundary of $\pi(F)$ in E^2 . For the same reason, $\pi(G) \cap \pi(M) = \emptyset$. Let U be a disk in E^2 and let p' be a point of $\text{Int } U$ such that $U \cap (\pi(\text{Bd } F)) \subset \pi(M)$ and $p' \in \pi(M)$. Choose a point p in $M \cap \pi^{-1}(p')$. In the next paragraph we show the existence of a disk E in S with $p \in \text{Int } E$ and $\pi(E) \subset U \cap \pi(F)$.

The difficulty in choosing E is the requirement that $\pi(E) \subset \pi(F)$. If no such E exists there must exist a sequence $\{p_i\}$ of points of $\text{Int } S$ converging to p such that $\pi(p_i) \in U - \pi(F)$ for each i . Using the 0-ULC of $\text{Int } S$ it is easy to select a point $g \in G \subset \text{Int } S$ close enough to p and an integer N large enough that g and p_N are the end points of an arc A in $\text{Int } S$ where $\pi(A) \subset U$. Now $\pi(A)$ contains an arc with one end point a in $\pi(G)$ and the other end point b in $U - \pi(F)$. If this arc is traversed from b to a , then there is a first point f of $\pi(F)$ encountered. This point f clearly belongs to $\text{Bd } \pi(F)$. This contradiction establishes the existence of E .

Now that the existence of E is clear we proceed by using Lemma 3.3 to pick a point q in $E \cap (W - M)$. Let V be an open ball centered at q such that $V \cap S \subset E$ and $V \cap F = \emptyset$. Since $L(q) \cap \text{Int } S = \emptyset$ (see Lemma 3.1) there are open balls B_1 and B_2 centered at points above and below q , respectively, that lie in $(\text{Ext } S) \cap V$. We choose a disk D in $V \cap S$ with $q \in \text{Int } D$ vertically between B_1 and B_2 such that $\pi(D) \subset \pi(B_1) \cap \pi(B_2)$. We shall show that D is locally tame at q to obtain a contradiction to $q \in W$.

In order to apply Lemma 3.4 we must show that if a vertical line L pierces D at a point of V , then $L \cap \text{Int } S$ has exactly one

component whose closure intersects D . Suppose to the contrary that for some such line L there are two components X and Y of $L \cap \text{Int } S$ whose closures intersect D . Now $X \cup Y \subset V$ since D lies between B_1 and B_2 . Since $L \cap \text{Int } S = \emptyset$ and $\pi(D) \subset \pi(F)$, we see that $L \cap G \neq \emptyset$. Thus $L \cap (\text{Int } S)$ has a third component Z , different from both X and Y because Z lies either above B_1 or below B_2 . Now the only way to avoid there being 6 points in $L \cap S$ is for X and Y to share an end point x . In this case there is a point e of $\text{Ext } S$ close enough to x to insure that there are three components of $L(e) \cap \text{Int } S$ with pairwise disjoint closures. Now $L(e) \cap S$ contains 6 points contrary to the hypothesis.

4. Examples and questions. One can use a countably infinite null sequence of Fox-Artin [8] "feelers" whose wild points form a dense subset of an arc to see that a vertically countable 2-sphere can have an arc in its wild set. Thus Corollary 2.2 cannot be improved in this direction.

EXAMPLE 4.1. A wild 2-sphere S having vertical order 6 such that $W(S)$ is not 0-dimensional. In Figure 1 we see an embedding of




FIGURE 1.

the Alexander Horned Sphere, having vertical order 4, inside a wedge-shaped 3-cell in E^3 . We attach a null sequence of such wedges to a right circular cone, as indicated in Figure 2, to obtain the desired example S . Notice that $W(S)$ is the union of a tame simple closed curve with countably infinite number of tame Cantor sets. Furthermore, every point of S is a piercing point of S .

In Example 4.1 we see that every nondegenerate continuum in $W(S)$ is tame.




FIGURE 2.

Question 4.2. If S is a 2-sphere in E^3 having finite vertical order, then must every nondegenerate continuum in $W(S)$ be tame?

We do not know the answer to Question 4.2 even when “vertical order n ” replaces “finite vertical order”, unless $n \leq 5$ where Theorem 3.5 applies. The proof of Lemma 3.2 shows an affirmative answer to Question 4.2 if it is also known that $\pi(W(S)) \cap \pi(\text{Int } S) = \emptyset$.

REFERENCES

1. R. H. Bing, *Approximating surfaces with polyhedral ones*, Ann. of Math., (2) **65** (1957), 456-483.
2. ———, *Conditions under which a surface in E^3 is tame*, Fund. Math., **47** (1959), 105-139.
3. ———, *Each disk in E^3 contains a tame arc*, Amer. J. Math., **84** (1962), 583-590.
4. C. E. Burgess and J. W. Cannon, *Embeddings of surfaces in E^3* , Rocky Mountain J. Math., **1** (1971), 259-344.
5. J. W. Cannon, **-taming sets for crumpled cubes, I: Basic properties*, Tran. Amer. Math. Soc., **161** (1971), 429-440.
6. ———, **-taming sets for crumpled cubes, II: Horizontal sections in closed sets*, Trans. Amer. Math. Soc., **161** (1971), 441-446.
7. J. W. Cannon and L. D. Loveland, *A 2-sphere in E^3 with vertically connected interior is tame*, Trans. Amer. Math. Soc., (to appear).
8. R. H. Fox and E. Artin, *Some wild cells and spheres in three-dimensional space*, Ann. of Math., (2) **49** (1948), 979-990.
9. W. Hurewicz and H. Wallman, *Dimension Theory*, Princeton Math. Series, Vol. 4, Princeton Univ. Press, Princeton, N. J., 1941.
10. R. A. Jensen and L. D. Loveland, *Surfaces of vertical order 3 are tame*, Bull. Amer. Math. Soc., **76** (1970), 151-154.
11. L. D. Loveland, *A 2-sphere of vertical order 5 bounds a 3-cell*, Proc. Amer. Math. Soc., **26** (1970), 674-678.

12. M. H. A. Newman, *Elements of the Topology of Plane Sets of Points*, First Edition, Cambridge Univ. Press, Cambridge, 1937.

Received November 21, 1972.

UTAH STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$60.00 a year (6 Vols., 12 issues). Special rate: \$30.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Mustafa Agah Akcoglu, John Philip Huneke and Hermann Rost, <i>A counter example to the Blum Hanson theorem in general spaces</i>	305
Huzihiro Araki, <i>Some properties of modular conjugation operator of von Neumann algebras and a non-commutative Radon-Nikodym theorem with a chain rule</i>	309
E. F. Beckenbach, Fook H. Eng and Richard Edward Tafel, <i>Global properties of rational and logarithmico-rational minimal surfaces</i>	355
David W. Boyd, <i>A new class of infinite sphere packings</i>	383
K. G. Choo, <i>Whitehead Groups of twisted free associative algebras</i>	399
Charles Kam-Tai Chui and Milton N. Parnes, <i>Limit sets of power series outside the circles of convergence</i>	403
Allan Clark and John Harwood Ewing, <i>The realization of polynomial algebras as cohomology rings</i>	425
Dennis Garbanati, <i>Classes of circulants over the p-adic and rational integers</i>	435
Arjun K. Gupta, <i>On a "square" functional equation</i>	449
David James Hallenbeck and Thomas Harold MacGregor, <i>Subordination and extreme-point theory</i>	455
Douglas Harris, <i>The local compactness of vX</i>	469
William Emery Haver, <i>Monotone mappings of a two-disk onto itself which fix the disk's boundary can be canonically approximated by homeomorphisms</i>	477
Norman Peter Herzberg, <i>On a problem of Hurwitz</i>	485
Chin-Shui Hsu, <i>A class of Abelian groups closed under direct limits and subgroups formation</i>	495
Bjarni Jónsson and Thomas Paul Whaley, <i>Congruence relations and multiplicity types of algebras</i>	505
Lowell Duane Loveland, <i>Vertically countable spheres and their wild sets</i>	521
Nimrod Megiddo, <i>Kernels of compound games with simple components</i>	531
Russell L. Merris, <i>An identity for matrix functions</i>	557
E. O. Milton, <i>Fourier transforms of odd and even tempered distributions</i>	563
Dix Hayes Pettey, <i>One-one-mappings onto locally connected generalized continua</i>	573
Mark Bernard Ramras, <i>Orders with finite global dimension</i>	583
Doron Ravdin, <i>Various types of local homogeneity</i>	589
George Michael Reed, <i>On metrizable complete Moore spaces</i>	595
Charles Small, <i>Normal bases for quadratic extensions</i>	601
Philip C. Tonne, <i>Polynomials and Hausdorff matrices</i>	613
Robert Earl Weber, <i>The range of a derivation and ideals</i>	617