

Pacific Journal of Mathematics

POLYNOMIALS AND HAUSDORFF MATRICES

PHILIP C. TONNE

POLYNOMIALS AND HAUSDORFF MATRICES

PHILIP C. TONNE

If f is a function from the rational numbers in $[0, 1]$ to the complex plane and c is a complex sequence, then the Hausdorff matrix $H(c)$ for c and a sequence $L(f, c)$ are defined:

$$H(c)_{np} = \binom{n}{p} \sum_{q=0}^{n-p} (-1)^q \binom{n-p}{q} c_{p+q} = \binom{n}{p} \Delta^{n-p} c_p$$

$$L(f, c)_n = \sum_{p=0}^n H(c)_{np} f(p/n).$$

THEOREM. If f is a function from the rationals in $[0, 1]$ to the plane and $L(f, c)$ converges for each complex sequence c , then f is a subset (contraction) of a polynomial.

J. S. MacNerney [2, p. 56] and A. Jakimovski [1] have shown:

THEOREM A. If f is a polynomial and c is a complex sequence then $L(f, c)$ converges.

Our theorem is a converse to Theorem A and an improvement of Theorem 7 of [3] where we proved that if f is a continuous function from $[0, 1]$ to the complex plane and $L(f, c)$ converges for each complex sequence c , then f is a subset of a polynomial.

LEMMA 1. Suppose that M is an infinite complex matrix and M transforms each sequence to a bounded sequence. Then there is a positive integer P such that if p is an integer exceeding P then $M_{np} = 0$, $n = 0, 1, \dots$.

Lemma 2, which follows, is not necessary for our result. It is included because it gives an interesting way of proving Lemma 3. The proof for Lemma 3 given here is due to the referee. It is simpler and more direct than the author's argument which uses Lemma 2. The author is grateful to the referee for the pretty proof of Lemma 3 which appears in this paper.

LEMMA 2. Suppose that n is a positive integer, each of $\{a_p\}_1^n$ and $\{b_p\}_1^n$ is an increasing sequence of nonnegative integers, M is an $n \times n$ matrix such that

$$M_{mp} = \begin{pmatrix} a_m \\ b_p \end{pmatrix} \quad (m, p = 1, 2, \dots, n)$$

and $a_k \geq b_k$, $k = 1, 2, \dots, n$. Then the determinant of M is positive.

LEMMA 3. Suppose that f is a function from the rationals in $[0, 1]$ to the complex plane, P is a nonnegative integer, and

$$\sum_{q=0}^p (-1)^q \binom{p}{q} f(q/n) = 0$$

for each integer-pair $\{n, p\}$ satisfying $P < p \leq n$. Then f is a subset of a polynomial.

Lemma 1 is known. Also, its proof is not difficult.

Lemma 2 is Theorem 2 of [4]. (This present paper was the motivation for [4].)

Proof of Lemma 3. Suppose that each of p and n is an integer and $P < p \leq n$. For each number x let $Q(x)$ be

$$\sum_{k=0}^P \binom{x}{k} \sum_{q=0}^k (-1)^{k+q} \binom{k}{q} f(q/n)$$

(here $\binom{x}{0} = 1$ and $\binom{x}{k+1} = (x-k)/(k+1) \binom{x}{k}$, $k = 0, 1, 2, \dots$), and let $H_n(x)$ be $Q(nx)$. H_n is a polynomial of degree P , at most, and

$$H_n(j/n) = f(j/n) \quad (j = 0, 1, \dots, n).$$

Now, if m is a multiple of n , $H_m = H_n$, since each of H_m and H_n is a polynomial of degree P , at most, $P < n$, and H_m and H_n agree at $0, 1/n, 2/n, \dots, n/n$.

Consequently, if n and m are integers exceeding P , $H_n = H_{m \cdot n} = H_m$. Now, let n be $P + 1$ and let s be a positive integer and let r be an integer in $[0, s]$. Then

$$H_n\left(\frac{r}{s}\right) = H_n\left(\frac{rn}{sn}\right) = H_{sn}\left(\frac{rn}{sn}\right) = f\left(\frac{rn}{sn}\right) = f\left(\frac{r}{s}\right),$$

and f is a subset of the polynomial H_n .

Proof of Theorem. Let c be a complex sequence. For each positive integer n ,

$$L(f, c)_n = \sum_{p=0}^n c_p \binom{n}{p} \sum_{q=0}^p (-1)^{p+q} \binom{p}{q} f(q/n).$$

Define

$$M_{n,p} = \binom{n}{p} \sum_{q=0}^p (-1)^{p+q} \binom{p}{q} f(q/n) \quad (n = 1, 2, \dots; p = 0, 1, \dots, n)$$

and $M_{np} = 0$ if $n = 0$ or $p > n$. By hypothesis M transforms each sequence to a convergent sequence, so, by Lemma 1, there is a positive integer P such that if p is an integer exceeding P then $M_{np} = 0$, $n = 0, 1, \dots$.

Thus, by Lemma 3, f is a subset of a polynomial.

REFERENCES

1. A. Jakimovski, *Some remarks on the moment problem of Hausdorff*, J. London Math. Soc., **33** (1958), 1-13.
2. J. S. Mac Nerney, *Hermitian moment sequences*, Trans. Amer. Soc., **103** (1962), 45-81.
3. Philip C. Tonne, *Power-series and Hausdorff matrices*, Pacific J. Math., **21** (1967), 189-198.
4. ———, *A regular determinant of binomial coefficients*, Proc. Amer. Math. Soc., **41** (1973), 17-23.

Received October 12, 1972 and in revised form February 26, 1973.

EMORY UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI*

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$60.00 a year (6 Vols., 12 issues). Special rate: \$30.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 50, No. 2

October, 1974

Mustafa Agah Akcoglu, John Philip Huneke and Hermann Rost, <i>A counter example to the Blum Hanson theorem in general spaces</i>	305
Huzihiro Araki, <i>Some properties of modular conjugation operator of von Neumann algebras and a non-commutative Radon-Nikodym theorem with a chain rule</i>	309
E. F. Beckenbach, Fook H. Eng and Richard Edward Tafel, <i>Global properties of rational and logarithmico-rational minimal surfaces</i>	355
David W. Boyd, <i>A new class of infinite sphere packings</i>	383
K. G. Choo, <i>Whitehead Groups of twisted free associative algebras</i>	399
Charles Kam-Tai Chui and Milton N. Parnes, <i>Limit sets of power series outside the circles of convergence</i>	403
Allan Clark and John Harwood Ewing, <i>The realization of polynomial algebras as cohomology rings</i>	425
Dennis Garbanati, <i>Classes of circulants over the p-adic and rational integers</i>	435
Arjun K. Gupta, <i>On a "square" functional equation</i>	449
David James Hallenbeck and Thomas Harold MacGregor, <i>Subordination and extreme-point theory</i>	455
Douglas Harris, <i>The local compactness of vX</i>	469
William Emery Haver, <i>Monotone mappings of a two-disk onto itself which fix the disk's boundary can be canonically approximated by homeomorphisms</i>	477
Norman Peter Herzberg, <i>On a problem of Hurwitz</i>	485
Chin-Shui Hsu, <i>A class of Abelian groups closed under direct limits and subgroups formation</i>	495
Bjarni Jónsson and Thomas Paul Whaley, <i>Congruence relations and multiplicity types of algebras</i>	505
Lowell Duane Loveland, <i>Vertically countable spheres and their wild sets</i>	521
Nimrod Megiddo, <i>Kernels of compound games with simple components</i>	531
Russell L. Merris, <i>An identity for matrix functions</i>	557
E. O. Milton, <i>Fourier transforms of odd and even tempered distributions</i>	563
Dix Hayes Pettey, <i>One-one-mappings onto locally connected generalized continua</i>	573
Mark Bernard Ramras, <i>Orders with finite global dimension</i>	583
Doron Ravdin, <i>Various types of local homogeneity</i>	589
George Michael Reed, <i>On metrizability of complete Moore spaces</i>	595
Charles Small, <i>Normal bases for quadratic extensions</i>	601
Philip C. Tonne, <i>Polynomials and Hausdorff matrices</i>	613
Robert Earl Weber, <i>The range of a derivation and ideals</i>	617