# Pacific Journal of Mathematics

LIPSCHITZ SPACES

JERRY ALAN JOHNSON

Vol. 51, No. 1

November 1974

# LIPSCHITZ SPACES

# JERRY A. JOHNSON

If (S, d) is a metric space and  $0 < \alpha < 1$ , Lip  $(S, d^{\alpha})$  is the Banach space of real or complex-valued functions f on S such that  $||f|| = \max(||f||_{\infty}, ||f||_{d^{\alpha}}) < \infty$ , where  $||f||_{d^{\alpha}} = \sup\{|f(s) - f(t)|d^{-\alpha}(s, t): s \neq t\}$ . The closed subspace of functions f such that  $\lim_{d(s,t)\to 0} |f(s) - f(t)| d^{-\alpha}(s, t) = 0$  is denoted by lip  $(S, d^{\alpha})$ . The main result is that, when  $\inf_{s\neq t} d(s, t) = 0$  lip  $(S, d^{\alpha})$  contains a complemented subspace isomorphic with  $c_0$  and Lip (S, d)contains a subspace isomorphic with  $l_{\infty}$ . From the construction, it follows that lip  $(S, d^{\alpha})$  is not isomorphic to a dual space nor is it complemented in Lip  $(S, d^{\alpha})$ .

If E is a normed space,  $E^*$  denotes its dual.  $c_0$ ,  $l_1$ , and  $l_{\infty}$  denote the usual sequence spaces; "isomorphism" means "linear homeomorphism"; "projection" means "continuous projection"; and F is a complemented subspace of E if there is a projection of E on F.

In recent years, much work has been done on the Banach space properties (isomorphic and isometric) of Lipschitz functions. Some of the main references are [1], [2], [3], [4], [10], [11], and [12]. There are still many outstanding conjectures concerning these spaces, some of which seem to be fairly difficult; especially certain questions about extreme points. The known results along these lines can be found in [2], [3], [4], [10], and [11].

It is established in [1] that if (S, d) is an infinite compact subset of Euclidean space, then  $\lim (S, d^{\alpha})(0 < \alpha < 1)$  is isomorphic with  $c_0$ and  $\lim (S, d^{\alpha})$  is isomorphic with  $l_{\infty}$ . The proof given in [1] is not evidently adaptable to arbitrary compact metric spaces. Thus, many of the natural conjectures one might make concerning properties that Lipschitz spaces may share with these sequence spaces are still unresolved. Of course the main one is whether  $\lim (S, d^{\alpha})$  with  $0 < \alpha < 1$  and S compact and infinite is isomorphic with  $c_0$ .

It is shown in [1, Remark, p. 319] that for (S, d) compact and  $0 < \alpha < 1$ , lip  $(S, d^{\alpha})$  and Lip  $(S, d^{\alpha})$  are isomorphic to subspaces of  $c_0$  and  $l_{\infty}$  respectively. (Although the sketch of the proof given there is not precisely right, Professor Frampton, in a private communication, has exhibited a correct one for which we thank him.) It is well known that an infinite dimensional subspace A of  $c_0$  contains a subspace B isomorphic to  $c_0$  and that, since A is separable, B is complemented. In §1 of this paper, we show that lip  $(S, d^{\alpha})$ ,  $0 < \alpha < 1$ , contains a complemented subspace isomorphic to  $c_0$  when inf  $\{d(s, t): s \neq t\} = 0$  (Theorem 1). It is also shown that lip  $(S, d^{\alpha})$ 

remark here that if  $(\overline{S}, d)$  is the completion of (S, d) then the restriction map is an isometric isomorphism of Lip  $(\overline{S}, d^{\alpha})$  onto Lip  $(S, d^{\alpha})$  which sends lip  $(\overline{S}, d^{\alpha})$  onto lip  $(S, d^{\alpha})$   $(0 < \alpha \leq 1)$ .)

In addition, we show in Theorem 1 that for any infinite metric space (S, d) and  $0 < \alpha \leq 1$ , Lip  $(S, d^{\alpha})$  contains a subspace isomorphic to  $l_{\infty}$ .

In §2, we discuss some open problems concerning the isomorphic types of the Lipschitz spaces.

In § 3, we consider some questions about the extreme points of the unit ball and dual unit ball of these spaces.

1. In this section we prove our main theorems. We begin by stating

THEOREM 1. Let (S, d) be a metric space with  $\inf \{d(s, t) : s \neq t\} = 0$ . Then Lip (S, d) contains a subspace isomorphic to  $l_{\infty}$  and lip  $(S, d^{\alpha})$ ,  $0 < \alpha < 1$ , contains a complemented subspace isomorphic to  $c_0$ .

REMARK 1. Theorem 1 has been announced in [5] along with Corollaries 1 and 2 below. It is pointed out in [4, Lemma 2.5] that if  $\{d(s, t): s \neq t\} > 0$ , then both  $\lim (S, d^{\alpha})$  and  $\lim (S, d^{\alpha})$  are isomorphic with the bounded functions on S.

The next two corollaries are immediate. We assume (S, d) and  $\alpha$  are as in Theorem 1.

COROLLARY 1. lip  $(S, d^{\alpha})$  is not isomorphic with a dual space.

*Proof.* It has been observed in [7, p. 16] by Lindenstrauss (and is not hard to see) that if a Banach space is complemented in some dual space, it is complemented in its second dual. Hence, if lip  $(S, d^{\alpha})$  is isomorphic with a dual space,  $c_0$  is complemented in  $l_{\infty} \cdots a$  contradiction.

COROLLARY 2. lip  $(S, d^{\alpha})$  is not complemented in Lip  $(S, d^{\alpha})$ .

*Proof.* Let E and  $E_0$  denote the subspaces isomorphic with  $l_{\infty}$  and  $c_0$  respectively. In the proof of Theorem 1, E and  $E_0$  are constructed so that  $E_0 \subset E$ . Let P be a projection of  $\lim (S, d^{\alpha})$  on  $E_0$  and suppose Q is a projection of  $\lim (S, d^{\alpha})$  on  $\lim (S, d^{\alpha})$ . It is then easy to see that PQ restricted to E is a projection of E onto  $E_0$ . This is a contradiction.

REMARK 2. The theorem also implies the author's result [4, Theorem 2.6].

#### LIPSCHITZ SPACES

REMARK 3.  $l_{\infty}$  is complemented in any Banach space containing it. The following notation is used throughout:

$$(S, d)$$
 is a metric space, $ar{B}(s, r) = \{t \in S \mid d(s, t) \leq r\}$ 

and

$$\mathring{B}(s, r) = \{t \in S \mid d(s, t) < r\}$$

If  $A \subset S$  and  $B \subset S$ ,  $d(A, B) = \inf \{d(s, t) \mid s \in A, t \in B\}$ .  $d(A, \{t\})$  is denoted by d(A, t).

In [5], we sketched the proof of Theorem 1 in the case where S has no nonconstant Cauchy sequences. We present here the proof where S is assumed to have a limit point. (Recall that completeness may be assumed without loss of generality.) Since the case  $\alpha = 1$  for Lip  $(S, d^{\alpha})$  is similar to the discrete case appearing in [5], we will omit it.

*Proof of the Theorem.* We begin by constructing a sequence of closed balls that will serve as supports for certain functions.

Let  $s_0$  be a fixed limit point of S. Choose  $s_1 \in S$  with  $0 < d(s_1, s_0)$ and define  $r_1 = (1/2)d(s_1, s_0)$ ,  $B_1 = \overline{B}(s_1, r_1)$  and  $p_1 = d(s_0, B_1)$ . Now, assume that  $s_j$   $(j \ge 1)$  has been chosen such that  $d(s_j, s_0) > 0$ . Set  $r_j = (1/2)d(s_j, s_0)$ ,  $B_j = \overline{B}(s_j, r_j)$  and  $p_j = d(s_0, B_j)$ . We first note that

$$(1) r_j \leq p_j \,.$$

*Proof.* Otherwise, there is a point  $t \in B_j$  with  $d(s_0, t) < r_j$ . Hence,  $d(s_0, s_j) \leq d(s_0, t) + d(t, s_j) < 2r_j$ , a contradiction.

Now,  $p_j > 0$  implies that there is a point  $s_{j+1}$  with  $0 < d(s_{j+1}, s_0) < (1/6)p_j$ . Set  $r_{j+1} = (1/2)d(s_{j+1}, s_0)$ ,  $B_{j+1} = \overline{B}(s_{j+1}, r_{j+1})$ , and  $p_{j+1} = d(B_{j+1}, s_0)$ .

We record the following facts concerning our construction which are needed later.

(2) 
$$p_j < \frac{1}{6} p_{j-1}$$
 for each  $j$ .

Proof.  $p_j \leq d(s_0, s_j) < (1/6)p_{j-1}$ .

(3) 
$$B_j \subset \overset{\circ}{B}\left(s_0, \frac{1}{2}p_{j-1}\right)$$
 for each  $j$ .

*Proof.* Let  $t \in B_j$ . Then  $d(t, s_0) \leq d(t, s_j) + d(s_j, s_0) \leq r_j + 2r_j \leq 3p_j < (1/2)p_{j-1}$  by (1) and (2).

*Proof.* If  $s \in B_i$ ,  $d(s, s_0) \ge p_i$ . If  $t \in B_j$ ,  $d(t, s_0) < (1/2)p_{j-1} \le (1/2)p_i$ by (3) and (2). Hence,  $d(s, t) \ge d(s, s_0) - d(s_0, t) \ge (1/2)p_i$ .

(5) 
$$d(s, \tilde{B}_j) \leq 3p_j \text{ for each } j \text{ and each } s \in S,$$
  
where  $\tilde{B}_j$  denotes the complement of  $B_j$ .

*Proof.* Assume  $s \in B_j$  since the assertion is otherwise trivial. Given  $\varepsilon > 0$ , there is  $t \in B_j$  such that  $d(t, s_0) < \varepsilon + d(B_j, s_0) = \varepsilon + p_j$ . Thus,  $d(s, \tilde{B}_j) \leq d(s, s_0) \leq d(s, s_j) + d(s_j, t) + d(t, s_0) \leq r_j + r_j + \varepsilon + p_j \leq 3p_j + \varepsilon$ , and the assertion follows.

If  $s \in B_i$  and  $t \in B_j$  with j > i, then

$$(6) \qquad \qquad \frac{d(s, \tilde{B}_i) + d(t, \tilde{B}_j)}{d(B_i, B_j)} < 7.$$

Proof. By (5) and (4),

$$rac{d(s,\,\widetilde{B}_i)+d(t,\,\widetilde{B}_j)}{d(B_i,\,B_j)} \leq rac{3p_i+3p_j}{(1/2)p_i} = 6 \Bigl[1+rac{p_j}{p_i}\Bigr]\,.$$

Since j > i,  $p_j < (1/6)p_{j-i} \leq (1/6)p_i$ . Hence, the assertion follows. We next proceed to construct the isomorphism. We will assume

in the proof that for each j,  $d(s, \tilde{B}_j) \leq 1$  for all  $s \in S$  and  $d(s, t) \leq 1$  for all  $s, t \in B_j$ . This clearly can be done by taking j large enough (see (5) and (1)).

First choose a sequence  $\{\beta_j\}$  converging to  $\alpha$  such that  $\alpha < \beta_j < 1$ ,  $r_j^{\beta_j-\alpha} \geq 1/2$ , and  $d^{\beta_j-\alpha}(s_j, \tilde{B}_j) \geq 1/2$  for each j. Now, define  $f_j(s) = d^{\beta_j}(s, \tilde{B}_j)$  for each j and  $s \in S$ . Then, given  $a = \{a_j\} \in l_{\infty}$ , define

$$f_a = \sum_{j=1}^{\infty} a_j f_j$$
.

It is easy to see that, since the nonzero functions  $f_j$  have disjoint supports  $B_j$ , the function  $f_a$  is well-defined and the mapping  $a \rightarrow f_a$  is one-to-one and linear.

Next, let us note that

$$||f_a|| \geq rac{1}{2^{1+lpha}}||a||$$

for each  $a \in l_{\infty}$ . To see this, observe that for each j,

$$egin{aligned} ||f_a||_{d^{lpha}} &\geq rac{|f_a(s_j) - f_a(s_0)|}{d^{lpha}(s_j,\,s_0)} = rac{|a_j|d^{eta_j}(s_j,\,\widetilde{B}_j)}{d^{lpha}(s_j,\,s_0)} \ &= rac{|a_j|d^{eta_j}(s_j,\,\widetilde{B}_j)}{(2r_j)^{lpha}} \geq rac{|a_j|r_j^{eta_j}}{2^{lpha}r_j^{lpha}} \geq rac{|a_j|}{2^{1+lpha}} \end{aligned}$$

by our choice of  $\beta_j$ . Since j was arbitrary, the desired inequality follows.

Now, we will show that  $a \to f_a$  is bounded and that  $f_a \in \text{lip}(S, d^{\alpha})$ when  $a \in c_0$ . The boundedness of  $a \to f_a$  will show that  $f_a$  is in fact in Lip  $(S, d^{\alpha})$ .

In what follows, assume that  $a \in l_{\infty}$  is fixed with  $|a_j| \leq 1$  for each j and set  $f = f_a$ . First note that

$$||f||_{\infty} = \sup_n ||a_n f_n||_{\infty} \leq \sup_n |a_n| \leq 1$$
 .

We next proceed to show that  $||f||_{d^{\alpha}} \leq 7^{\alpha} \cdot 2^{1-\alpha} \leq 7$ . Let  $s \in S$ ,  $t \in S$ ,  $s \neq t$ . If  $s \in B_{\epsilon}$  and

$$t 
ot\in igcup_{j=1}^\infty B_j$$
 ,

then

$$rac{|f(s)-f(t)|}{d^lpha(s,\,t)}=rac{|\,a_{\scriptscriptstyle lpha}\,|\,d^{eta_i}(s,\,\widetilde{B}_i)}{d^lpha(s,\,t)}\leq |\,a_{\scriptscriptstyle lpha}\,|\,d^{eta_i-lpha}(s,\,\widetilde{B}_i)\leq 1$$
 ,

while if  $t \in B_i$ , then

$$rac{|f(s)-f(t)|}{d^{lpha}(s,\,t)} \leq |\,a_i\,|\,d^{eta_i^{-lpha}}(s,\,t) \leq 1\;.$$

Thus, suppose  $s \in B_i$ ,  $t \in B_j$  and j > i. Then

$$egin{aligned} & rac{|f(s)-f(t)|}{d^lpha(s,t)} \leq rac{|f(s)|+|f(t)|}{d^lpha(s,t)} \leq rac{d^{eta i}(s,\, ilde{B}_i)+d^{eta j}(t,\, ilde{B}_j)}{d^lpha(B_i,\,B_j)} \ & \leq rac{d^{lpha}(s,\, ilde{B}_i)+d^lpha(t,\, ilde{B}_j)}{d^lpha(B_i,\,B_j)} \ , \end{aligned}$$

since  $\beta_i, \beta_j > \alpha$  and  $d(s, \tilde{B}_k) \leq 1$  for all k. Now, the last quotient does not exceed

$$2^{{\scriptscriptstyle 1-a}} \Big[ rac{d(s,\,\widetilde{B}_{\,\imath})\,+\,d(t,\,\widetilde{B}_{\,j})}{d(B_{_{\imath}},\,B_{\,j})} \Big]^a$$

since

$$rac{p^lpha+q^lpha}{2}\!\leq\!\left(rac{p+q}{2}
ight)^{\!lpha}$$

for  $p \ge 0$ ,  $q \ge 0$ . Hence, from (6) above,

$$\||f\||_{d^{lpha}} \leq 1 \lor 7^{lpha} \cdot 2^{\iota-lpha} \leq 7^{lpha} \cdot 2^{\iota-lpha} \leq 7$$
 .

Thus,  $\{f_a \mid a \in l_{\infty}\}$  is isomorphic with  $l_{\infty}$ .

Next, let  $a \in c_0$ ,  $||a|| \leq 1$ , and let  $\varepsilon > 0$  be given. We must find  $\delta > 0$  so that  $0 < d(s, t) < \delta$  implies that

$$rac{|f(s)-f(t)|}{d^lpha(s,\,t)} \leq arepsilon \; .$$

Since  $a \in c_0$ , there is a number N such that  $n \ge N$  implies  $|a_n| < (1/14)\varepsilon$ . There exists a  $\delta_0 > 0$  such that if  $i \ne j$  and  $d(B_i, B_j) < \delta_0$ , then  $i \ge N$  and  $j \ge N$ . This follows from (4) above. Having chosen N, take  $0 < \delta < \min \{\varepsilon^{1/\beta_j - \alpha} \mid 1 \le j < N\}$  and  $\delta < \delta_0$ . Then  $0 < d(B_i, B_j) < \delta$  still implies  $i, j \ge N$ .

Now, let  $0 < d(s, t) < \delta$ . Suppose  $s \in B_i$  and  $t \notin \bigcup_{j \neq i} B_j$ . If i < N, then

$$rac{|f(s)-f(t)\,|}{d^lpha(s,\,t)} \leq |\,a_i\,|\,d^{eta_i-lpha}(s,\,t) \leq \delta^{eta_i-lpha} \leq arepsilon \;.$$

If  $i \ge N$ ,  $|a_i| d^{\beta_i - \alpha}(s, t) \le |a_i| \delta^{\beta_i - \alpha} \le |a_i| < \varepsilon/14 < \varepsilon$ . Next, suppose  $s \in B_i$  and  $t \in B_j$  (j > i). Then

$$egin{aligned} rac{|f(s)-f(t)|}{d^lpha(s,t)} &\leq rac{|a_i|f_i(s)+|a_j|f_j(t)}{d^lpha(s,t)} \ &\leq (|a_i|+|a_j|) rac{d^{eta i}(s,\, ilde B_i)+d^{eta j}(t,\, ilde B_j)}{d^lpha(s,\,t)} \ &\leq \left(rac{arepsilon}{14}+rac{arepsilon}{14}
ight)\cdot 7=arepsilon \ ; \end{aligned}$$

this is because  $0 < d(s, t) < \delta$ ,  $s \in B_i$  and  $t \in B_j$  imply  $d(B_i, B_j) < \delta$ and hence  $i, j \ge N$ .

The only part now remaining is to show that " $c_0$ " is complemented in lip  $(S, d^{\alpha})$ . Our proof will entail the construction of a projection of Lip  $(S, d^{\alpha})$  onto the image of  $l_{\infty}$  which sends lip  $(S, d^{\alpha})$  onto the image of  $c_0$ . As mentioned before, since  $l_{\infty}$  is injective, it is already known that it must be complemented in Lip  $(S, d^{\alpha})$ .

Given  $f \in \text{Lip}(S, d^{\alpha})$ , define

$$Pf = \sum_{n} a_{n}f_{n}$$

where

$$a_n = \frac{f(s_n) - f(s_0)}{f_n(s_n)} \ .$$

It is easy to see that P is linear and that  $P^2 = P$ . Let  $||f|| \leq 1$ . We must first find a constant M such than  $|a_n| \leq M$  for each n.  $d(s_n, s_0) = 2r_n$ , for each n, by definition; thus,

$$d^lpha(s_n,\,s_0)=2^lpha r_n^lpha\leq 2^lpha d^lpha(s_n,\,ar{B}_n)\leq 2^{1+lpha}d^{eta_n}(s_n,\,ar{B}_n)$$

for each n, since  $\beta_n - \alpha$  was chosen small enough so that

$$d^{eta_n-lpha}(s_n,\,\widetilde{B}_n) \geqq rac{1}{2} \;.$$

Hence,

$$|f(s_n) - f(s_0)| \leq d^{lpha}(s_n, s_0) \leq 2^{1+lpha} d^{\beta_n}(s_n, \widetilde{B}_n) = 2^{1+lpha} f_n(s_n)$$

for each *n*. Therefore,  $|a_n| \leq 2^{1+\alpha}$  when  $||f|| \leq 1$ . Now, let  $f \in \text{lip}(S, d^{\alpha})$ . We must show that

$$\lim_{n o\infty}rac{|f(s_n)-f(s_0)|}{f_n(s_n)}=0 \;.$$

As above, we have  $d^{\alpha}(s_n, s_0) \leq 2^{1+\alpha} f_n(s_n)$ , so

$$\lim_{n \to \infty} \frac{|f(s_n) - f(s_0)|}{f_n(s_n)} \leq 2^{1+\alpha} \lim_{n \to \infty} \frac{|f(s_n) - f(s_0)|}{d^{\alpha}(s_n, s_0)} = 0 \, \, .$$

This completes the proof of the theorem.

We close this section with Theorem 2 which answers a question raised in [5].

THEOREM 2. The following are equivalent for  $0 < \alpha < 1$ .

(a) (S, d) is precompact.

(b)  $\lim (S, d^{\alpha})^*$  is separable.

(c)  $\lim (S, d^{\alpha})$  is separable.

*Proof.* In [3] Jenkins showed that if (S, d) is compact, the span of the point evaluations is dense in lip  $(S, d^{\alpha})^*$ . It is clear that  $|| \varepsilon_s - \varepsilon_t || \leq d^{\alpha}(s, t)$ , where  $\varepsilon_s(f) = f(s)$ . Thus,  $\{\varepsilon_s | s \in S\}$ , and hence lip  $(S, d^{\alpha})^*$ , is separable. (See [4] for further discussion.) Thus  $(a) \Rightarrow (b)$ .

(b)  $\Rightarrow$  (c) is true for any Banach space, so assume (a) fails. Then there exists a sequence  $\{s_n\} \subset S$  and a number p > 0 such that  $d(s_n, s_m) \geq p$  for each  $n \neq m$ . Let  $\{s_{n_k}\}$  be any subsequence of  $\{s_n\}$  and let  $A = \{s_{n_{2k+1}}\}$ . Now, the function f defined by  $f(s) = \min \{d(s, A), 1\}$ is an element of lip  $(S, d^{\alpha})$ , where  $d(s, A) = \inf \{d(s, t) \mid t \in A\}$ . However,  $f(s_{n_{2k}}) \geq \min (p, 1) > 0$  for each k, while  $f(s_{n_{2k+1}}) = 0$  for each k. Thus,  $\lim_k f(s_{n_k}) = \lim_k \varepsilon_{s_{n_k}}(f)$  does not exist; i.e.,  $\{\varepsilon_{s_n}\}$  has no weak\*-convergent subsequence. This implies that the dual unit ball of lip  $(S, d^{\alpha})$ is not  $w^*$ -metrizable, which completes the proof of the proposition by contradicting (c).

2. An investigation of the Banach space properties of the Lipschitz spaces is far from complete, and questions concerning those properties of  $(S, d^{\alpha})$  that give rise to corresponding properties of the Lipschitz spaces are abundant. The ultimate problem of classifying these spaces as to isomorphic type does not appear easy. Even the following two problems are still open:

If (S, d) is compact and infinite, and  $0 < \alpha < 1$ ,

(1) is  $\lim (S, d^{\alpha})$  isomorphic with  $c_0$  and

(2) is Lip  $(S, d^{\alpha})$  isomorphic with  $l_{\infty}$ ?

By [4, Theorem 4.7] it is known that Lip  $(S,d^{\alpha})$  is isometrically isomorphic with the bidual of lip  $(S, d^{\alpha})$ . Hence, a positive answer to (1) yields a positive answer to (2). As we mentioned in the introduction, the best result in this direction appears in [1].

One possible avenue of attack on question (2) may be furnished by the following proposition, since it seems that the problem of showing (b) or (c) may be more tractable than showing (e) directly. (For the definitions of  $\mathscr{L}_1$  and  $\mathscr{L}_{\infty}$  spaces see [9]. A Banach space is injective if it is complemented in every Banach space containing it.)

**PROPOSITION 1.** Let (S, d) be compact and infinite. If  $0 < \alpha < 1$ , the following assertions are equivalent.

- (a) Lip  $(S, d^{\alpha})$  is injective.
- (b) Lip (S,  $d^{\alpha}$ ) is an  $\mathscr{L}_{\infty}$  space.
- (c)  $\lim (S, d^{\alpha})$  is an  $\mathscr{L}_{\infty}$  space.
- (d) lip  $(S, d^{\alpha})^*$  is an  $\mathscr{L}_1$  space.
- (e) Lip  $(S, d^{\alpha})$  is isomorphic with  $l_{\infty}$ .

(a) and (b) are equivalent even if  $\alpha = 1$  and (S, d) is arbitrary.

*Proof.* [9, Remark 2, p. 337] yields (a)  $\Leftrightarrow$  (b) immediately since Lip (S, d) is a dual space for *any* metric space [4]. (a)  $\Leftrightarrow$  (c) is [9, Corollary, p. 335]. (d)  $\Leftrightarrow$  (b) is [9, Theorem I (iii), p. 327]. (d)  $\Leftrightarrow$  (e) is from the observation in [9, Problem 2a, p. 344] and the fact that lip  $(S, d^{\alpha})^*$  is separable (see [3]).

Although questions (1) and (2) are the most important, the following questions are also open in general.

(3) Does Lip (S, d) have the approximation property?

(4) If (S, d) is compact and  $0 < \alpha < 1$ , do lip  $(S, d^{\alpha})^*$  and lip  $(S, d^{\alpha})$  have Schauder bases?

Let us remark that in [3] it was shown that  $\lim(S, d^{\alpha})^*$  is separable.

Added in proof: By Enflo's example, there is a (non-compact) metric space for which  $\operatorname{Lip}(S, d)$  fails the approximation property. Using an idea due to Lindenstrauss it can be shown that  $\operatorname{Lip}(S, d)$  is not injective if (S, d) is the Hilbert cube.

3. In addition to the questions in §2 concerning the isomorphism types of the Lipschitz spaces, there are some interesting problems dealing with the extreme points of their unit balls and dual unit balls.

We begin by stating a theorem due to Lindenstrauss and Phelps

[8, Theorem 3.1]:

(I) If E is a normed space whose dual unit ball has countably many extreme points, then  $E^*$  is separable and E contains no infinite dimensional reflexive subspaces.

Quite recently William Johnson and Haskell Rosenthal [6] proved:

(II) If E is an infinite dimensional Banach space with  $E^{**}$  separable, then E and  $E^*$  have infinite dimensional reflexive subspaces.

(The author would like to thank Professors Johnson and Rosenthal for access to a preprint of [6].)

In view of (I), (II) now has as an immediate corollary the following: (III) The unit ball of  $E^{**}$ , for any infinite dimensional Banach space, has uncountably many extreme points.

The aforementioned results have some immediate applications to Lipschitz spaces. We proceed to mention a few.

Since it is known that Lip  $(S, d^{\alpha})$  is a second dual space for  $0 < \alpha < 1$  (see [3] and [4]), it follows from (III) that its unit ball has uncountably many extreme points. In view of (I), Theorem 1 and the fact [4, Theorem 4.1] that Lip (S, d) is a dual space for any metric space, we can state the following:

**PROPOSITION 2.** If (S, d) is any metric space with S infinite, then the unit ball of Lip (S, d) has uncountable many extreme points.

Of course, since Lip (S, d) is a dual space, its unit ball is the  $w^*$ -closed convex hull of its extreme points. As shown in [4, Corollary 4.4], convergence of bounded nets in the  $w^*$ -topology coincides with pointwise convergence in general, and with uniform convergence when (S, d) is compact. Thus, in both senses, the unit ball of Lip (S, d) has many extreme points. The problem of characterizing the extreme points of the unit ball of Lip (S, d) appears to be quite difficult. The only results we know of this kind are in [10] and [11], and these are for S = [0, 1]. In both papers a proof is given that the unit ball of Lip [0, 1] is the norm-closed convex hull of its extreme points. This problem is also open for more general metric spaces.

Assuming S is compact and countable, (II) yields another previously unknown result. Again in [3] the extreme points of the unit ball of Lip  $(S, d)^*$  are shown to be of two types: one corresponding to a subset of  $S \cup [(S \times S) \sim \Delta]$  and the other a set Q "arising from" the Stone-Čech compactification of  $(S \times S) \sim \Delta$  (see [3] or [4]). It was shown in [4] that, in general, Q must be nonempty. It now follows from (III) that if S is compact and countably infinite, Q must be uncountable. The work of Sherbert [12] shows that the functionals in Q must be point derivations. However, a complete description of Q still appears difficult. We sum up this result in

### J. A. JOHNSON

PROPOSITION 3. If S is compact and countably infinite, the set Q of extreme points of the unit ball of  $\text{Lip}(S, d)^*$  that are point derivations is uncountable.

## References

1. R. Bonic, J. Frampton, and A. Tromba, *A-Manifolds*, J. Functional Analysis, **3** (1969), 310-320.

2. K. deLeeuw, Banach spaces of Lipschitz functions, Studia Math., 21 (1961/62), 55-66.

3. T. M. Jenkins, Banach spaces of Lipschitz functions on an abstract metric space, Thesis, Yale Univ., New Haven, Conn., 1967.

4. J. A. Johnson, Banach spaces of Lipschitz functions and vector-valued Lipschitz functions, Trans. Amer. Math. Soc., **148** (1970), 147-169.

5. \_\_\_\_, Lipschitz function spaces for arbitrary metrics, Bull. Amer. Math. Soc., 78 (1972), 702-705,

6. W. B. Johnson and H. P. Rosenthal, On w\*-basic sequences and their applications to the study of Banach spaces, Studia Math., 43 (1972), 77-92.

7. J. Lindenstrauss, *Extentions of compact operators*, Memoirs Amer. Math. Soc., **48** (1964).

8. J. Lindenstrauss and R. Phelps, Extreme point properties of convex bodies in reflexive Banach spaces, Israel J. Math., 6 (1968), 39-48.

9. J. Lindenstrauss and H. P. Rosenthal, The  $\mathcal{L}_p$  spaces, Israel J. Math., 7 (1969), 325-349.

10. N. V. Rao and A. K. Roy, Extreme Lipschitz functions, Math. Ann., 189 (1970), 26-46.

11. A. K. Roy, Extreme points and linear isometries of the Banach space of Lipschitz functions, Canad. J. Math., **20** (1968), 1150-1164.

12. D. R. Sherbert, The structure of ideals and point derivations in Banach algebras of Lipschitz functions, Trans. Amer. Math. Soc., **111** (1964), 240-272.

Received November 29, 1972 and in revised form March 8, 1973. This research was supported in part by a grant from the Oklahoma State University Research Foundation.

#### OKLAHOMA STATE UNIVERSITY

# PACIFIC JOURNAL OF MATHEMATICS

# EDITORS

RICHARD ARENS (Managing Editor) University of California Los Angeles, California 90024 J. DUGUNDJI\* Department of Mathematics University of Southern California Los Angeles, California 90007

D. GILBARG AND J. MILGRAM Stanford University Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH B. H. NEUMANN F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON \* \* \* \* AMERICAN MATHEMATICAL SOCIETY

AMERICAN MATHEMATICAL SOCIETY NAVAL WEAPONS CENTER

\* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Printed in Japan by Intarnational Academic Printing Co., Ltd., Tokyo, Japan

R. A. BEAUMONT University of Washington Seattle, Washington 98105

# Pacific Journal of Mathematics Vol. 51, No. 1 November, 1974

Zvi Arad, $\pi$ -homogeneity and $\pi'$ -closure of finite groups	1
Ivan Baggs, A connected Hausdorff space which is not contained in a maximal connected space	11
Eric Bedford, The Dirichlet problem for some overdetermined systems on the unit ball in $C^n$	19
R. H. Bing, Woodrow Wilson Bledsoe and R. Daniel Mauldin, Sets generated by rectangles	27
Carlo Cecchini and Alessandro Figà-Talamanca, <i>Projections of uniqueness for</i> $L^{p}(G)$	37
Gokulananda Das and Ram N. Mohapatra, <i>The non absolute Nörlund summability of</i> Fourier series	49
Frank Rimi DeMever. On separable polynomials over a commutative ring	57
Richard Detmer. Sets which are tame in arcs in $F^3$	67
William Frh Dietrich Ideals in convolution algebras on Abelian groups	75
Bryce I Elkins A Galois theory for linear topological rings	89
William Alan Feldman, A characterization of the topology of compact convergence on $C(X)$	100
Hillel Halkin Gershenson, A problem in compact Lie groups and framed	109
Cobolaism	121
Mamin L. Croonbarg. Stuigtly logal solutions of Displanting soughting	131
Infarvin J. Greenberg, Strictly local solutions of Diophantine equations	143
Domingo Antonio Horroro, Inner functione un den uniform topology	155
Johningo Antonio Henero, Inner junctions under uniform topology	107
Marvin Stanford Keeper, Oscillatory solutions and multi point boundary value	1//
functions for certain nth-order linear ordinary differential equations	187
John Cronan Kieffer, A simple proof of the Moy-Perez generalization of the Shannon-McMillan theorem	203
Joong Ho Kim, <i>Power invariant rings</i>	207
Gangaram S. Ladde and V. Lakshmikantham, On flow-invariant sets	215
Roger T. Lewis, Oscillation and nonoscillation criteria for some self-adjoint even order linear differential operators	221
Jürg Thomas Marti, On the existence of support points of solid convex sets	235
John Rowlay Martin, Determining knot types from diagrams of knots	241
James Jerome Metzger, Local ideals in a topological algebra of entire functions	251
K. C. O'Moore, Intrinsic artensions of prime rings	251
K. C. O Meara, Intrinsic extensions of prime rings	237
Statiley Foreda, A note on the continuity of best polynomial approximations	271
of mans	273
Fric Peter Smith The Garabedian function of an arbitrary compact set	275
Arne Stray, Pointwise bounded approximation by functions satisfying a side	207
	301
Jonn St. Clair Werth, Jr., <i>Maximal pure subgroups of torsion complete abelian</i> <i>p-groups</i>	307
Robert S. Wilson, On the structure of finite rings. II	317
Kari Ylinen, The multiplier algebra of a convolution measure algebra	327