

Pacific Journal of Mathematics

**POINTWISE BOUNDED APPROXIMATION BY FUNCTIONS
SATISFYING A SIDE CONDITION**

ARNE STRAY

POINTWISE BOUNDED APPROXIMATION BY FUNCTIONS SATISFYING A SIDE CONDITION

A. STRAY

In this paper necessary and sufficient conditions on a subset S of the unit disc D are given such that every bounded analytic function f on D is a pointwise limit of a sequence $\{f_n\}_{n=1}^\infty$ of uniformly continuous analytic functions on D bounded by the sup norm of f and in addition satisfying $\sup\{|f_n(z)|, z \in S\} \leq \sup\{|f(z)|, z \in S\}$ for all n .

Let $D = \{z: |z| < 1\}$ denote the open unit disc and $T = \{z: |z| = 1\}$ the unit circle. $H^\infty(D)$ denotes all bounded analytic functions on D and $A(D)$ consists of all uniformly continuous f in $H^\infty(D)$. For any $f \in H^\infty(D)$ and any subset S of D we put $\|f\|_S = \sup\{|f(z)|, z \in S\}$ and we set $\|f\| = \|f\|_D$.

A sequence $\{z_n\}_{n=1}^\infty$ in D converging to $z \in T$ converges *nontangentially* to z if for some constant λ we have $|z - z_n| \leq \lambda(1 - |z_n|)$ for all n . If $f \in H^\infty(D)$ then Fatou's theorem [2, page 34] tells us that f has a nontangential limit at almost every boundary point. Thus at almost every boundary point $\lim f(z_n)$ exists and is independent of the choice of sequence. If $f \in H^\infty(D)$ is known a.e. on T we recapture its values in D by the Cauchy or Poisson integral formula. We will therefore consider functions in $H^\infty(D)$ as defined in D and a.e. on T .

A relatively closed subset S of D is called a *Farrell set* if for each $f \in H^\infty(D)$ there are $f_n \in A$, $n = 1, 2, \dots$, converging pointwise to f on D with $\|f_n\| \leq \|f\|$ and such that $\|f_n\|_S \leq \|f\|_S$. This concept was introduced to us by Professor L. A. Rubel who also raised the question of describing such sets. The object here is to characterize Farrell sets in terms of their cluster points on T . The author is very grateful to Dr. A. M. Davie for valuable conversations on this subject.

First we observe that if $rz \in S$ whenever $0 < r < 1$ and $z \in S$, then S is a Farrell set. Indeed, letting $f_r(z) = f(rz)$ ($0 < r < 1$), we have: $f_r(z) \rightarrow f(z)$ as $r \rightarrow 1$.

On the other hand, let $S = \{z_n\}_{n=1}^\infty$ where $\sum_n (1 - |z_n|) < \infty$ and assume that the set of cluster points of S on T has positive linear measure. Then there are $f \in H^\infty(D)$ with $f = 0$ on S , but $f \neq 0$, while if $f \in A(D)$ and $f = 0$ on S we must have $f \equiv 0$. The set of cluster points of S on T which are not nontangential limits of sequences from S is here too large. In fact we will prove the following.

THEOREM. *A relatively closed subset S of the open unit disc D is*

a Farrell set if and only if the set S_1 of cluster points of S on T which are nontangential limits of sequences from S has zero linear measure.

Proof. First assume S_1 has positive linear measure. Using ideas from [1] we construct a function $f \in H^\infty(D)$ which shows that S is not a Farrell set. If $t > 0$ and $z \in T$ we define

$$\Delta(z, t) = \{w: 0 < 1 - |w| \leq t, |z - w| \leq 2(1 - |w|)\}.$$

For $t > 0$ we also define $E_t = \{z \in \bar{S} \cap T: \Delta(z, t) \cap S = \emptyset\}$.

Since $S_1 \subseteq \bigcup_t E_t$ has positive measure, we can find for some t_0 a set $E = E_{t_0}$ of positive measure.

Let u be the harmonic extension to D of the function on T which is zero on E and -1 on $T \setminus E$. Define $f = \exp[u + iv]$ where v is a harmonic conjugate to u in D . Then $f \in H^\infty(D)$ and by the theorem of Fatou we have $|f| = 1$ a.e. on E and $|f| = e^{-1}$ a.e. on $T \setminus E$. In particular $\|f\| = 1$.

We claim that $\|f\|_S < 1$. Assume this proved. Let $\{f_n\}_{n=1}^\infty \subset A(D)$, $\|f_n\| \leq 1$ and assume $f_n(z) \rightarrow f(z)$ $z \in D$. We prove that $\|f_n\|_S \rightarrow 1$ which in turn will show that S is not a Farrell set.

Given $\varepsilon > 0$ it follows from Fatou's theorem that there exists $z \in D$ such that

$$(1) \quad |f(z)| > 1 - \varepsilon$$

$$(2) \quad m_z(E) > 1 - \varepsilon.$$

Here m_z denotes harmonic measure on T w.r.t. z .

There exists a number N such that

$$(3) \quad |f_n(z) - f(z)| < \varepsilon \quad \text{if } n > N.$$

By (1) and (3) we have $|f_n(z)| > 1 - 2\varepsilon$ if $n > N$.

But on the other hand we have

$$|f_n(z)| \leq \int_E |f_n| dm_z + \int_{T \setminus E} |f_n| dm_z \leq \|f_n\|_E + \varepsilon$$

so that $1 \geq \|f_n\|_S \geq \|f_n\|_E \geq 1 - 3\varepsilon$ if $n \geq N$.

It remains to prove that $\|f\|_S < 1$. Choose $z \in S$ such that $|z| > 1 - t_0$.

Define $J_z = \{w \in T: |w - z| < 2(1 - |z|)\}$. By a well-known estimate of harmonic measure there exists a constant $c \in (0, 1)$ depending only on t_0 such that

$$(4) \quad m_z(J_z) \geq c.$$

Since $z \in S$ we have $J_z \cap E = \emptyset$.

By (4) we therefore have since $|f| \leq 1$ a.e. on E and $|f| \leq e^{-1}$ a.e. on $T \setminus E$:

$$\begin{aligned} |f(z)| &\leq \int_E |f| dm_z + \int_{T \setminus E} |f| dm_z \leq m_z(E) + \frac{1}{e} m_z(T \setminus E) \\ &\leq 1 - c + ce^{-1} < 1 - c2^{-1} \quad \text{if } z \in S. \end{aligned}$$

Since clearly $\sup \{|f(z)| : |z| \leq 1 - t_0\} < 1$, the claim is proved.

Suppose now S_1 has zero linear measure.

We first consider the special case where $B = \bar{S} \cap T$ itself has zero measure.

Given $f \in H^\infty(D)$ we first choose functions g_n continuous on $D \cup T \setminus B$ and analytic in D such that $\|g_n\|_D \leq \|f\|$, $\|g_n\|_S \leq \|f\|_S$ and $g_n(z) \rightarrow f(z)$ if $z \in D$. (See [3] § 1.)

We then choose $\{a_n\}_{n=1}^\infty \subset A(D)$ converging pointwise to 1 on D such that $\|a_n\| = 1$ and $a_n = 0$ on B . See [2], p. 80 for a construction of the a_n 's.

Define $f_n = 0$ on B and $f_n = a_n g_n$ in $D \setminus B$. Then $f_n \in A(D)$ and $\{f_n\}_{n=1}^\infty$ approximates f as required.

Finally we consider the case where B has positive linear measure.

Define $H(\theta, z) = (e^{i\theta} + z)/(e^{i\theta} - z)$ if $-\pi < \theta \leq \pi$ and $z \in D$.

Choose $f \in H^\infty(D)$ with $\|f\| = 1$. Let $\eta = \|f\|_S$. If $\eta = 0$ it follows from Fatou's theorem and that S_1 has zero linear measure, that $|f| = 0$ a.e. on B which is impossible since $f \neq 0$ and B has positive measure. Hence $\eta > 0$.

We can factorize f as $f = I \cdot F$ where I and F denotes the "inner" and "outer" part of f . [[2], Ch. 5.]

Here F is given by

$$F(z) = \exp \left[\frac{1}{2\pi} \int_{-\pi}^{\pi} H(\theta, z) u(\theta) d\theta \right]$$

and it follows from the hypothesis that $u \leq 0$ a.e. on T and $u \leq \log \eta$ a.e. on B .

Fix a positive integer n . Let u_1 be a continuously differentiable function on T and $V \supset B$ a set open in T such that $u_1 \leq 0$, $u_1 \leq \log \eta$ on V and with the following properties:

$$(5) \quad \int_{-\pi}^{\pi} |u(\theta) - u_1(\theta)| d\theta < \frac{1}{n}$$

$$(6) \quad \int_{T \setminus V} |H(\theta, z)(u(\theta) - u_1(\theta))| d\theta < \frac{1}{n} \text{ for all } z \in S.$$

To obtain u_1 on V one first defines u_1 in a neighborhood of B such that $\int_B |u(\theta) - u_1(\theta)| d\theta < 1/2n$. This inequality will still hold if B is

replaced by some open set V containing it if linear measure of $V \setminus B$ is small enough. But then it is easy to extend u_1 to T so that (5) and (6) hold.

Since $u_1 \leq \log \eta$ on V and $u_1 \leq 0$ there is a compact subset K of S such that

$$(7) \quad \int u_1 dm_z \leq \log \left(\eta + \frac{1}{n} \right) \quad \text{if } z \in S \setminus K.$$

Let $A = \{z \in V: u(z) \leq u_1(z)\}$ and M be a compact subset of A . We choose a continuously differentiable function $a \leq 0$ such that $\text{supp } a$ (the support of a) is contained in V and we have

$$(8) \quad \left| \int_V H(\theta, z)(u_1 + a - u)(\theta) d\theta \right| < \frac{1}{n} \quad \text{if } z \in K.$$

If a approximates $u - u_1$ sufficiently well on M and if the linear measures of $A \setminus M$ and $\text{supp } a \setminus M$ are sufficiently small we can obtain that (5) and (6) still hold if we replace u_1 by $(u_1 + a)$ there.

We define $g_n(z) = \exp \left[1/2\pi \int_{-\pi}^{\pi} (H(\theta, z)(u_1 + a)(\theta) d\theta) \right]$, $z \in D$.

Since $u_1 + a$ is smooth $g_n \in A(D)$ and $\|g_n\| \leq 1$ since $u_1 + a \leq 0$.

By (5), $g_n(z) \rightarrow f(z)$ if $z \in D$. When $z \in S \setminus K$, (7) implies that $|g_n(z)| \leq \eta + 1/n$ and if $z \in K$ we get from (6) and (8) that $|g_n(z)| \leq \eta e^{1/n}$.

Define now $h_n = (\eta/(\eta + 3/n))g_n$ and $B_n(z) = I(z(1 - n^{-1}))$ if $z \in D$ and $n = 1, 2, \dots$.

The sequence $f_n = B_n h_n n = 1, 2, \dots$ approximates f as required.

We have completed the proof that S is a Farrell set if S_1 has zero linear measure and the proof also shows how to construct the functions $\{f_n\}$ given f such that they satisfy the requirements given in the definition of a Farrell set.

If S_1 has zero linear measure and B has positive linear measure there is a proof based on functional analysis showing that S is a Farrell set. This proof is due to Dr. A. M. Davie. Since the proof is short and rather different from the one given above we would like to include it here.

So we assume $f \in H^\infty(D)$, $|f| = 1$ and $\|f\|_S = \eta > 0$.

Let $N = \{f \in C(\bar{D}) \mid \|f\|_S \leq \eta \text{ and } \|f\| \leq 1\}$. We have to show that f is in the closure of $N \cap A(D)$ in the topology of uniform convergence on compact subsets of D .

By the separation theorem and Riez representation theorem it is sufficient to prove that $|\mu(f)| \leq 1$ whenever μ is a regular complex Borel measure with compact support in D such that $|\mu(h)| \leq 1$ for all $h \in N \cap A(D)$. ($N \cap A(D)$ is a convex set in the space of all continuous functions in D , with the topology of uniform convergence on compact subsets of D and the dual space is the space of regular

complex Borel measures with compact support in D .)

N is the unit ball w.r.t. some norm on $C(\bar{D})$ which is equivalent to sup norm on \bar{D} since $\eta > 0$.

Hence we can extend the functional $g \rightarrow \mu(g)$ from $A(D)$ to $C(\bar{D})$ and represent it by a measure ν on \bar{D} such that $|\nu(g)| \leq 1$ for all $g \in N$.

But the last fact implies

$$(I): |\nu|(\bar{D} \setminus \bar{S}) + \eta |\nu| \bar{S} \leq 1$$

where $|\nu|$ denotes the total variation of ν .

We claim that $\nu(E) = 0$ if $E \subset T$ has zero linear measure. To see this let $K \subset E$ be compact. Choose $\{a_n\}_{n=1}^{\infty} \subset A(D)$ such that $\|a_n\| = 1$, $a_n = 0$ on K , and $a_n \rightarrow 1$ on $\bar{D} \setminus K$. (The sequence $\{a_n\}$ mentioned in the previous proof will do.) Then $0 = (\mu - \nu)(1 - a_n) \rightarrow (\mu - \nu)(K) = \nu(K)$ by dominated convergence.

This means that f is defined a.e. ν and if $\{f_n\} \subset A(D)$ converges pointwise to f on D and a.e. to f on T such that $\|f_n\| \leq \|f\|$ we again have by dominated convergence that $0 = \mu(f) - \nu(f)$. But $|f| \leq 1$ a.e. ν and $|f| \leq \eta$ a.e. ν on S so $|\mu(f)| = |\nu(f)| \leq 1$ by (I).

REFERENCES

1. L. Brown, A. Shields, and K. Zeller, *On absolutely convergent exponential sums*, Trans. Amer. Math. Soc., **96** (1960), 162-183.
2. K. Hoffman, *Banach Spaces of Analytic Functions*, Prentice-Hall, Inc., Englewood Cliffs, N.J. 1962
3. A. Stray, *Approximation and interpolation*, Pacific. J. Math., **40** (1972), 463-475.

Received October 27, 1972 and in revised form February 20, 1973.

AGDER DISTRIKSHOGSKOLE
 POSTBOKS 332
 4600 KRISTIANSAND S.
 NORWAY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Pacific Journal of Mathematics

Vol. 51, No. 1

November, 1974

Zvi Arad, π -homogeneity and π' -closure of finite groups	1
Ivan Baggs, A connected Hausdorff space which is not contained in a maximal connected space	11
Eric Bedford, The Dirichlet problem for some overdetermined systems on the unit ball in C^n	19
R. H. Bing, Woodrow Wilson Bledsoe and R. Daniel Mauldin, Sets generated by rectangles	27
Carlo Cecchini and Alessandro Figà-Talamanca, Projections of uniqueness for $L^p(G)$	37
Gokulananda Das and Ram N. Mohapatra, The non absolute Nörlund summability of Fourier series	49
Frank Rimi DeMeyer, On separable polynomials over a commutative ring	57
Richard Detmer, Sets which are tame in arcs in E^3	67
William Erb Dietrich, Ideals in convolution algebras on Abelian groups	75
Bryce L. Elkins, A Galois theory for linear topological rings	89
William Alan Feldman, A characterization of the topology of compact convergence on $C(X)$	109
Hillel Halkin Gershenson, A problem in compact Lie groups and framed cobordism	121
Samuel R. Gordon, Associators in simple algebras	131
Marvin J. Greenberg, Strictly local solutions of Diophantine equations	143
Jon Craig Helton, Product integrals and inverses in normed rings	155
Domingo Antonio Herrero, Inner functions under uniform topology	167
Jerry Alan Johnson, Lipschitz spaces	177
Marvin Stanford Keener, Oscillatory solutions and multi-point boundary value functions for certain n th-order linear ordinary differential equations	187
John Cronan Kieffer, A simple proof of the Moy-Perez generalization of the Shannon-McMillan theorem	203
Joong Ho Kim, Power invariant rings	207
Gangaram S. Ladde and V. Lakshmikantham, On flow-invariant sets	215
Roger T. Lewis, Oscillation and nonoscillation criteria for some self-adjoint even order linear differential operators	221
Jürg Thomas Marti, On the existence of support points of solid convex sets	235
John Rowlay Martin, Determining knot types from diagrams of knots	241
James Jerome Metzger, Local ideals in a topological algebra of entire functions characterized by a non-radial rate of growth	251
K. C. O'Meara, Intrinsic extensions of prime rings	257
Stanley Poreda, A note on the continuity of best polynomial approximations	271
Robert John Sacker, Asymptotic approach to periodic orbits and local prolongations of maps	273
Eric Peter Smith, The Garabedian function of an arbitrary compact set	289
Arne Stray, Pointwise bounded approximation by functions satisfying a side condition	301
John St. Clair Werth, Jr., Maximal pure subgroups of torsion complete abelian p -groups	307
Robert S. Wilson, On the structure of finite rings. II	317
Kari Ylinen, The multiplier algebra of a convolution measure algebra	327