Pacific Journal of Mathematics

REGULAR COMPLETIONS OF CAUCHY SPACES

DARRELL CONLEY KENT AND GARY DOUGLAS RICHARDSON

Vol. 51, No. 2

December 1974

REGULAR COMPLETIONS OF CAUCHY SPACES

D. C. KENT AND G. D. RICHARDSON

A uniform convergence space is a generalization of a uniform space. The set of all Cauchy filters of some uniform convergence space is called a Cauchy structure. We give necessary and sufficient conditions for the Cauchy structure of some totally bounded uniform convergence space to be precompact; i.e., have a regular completion. Also, it is shown that there is an isomorphism between the set of ordered equivalence classes of strict regular compactifications of a completely regular convergence space and the set of ordered precompact Cauchy structures inducing the given convergence structure.

Preliminaries. Kowalsky [5] has studied completions using only Cauchy filters, described axiomatically, and not necessarily those of a uniform convergence space. This has led others to the notion of a Cauchy space, which is described axiomatically in [2]. The reader is referred to [6], [7], and [8] for a discussion of completions of Cauchy spaces.

For basic definitions of convergence spaces and uniform convergence space, see [3] and [1]. A Hausdorff convergence space (S, q)is compatible with a uniform convergence space iff it satisfies the "Limitierungsaxiom": $\mathfrak{F} \cap \mathfrak{G}$ q-converges to x whenever \mathfrak{F} and \mathfrak{G} both q-converge to x. We will make the assumption that all convergence spaces in this paper satisfy this axiom. The closure operator in a convergence space (S, q) will be denoted by Γ_q . A Hausdorff convergence space (S, q) is called regular if it has the property that $\Gamma_q \mathfrak{F}$ (the filter generated by $\{\Gamma_q F | F \in \mathfrak{F}\}$) q-converges to x whenever \mathfrak{F} q-converges to x. The filter \dot{x} denotes the set of all subsets of S containing the set $\{x\}$. If filters \mathfrak{F} and \mathfrak{G} contain disjoint sets, we write " $\mathfrak{F} \vee \mathfrak{G} = 0$ ". The term "ultrafilter" will be abbreviated "u.f."; uniform convergence space will be abbreviated "u.c.s.".

A Cauchy structure \mathscr{C} on a set S is characterized axiomatically in [2] as follows: (1) $\dot{x} \in \mathscr{C}$ for each $x \in S$; (2) $\mathfrak{F} \in \mathscr{C}$ and \mathfrak{G} finer than \mathfrak{F} implies $\mathfrak{G} \in \mathscr{C}$; (3) $\mathfrak{F}, \mathfrak{G} \in \mathscr{C}$ and $\mathfrak{F} \vee \mathfrak{G} \neq 0$ implies $\mathfrak{F} \cap \mathfrak{G} \in \mathscr{C}$. The pair (S, \mathscr{C}) is called a Cauchy space. It should be pointed out that the Cauchy space axioms of [2] are stricter than those of [5] and [7].

A Cauchy space (S, \mathscr{C}) induces a convergence structure q in the following way: $\mathfrak{F} q$ -converges to x iff $\mathfrak{F} \cap \dot{x} \in \mathscr{C}$. Conversely, if (S, q) is a Hausdorff convergence space, then define the associated Cauchy structure \mathscr{C} on $S: \mathfrak{F} \in \mathscr{C}$ iff $\mathfrak{F} q$ -converges. Note that (S, \mathscr{C}) induces

q on S. A Cauchy space (S, \mathscr{C}) is called Hausdorff if the induced convergence space is Hausdorff, and complete if each Cauchy filter converges. We will assume that all spaces are Hausdorff unless otherwise indicated. The above describes a one-to-one correspondence between the convergence spaces and the complete Cauchy spaces. If \mathscr{C} is a Cauchy structure on S, then we often write C_q for C if q is the induced convergence structure. (S, \mathscr{C}_q) is called regular if $\Gamma_q \mathfrak{F} \in \mathscr{C}$ whenever $\mathfrak{F} \in \mathscr{C}$. This definition was suggested by the referee, and corresponds to the definition of regularity for u.s.c.'s given in [10] and [12].

Let (S, \mathscr{C}) be a Cauchy space and define (as in [1]): $\mathfrak{F} \sim \mathfrak{G}$ iff $\mathfrak{F} \cap \mathfrak{G} \in \mathscr{C}$. This equivalence relation partitions \mathscr{C} into equivalence classes of the form $[\mathfrak{F}] = \{\mathfrak{G} \in \mathscr{C} \mid \mathfrak{G} \sim \mathfrak{F}\}$. Let T be the set of equivalence classes, and let $j: S \to T$ denote the canonical mapping, i.e., $j(x) = [\dot{x}]$.

DEFINITION 1.1. (P, \mathcal{D}_r, f) is called a *completion* of the Cauchy space (S, \mathcal{C}) , if (P, \mathcal{D}) is complete, and f is a dense embedding from (S, \mathcal{C}) into (P, \mathcal{D}) . If in addition, whenever a filter \mathcal{F} r-converges to y in P, there is a filter \mathfrak{G} on fS which r-converges to y and such that $\Gamma_r \mathfrak{G} \leq \mathfrak{F}$, then (P, \mathcal{D}_r, f) is called a *strict completion*.

The notion of a completion of a u.c.s. is defined similarly. Wyler [11] has shown that each u.c.s. has a completion, with the universal property, and so each Cauchy space has a completion. If P denotes any convergence space property, then we say that a (strict) completion is a (strict) P completion if it possesses property P.

The next two definitions follow the terminology of [8]. Analogous definitions apply in the u.c.s. setting.

DEFINITION 1.2. A completion (P, \mathcal{D}, f) of the Cauchy space (S, \mathcal{C}) is said to be in *standard form* if P = T, f = j, and f converges to $[\mathcal{F}]$ for each $\mathcal{F} \in \mathcal{C}$.

DEFINITION 1.3. The completions $(P_i, \mathscr{D}_i, f_i)$, i = 1, 2, of the Cauchy space (S, \mathscr{C}) are said to be *equivalent* if there is an isomorphism g from (P_1, \mathscr{D}_1) onto (P_2, \mathscr{D}_2) such that $gf_1 = f_2$.

PROPOSITION 1.4 ([8]). Each Cauchy space (u.c.s.) completion is equivalent to exactly one in standard form.

Let (S, \mathscr{C}) be a Cauchy space and T, j defined as above. Let A be a subset of S; then define ΣA to be $\{[\mathfrak{F}] \in T \mid A \in \mathfrak{G} \text{ for some } \mathfrak{G} \in [\mathfrak{F}]\}$. If \mathfrak{G} is a filter on S, then $\Sigma \mathfrak{G}$ denotes the filter on T

generated by $\{\Sigma G \mid G \in \mathfrak{G}\}$. Further, the convergence structure p on T is defined as follows: \mathscr{H} p-converges to $[\mathfrak{F}]$ iff $\mathfrak{F} \geq \Sigma \mathfrak{G}$ for some $\mathfrak{G} \in [\mathfrak{F}]$. In general p is not Hausdorff, and so we use the notation \mathscr{C}_p only whenever p is Hausdorff. The following is straightforward to verify: if p is Hausdorff, then (T, \mathscr{C}_p, j) is a completion of (S, \mathscr{C}) iff (S, \mathscr{C}) is regular. Our next result follows immediately from the definitions.

PROPOSITION 1.5. Let (T, \mathcal{D}_s, j) be any completion of (S, \mathcal{C}_q) in standard form. The following are true.

(1) If $A \subset S$, then $\Gamma_s jA = \Sigma A$ and $\Gamma_g A = j^{-1} \Sigma A$.

(2) The completion is strict iff $s \ge p$.

COROLLARY 1.6. (T, \mathcal{C}_p, j) is the only possible candidate for a strict regular completion of (S, \mathcal{C}) in standard form. Moreover, (T, \mathcal{C}_p, j) has the universal property for regular Cauchy spaces.

Proof. The first part follows from (2) of Proposition 1.5 since the convergence structure induced by any regular completion on Tmust be coarser than p. The second part following from Theorem 4.11 of [7], since (T, \mathcal{C}_p) is the quotient space of the quasi-completion mentioned there.

Finally, we remark that if (T, \mathcal{C}_p, j) is a completion of (S, \mathcal{C}) , and if \mathscr{I} is a u.c.s. with Cauchy filters \mathscr{C} , then by Theorem 15 of [8], (S, \mathscr{I}) has a u.c.s. completion with Cauchy structure \mathcal{C}_p .

Almost topological completions. In [9] it is shown that a regular compactification (R, r, f) of a convergence space (S, q) is almost topological, which means that r and its topological modification, λr , coincide relative to the convergence of u.f.'s. The next theorem characterizes those Cauchy spaces which have almost topological completions. The proof of this theorem uses the following lemma proved in [4].

LEMMA 2.1. Let (S, q) be a convergence space, $A \subset S$, \mathfrak{F} an u.f. on S. $\Gamma_q A \in \mathfrak{F}$, then there is an u.f. \mathfrak{G} containing A such that $\mathfrak{F} \geq \Gamma_q \mathfrak{G}$.

THEOREM 2.2. The following conditions are equivalent for a regular Cauchy space (S, \mathcal{C}) .

- (1) (S, \mathcal{C}) has an almost topological regular completion.
- (2) If $\Sigma \mathfrak{F} \vee \Sigma \mathfrak{G} \neq 0$ for $\mathfrak{F} \in \mathscr{C}$, \mathfrak{G} an u.f. on S, then $\mathfrak{F} \cap \mathfrak{G} \in \mathscr{C}$.
- (3) (T, \mathcal{C}_{p}, j) is an almost topological regular completion.

Proof. (1) *implies* (2). Let (T, \mathcal{D}_r, j) be such a completion in standard form, and let \mathfrak{F} , \mathfrak{G} be as in the hypothesis of (2). Then $\Gamma_r j\mathfrak{F} \vee \Gamma_r j\mathfrak{G} \neq 0$, and so $[\mathfrak{F}]$ is an adherent point of the u.f. $j\mathfrak{G}$ in $(T, \lambda r)$. Thus $j\mathfrak{G} \lambda r$ -converges to $[\mathfrak{F}]$, and so by hypothesis *r*-converges to [F]. Hence $\mathfrak{F} \cap \mathfrak{G} \in \mathfrak{C}$, and (2) is satisfied.

(2) implies (3). First we show that if $\Sigma \mathfrak{F} \vee \Sigma \mathfrak{G} \neq 0$, for $\mathfrak{F}, \mathfrak{G} \in \mathscr{C}$, then $[\mathfrak{F}] = [\mathfrak{G}]$. Let \mathfrak{U} be an u.f. on T such that $\mathfrak{U} \geq \Sigma \mathfrak{F} \vee \Sigma \mathfrak{G}$. From Lemma 2.1 there is an u.f. \mathfrak{F} on S such that $\Sigma \mathfrak{F} \leq \mathfrak{U}$. Thus $\Sigma \mathfrak{F} \vee \Sigma \mathfrak{F} \neq 0$, and by condition (2) $[\mathfrak{F}] = [\mathfrak{F}]$. Similarly, $[\mathfrak{F}] = [\mathfrak{G}]$, and so $[\mathfrak{F}] = [\mathfrak{G}]$. Thus (T, p) is Hausdorff, and so (T, \mathscr{C}_p, j) is a completion of (S, \mathscr{C}) .

Two steps are needed to prove Γ_p is idempotent. First let $A \subset S$ and $[\mathfrak{F}] \in \Gamma_p^2 jA = \Gamma_p \Sigma A$. Then there is an u.f. \mathfrak{U} p-converging to $[\mathfrak{F}]$ such that $\Sigma A \in \mathfrak{U}$. Thus $\mathfrak{U} \geq \Sigma \mathfrak{G}$ for some $\mathfrak{G} \in [\mathfrak{F}]$. By Lemma 2.1, there is an u.f. \mathfrak{F} on A such that $\mathfrak{U} \geq \Sigma \mathfrak{F}$, and we have $\Sigma \mathfrak{G} \vee \Sigma \mathfrak{F} \neq 0$. By condition (2), $[\mathfrak{F}] = [\mathfrak{F}]$, and since $A \in \mathfrak{F}$, then $[\mathfrak{F}] \in \Sigma A$. Thus $\Gamma_p^2 jA = \Gamma_p jA$ whenever $A \subset S$.

Next let $B \subset T$ and $[\mathfrak{F}] \in \Gamma_p^* B$. Then there is an u.f. \mathfrak{U} p-converging to $[\mathfrak{F}]$ such that $\Gamma_p B \in \mathfrak{U}$. Thus $\mathfrak{U} \geq \Sigma \mathfrak{G}$ for some $\mathfrak{G} \in [\mathfrak{F}]$. Using Lemma 2.1 again, there is an u.f. \mathfrak{K} on B such that $\mathfrak{U} \geq \Gamma_p \mathfrak{K}$, and also an u.f. \mathfrak{G} on S such that $\mathfrak{K} \geq \Sigma \mathfrak{G}$. Thus $\mathfrak{U} \geq \Gamma_p \mathfrak{K} \geq \Gamma_p \mathfrak{L} \mathfrak{G} = \Sigma \mathfrak{G}$, and so $\Sigma \mathfrak{G} \vee \Sigma \mathfrak{G} \neq 0$, which implies that $[\mathfrak{F}] = [\mathfrak{G}]$. Since \mathfrak{K} p-converges to $[\mathfrak{G}]$ and $B \in \mathfrak{K}$, then $[\mathfrak{F}] \in \Gamma_p B$. Thus $\Gamma_p^* B = \Gamma_p B$, for $B \subset T$. If $\mathfrak{F} \in \mathfrak{C}$, then $\Gamma_p \Sigma \mathfrak{F} = \Gamma_p^* j \mathfrak{F} = \Gamma_p j \mathfrak{F} = \Sigma \mathfrak{F}$, and so (T, \mathfrak{C}_p, j) is a regular completion of (S, \mathfrak{C}) .

Finally, we show that p and λp coincide on u.f.'s. Let $[\mathfrak{F}] \in T$ and \mathfrak{U} an u.f. such that $\mathfrak{U} \geq \bigcap \{ \mathfrak{F} \mid \mathfrak{F} p$ -converges to $[\mathfrak{F}] \}$. The latter intersection is the *p*-neighborhood filter at $[\mathfrak{F}]$, and since Γ_p is idempotent, the λp -neighborhood filter at $[\mathfrak{F}]$. Note that $[\mathfrak{F}] \geq \Gamma_p \mathfrak{U} \vee \Sigma \mathfrak{F}$. From Lemma 2.1, there is an u.f. \mathfrak{G} on S such that $\mathfrak{U} \geq \Sigma \mathfrak{G}$, and so $\Gamma_p \mathfrak{U} \geq \Gamma_p \Sigma \mathfrak{G} = \Sigma \mathfrak{G}$. Thus $\Sigma \mathfrak{G} \vee \Sigma \mathfrak{F} \neq 0$, which implies $[\mathfrak{G}] = [\mathfrak{F}]$, and so \mathfrak{U} *p*-converges to $[\mathfrak{F}]$, which completes the proof.

PROPOSITION 2.3. If (T, \mathcal{C}_s, j) is any almost topological regular completion of (S, \mathcal{C}) in standard form, then $s \leq p$ and s = p on u.f.'s.

Proof. Clearly $s \leq p$. Let \mathfrak{U} be an u.f. which s-converges to $[\mathfrak{F}]$ in T. Then from Lemma 2.1, there is an u.f. \mathfrak{G} on S such that $\Gamma_s j \mathfrak{G} \leq \mathfrak{U}$. Thus the u.f. $j \mathfrak{G} \lambda s$ -converges to $[\mathfrak{F}]$, which implies by hypothesis that $j \mathfrak{G} s$ -converges to $[\mathfrak{F}]$. Hence $\mathfrak{G} \in [\mathfrak{F}]$, and from Proposition 1.5 $\Gamma_s j \mathfrak{G} = \Sigma \mathfrak{G}$, which implies that \mathfrak{U} p-converges to $[\mathfrak{F}]$, and so s = p on u.f.'s. Precompact Cauchy spaces. A Cauchy space (u.c.s.) is said to be *totally bounded* if every u.f. is Cauchy. A totally bounded Cauchy space with a regular completion will be termed *precompact*. From Theorem 2.2 we conclude the following.

PROPOSITION 3.1. A precompact Cauchy space is almost topological and has an almost topological regular completion.

Another characterization of precompact Cauchy spaces is given by Theorem 3.4. First, we need two preliminary results, the first of which is proved in [9].

LEMMA 3.2. A convergence space (S, q) is compact and regular iff (S, q) is almost topological and λq is a compact Hausdorff topology.

PROPOSITION 3.3. Let (S, \mathscr{I}) be a u.c.s. with a compact regular induced convergence structure q, and let \mathscr{U} be the filter of λq -neighborhoods of the diagonal \varDelta_s in $S \times S$. Then each element of \mathscr{I} is finer than \mathscr{U} .

Proof. It follows from Lemma 3.2 that \mathscr{U} is a Hausdorff uniformity. Let $\emptyset \in \mathscr{I}$, and assume $\mathscr{U} \leq \emptyset$. Then there is an u.f. \mathfrak{F} on $S \times S$ such that $\emptyset \leq \mathfrak{F}$ and $\mathscr{U} \leq \mathfrak{F}$. Let \mathfrak{F}_1 and \mathfrak{F}_2 be the first and second projections, respectively, of \mathfrak{F} onto S; then by the assumption of compactness, there are points x and y in S such that \mathfrak{F}_1 q-converges to x and \mathfrak{F}_2 q-converges to y. Since $\mathscr{U} \leq \mathfrak{F}, x$ and y must be distinct. But $(\mathfrak{F}_1 \times \mathfrak{F}_2) \vee \emptyset \neq 0$, and so $\dot{x} \times \dot{y} = (\dot{x} \times \mathfrak{F}_1) \circ \emptyset \circ (\mathfrak{F}_2 \times \dot{y}) \in \mathscr{I}$. This contradicts the fact that (S, q) is Hausdorff.

THEOREM 3.4. The Cauchy structure (S, \mathscr{C}) of a totally bounded u.c.s. (S, \mathscr{I}) is precompact iff the following conditions are satisfied. (1) $\mathscr{U} = \bigcap \{ \Phi \mid \Phi \in \mathscr{I} \}$ is a Hausdorff uniformity on S.

(1) $Z = \prod_{i=1}^{n} [\psi_{i} \psi_{i} \psi_{i} \psi_{i}]$ (2) (S, \mathscr{C}) is regular.

(3) If F and S are u.f.'s on S such that $\mathfrak{F} \times \mathfrak{S} \geq \mathbb{Z}$, then $\mathfrak{F} \times \mathfrak{S} \in \mathscr{I}$.

Proof. Assume the three conditions. Let (S', \mathscr{W}) denote the Hausdorff uniform completion of (S, \mathscr{U}) . For each \mathscr{U} -Cauchy filter \mathfrak{F} on S, let $[\mathfrak{F}]_{\mathscr{U}} = \{\mathfrak{G} \mid \mathfrak{G} \text{ is } \mathscr{U}$ -Cauchy and $\mathfrak{F} \times \mathfrak{G} \geq \mathscr{U}\}$. From condition (3) and the fact that (S, \mathfrak{F}) is totally bounded, it follows that an u.f. \mathfrak{G} is in $[\mathfrak{F}]$ (the \mathscr{C} -equivalence class) iff $\mathfrak{G} \in [\mathfrak{F}]_{\mathscr{U}}$. Since S' can be identified with $\{[\mathfrak{F}]_{\mathscr{U}} \mid \mathfrak{F} \text{ is an u.f. on } S\}$, we can identify S' with $T = \{[\mathfrak{F}] \mid \mathfrak{F} \in \mathscr{C}\}$. If r is the topology on T associated with \mathscr{W} , then $\Sigma A \subset \Gamma_r jA, A \subset S$. If $\mathfrak{F} \in \mathscr{C}$ and \mathfrak{G} is an u.f. on S such that

 $\Sigma \mathfrak{F} \vee \Sigma \mathfrak{G} \neq 0$, then $\Gamma_r j \mathfrak{F} \vee \Gamma_r j \mathfrak{G} \neq 0$, and so $[\mathfrak{F}]_{\mathfrak{X}} = [\mathfrak{G}]_{\mathfrak{X}}$. Hence $[\mathfrak{F}] = [\mathfrak{G}]$, or $\mathfrak{F} \cap \mathfrak{G} \in \mathfrak{C}$, and by Theorem 2.2, (S, \mathfrak{C}) is precompact.

Conversely, assume (S, \mathscr{C}) is precompact. From our previous results, (T, \mathscr{C}_p, j) is a regular completion of (S, \mathscr{C}) . From Theorem 15 of [8], there is a u.c.s. \mathscr{J} on T which has Cauchy filters \mathscr{C}_p and such that (T, \mathscr{J}, j) is a completion of (S, \mathscr{I}) . Note that \mathscr{J} induces the convergence structure p on T. Let \mathscr{W} be the uniformity on Tof λp -neighborhoods of the diagonal Δ_T . By Proposition 3.3, each $\psi \in \mathscr{J}$ is finer than \mathscr{W} . Let $\mathscr{U} = (j \times j)^{-1}(\mathscr{W})$; then \mathscr{U} is a uniformity and each $\varphi \in \mathscr{I}$ is finer than \mathscr{U} . Thus $\mathscr{U} \leq \bigcap {\varphi | \varphi \in \mathscr{I}}$. If \mathscr{U} is strictly coarser than $\bigcap \varphi$, then there is an u.f. \mathfrak{F} on $S \times S$ such that $\mathfrak{F} \geq \mathscr{U}$, but $\mathfrak{F} \not\geq \bigcap \varphi$. Let $\mathfrak{F}_1, \mathfrak{F}_2$ denote the projections of \mathfrak{F} ; then since $(j \times j)\mathfrak{F} \geq \mathscr{W}, j\mathfrak{F}_1$ and $j\mathfrak{F}_2$ converge to the same point in (T, p). Thus $\mathfrak{F}_1 \times \mathfrak{F}_2 \in \mathscr{I}$, and so $\mathfrak{F} \in \mathscr{I}$, which contradicts $\mathfrak{F} \not\geq \bigcap \varphi$. Hence $\mathscr{U} = \bigcap \varphi$ is a Hausdorff uniformity on S, and (1) follows.

Of course (2) is clear. If $\mathfrak{F}, \mathfrak{G}$ are u.f.'s on S such that $\mathfrak{F} \times \mathfrak{G} \geq \mathscr{U}$, then $j\mathfrak{F} \times j\mathfrak{G} \geq \mathscr{W}$, and so they λp -converge to the same point. Since (T, p) is almost topological, then $j\mathfrak{F}$ and $j\mathfrak{G}$ p-converge to the same point, and so $\mathfrak{F} \times \mathfrak{G} \in \mathscr{I}$, which implies (3).

Strict regular compactifications. One of the more significant results in uniform space theory is the existence of an isomorphism from the ordered set of equivalence classes of the Hausdorff compactifications of a completely regular topological space and the ordered set of compatible precompact uniformities.

(R, r, f) is said to be a strict compactification of the convergence space (S, q), if f is a dense embedding, (R, r) is a compact convergence space, and if \mathfrak{F} r-converges to $y \in R$, then there is a filter \mathfrak{G} on fSwhich r-converges to y and $\Gamma_r \mathfrak{G} \leq \mathfrak{F}$. We define equivalence classes of compactifications of (S, q), and an ordering among the classes, analogous to the topological setting. Also if (S, \mathfrak{C}_1) , (S, \mathfrak{C}_2) are two Cauchy spaces, then $\mathfrak{C}_1 \geq \mathfrak{C}_2$ is defined to be $\mathfrak{C}_1 \subset \mathfrak{C}_2$.

A convergence space will be called *completely regular* if it has a strict regular compactification.

PROPOSITION 4.1. A convergence space (S, q) is completely regular iff it is almost topological and λq is a completely regular topology.

This result is essentially proved in [9], but the following two points need to be added. The compactification in [9] is in fact a strict regular compactification. The "Limitierungsaxiom" was not assumed in [9], but causes no difficulty if imposed.

THEOREM 4.2. If (S, q) is a completely regular convergence space,

then the ordered set of equivalence classes of strict regular compactifications of (S, q) is isomorphic to the ordered set of precompact Cauchy structures on S which induce q.

Proof. Let (S, \mathcal{C}) be a precompact Cauchy space which induces q on S; then (T, \mathcal{C}_p, j) is a strict regular completion of (S, \mathcal{C}) . Thus (T, p, j) is a strict regular compactification of (S, q). Define $\gamma(S, \mathcal{C}) = (T, p, j)$. We show that γ is an isomorphism.

Suppose \mathscr{C}_1 and \mathscr{C}_2 are distinct Cauchy structures on S; with no loss of generality assume $\mathfrak{F} \in \mathscr{C}_1 - \mathscr{C}_2$. We claim that $\gamma(S, \mathscr{C}_1) =$ (T_1, p_1, j_1) is not equivalent to $\gamma(S, \mathscr{C}_2) = (T_2, p_2, j_2)$. Assume, on the contrary, that there is a homeomorphism $f: (T_1, p_1) \to (T_2, p_2)$ such that $fj_1 = j_2$. Then $j_1\mathfrak{F}$ p_1 -converges to $[\mathfrak{F}]_1 \in T_1$, and so $fj_1\mathfrak{F} = j_2\mathfrak{F}$ p_2 -converges to an element in T_2 . This can occur only if $\mathfrak{F} \in \mathscr{C}_2$, which contradicts the choice of \mathfrak{F} , and it follows that γ is injective.

Next to show γ is onto. Let (R, r, f) be any strict regular compactification of (S, q). Let \mathscr{C} be the set of all filters \mathfrak{F} on S such that $f\mathfrak{F}$ r-converges in R. By a straightforward argument, it can be shown that \mathscr{C} satisfies the Cauchy space axioms, and is also totally bounded and induces q. Since (R, \mathscr{C}_r, f) is strict regular completion of (S, \mathscr{C}) , then by Theorem 2.2 (T, \mathscr{C}_p, j) is also a strict regular completion of (S, \mathscr{C}) . By Corollary 1.6, (P, p, j) and (R, r, f) are equivalent. Hence γ is subjective.

Finally to show that γ and γ^{-1} are order preserving. Suppose $\mathscr{C}_1 \geq \mathscr{C}_2$, i.e., $\mathscr{C}_1 \subset \mathscr{C}_2$. Let $\gamma(S, \mathscr{C}_i) = (T_i, p_i, j_i)$, i = 1, 2. It is straightforward to check that if $f: T_1 \to T_2$ such that $f([\mathfrak{F}]_1) = [\mathfrak{F}]_2$, where $[\mathfrak{F}]_i$ is the equivalence class in T_i of $\mathfrak{F} \in \mathscr{C}_i$, then $f(\mathfrak{L}_1 A) \subset \mathfrak{L}_2 A$. Thus $f(\mathfrak{L}_1 \mathfrak{F}) \geq \mathfrak{L}_2 \mathfrak{F}$, and so f is continuous. It follows that $(T_1, p_1, j_1) \geq (T_2, p_2, j_2)$. The proof that γ^{-1} is order preserving is straightforward, and the theorem follows.

We conclude with the following remarks concerning Theorem 4.2. In either of the ordered sets of Theorem 4.2, each nonempty subset has a supremum; the finest precompact Cauchy structure on S which induces q corresponds to the Stone-Čech compactification (S, q).

Acknowledgment. The authors are indebted to the referee for a number of corrections and improvements in the original manuscript.

References

2. H. H. Keller, Die Limes-Uniformisierbarkeit der Limesräume, Math. Ann., 176 (1968), 334-341.

3. D. C. Kent, Convergence quotient maps, Fund Math., 65 (1969), 197-205.

^{1.} C. H. Cook and H. R. Fischer, Uniform convergence structures, Math. Ann., 173 (1967), 290-306.

4. D. C. Kent and G. D. Richardson, *The decomposition series of a convergence space*, Czech. Math. J., (to appear).

H-J. Kowalsky, Limesräume und Komplettierung, Math. Nachr., 12 (1954), 301-340.
F. R. Miller, The Approximation of Topologies in Functional Analysis, Doctoral Dissertation, University of Massachusetts, 1968.

7. J. F. Ramaley and O. Wyler, Cauchy spaces II. Regular completions and compactifications, Math. Ann., 187 (1970), 187-199.

8. E. E. Reed, Completions of uniform convergence spaces, Math. Ann., **194** (1971), 83-108.

9. G. D. Richardson and D. C. Kent, *Regular compactifications of convergence spaces*, Proc. Amer. Math. Soc., **31** (1972), 571-573.

10. B. Sjöberg. Über die Fortsetzbarkeit gleichmässig stetiger Abbildungen in Limeräumen, Comm. Phys.-Math., **40** (1970), 41-46.

11. O. Wyler, Ein Komplettierungsfunktor für uniforme Limesräume, Math. Nachr., 46 (1970), 1-12.

12. _____, Filter Space Monads, Regularity, Completions, Technical Report 73-1, Department of Mathematics, Carnegie-Mellon University, 1973.

Received November 29, 1972, and in revised form August 22, 1973.

WASHINGTON STATE UNIVERSITY AND EAST CAROLINA UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor) University of California Los Angeles, California 90024 J. Dugundji*

Department of Mathematics University of Southern California Los Angeles, California 90007

D. GILBARG AND J. MILGRAM Stanford University Stanford, California 94305

University of Washington Seattle, Washington 98105

R. A. BEAUMONT

ASSOCIATE EDITORS

E. F. BECKENBACH B. H. NEUMANN

F. WOLF

K. Yoshida

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON * * * * AMERICAN MATHEMATICAL SOCIETY NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The Pacific of Journal Mathematics is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

> Copyright © 1973 by Pacific Journal of Mathematics Manufactured and first issued in Japan

Pacific Journal of Mathematics Vol. 51, No. 2 December, 1974

Robert F. V. Anderson, Laplace transform methods in multivariate spectral	220
theory	339
William George Bade, Two properties of the Sorgenfrey plane	349
John Robert Baxter and Rafael Van Severen Chacon, <i>Functionals on continuous</i> <i>functions</i>	355
Phillip Wayne Bean, Helly and Radon-type theorems in interval convexity	363
James Robert Boone On k-auotient mannings	369
Ronald P Brown <i>Extended prime</i> spots and auadratic forms	379
William Hugh Cornish, Crawley's completion of a conditionally upper continuous lattice	397
Robert S Cunningham <i>On finite left localizations</i>	407
Robert Jay Daverman, Approximating polyhedra in codimension one spheres embedded in s ⁿ by tame polyhedra	417
Burton I Fein Minimal splitting fields for group representations	427
Peter Fletcher and Robert Allen McCoy, Conditions under which a connected representable space is locally connected	433
Ionathan Samuel Golan, Tonologies on the torsion-theoretic spectrum of a	155
noncommutative ring	439
Manfred Gordon and Edward Martin Wilkinson. <i>Determinants of Petrie</i>	
<i>matrices</i>	451
Alfred Peter Hallstrom, A counterexample to a conjecture on an integral condition	
for determining peak points (counterexample concerning peak points)	455
E. R. Heal and Michael Windham, Finitely generated F-algebras with applications	
to Stein manifolds	459
Denton Elwood Hewgill, On the eigenvalues of a second order elliptic operator in an unbounded domain	467
Charles Royal Johnson, <i>The Hadamard product of A and A</i> *	477
Darrell Conley Kent and Gary Douglas Richardson, <i>Regular completions of Cauchy</i>	483
Alan Greenwell Law and Ann I. McKerracher. Sharpened polynomial	405
approximation	491
Bruce Stephen Lund. Subalgebras of finite codimension in the algebra of analytic	
functions on a Riemann surface	495
Robert Wilmer Miller. TTF classes and quasi-generators	499
Roberta Mura and Akbar H. Rhemtulla, Solvable groups in which every maximal	
partial order is isolated	509
Isaac Namioka, Separate continuity and joint continuity	515
Edgar Andrews Rutter, A characterization of OF – 3 rings	533
Alan Saleski, Entropy of self-homeomorphisms of statistical pseudo-metric spaces	537
Ryōtarō Satō An Abel-maximal ergodic theorem for semi-groups	543
H A Seid Cyclic multiplication operators on L -spaces	549
H B Skerry On matrix mans of entire sequences	563
John Brendan Sullivan, A proof of the finite generation of invariants of a normal	571
subgroup	571
solvable, VI	573
Ronson Joseph Warne, Generalized $\omega - \mathcal{L}$ -unipotent bisimple semigroups	631
Toshihiko Yamada, On a splitting field of representations of a finite group	649